

# Fungi aliquot saxonici novi lecti a cl. W. Krieger.

(Contributio Va ad Floram Mycol. Saxoniam)

auctore J. Bresadola.

*Diplodia Aristolochiae* Bres. et Krieg. n. sp.

Peritheciis subcutaneis, pustulatim exsurgentibus, dense gregariis, sphaeroideis,  $\frac{1}{4}$  mm latis, contextu parenchymatico, fuligineo; sporulis fuscis, 1-septatis, subconstrictis, ellipticis vel clavatis,  $25-29 = 10-12 \mu$ ; basidiis cylindraceis,  $10-12 = 4 \mu$ .

Hab. in ramis Aristolochiae Siphonis in hortis „Königstein“.

*Septoria Pyrethri* Bres. et Krieg. n. sp.

Epiphylla; maculis sparsis, irregularibus, ex isabellino fuscis, vix marginatis; peritheciis sparsis, nigris, globoso-lenticularibus, poro prominulo pertusis, contextu parenchymatico; sporulis cylindraccis, apicibus attenuatis, obtusis, decimam 2-pluriseptatis, hyalinis,  $50-75 = 2-3 \mu$ .

Hab. in foliis Pyrethri Parthenii „Königstein“.

*Phlyctena Jasiones* Bres. n. sp.

Peritheciis sparsis vel gregariis, hypophyllis, centro nigris contextu parenchymatico, margine pallidis cellulis tenuissimis, oblongatis, conflato, obovatis,  $85-100 \mu$ ; sporulis oblongatis,  $18-24 = 1-1\frac{1}{2} \mu$ .

Hab. in foliis exsiccatis Jasiones montanac prope „Königstein“.

Obs. Species haec ob perithecia tenuiora, ambitu subincompleto ad Phlyctaenam vagabundam accedit, a qua tamen optime distincta.

*Gleosporium Spiraeae* Bres. n. sp.

Maculis fuscidulo-umbrinis, amphigenis, marginalibus indeterminatis, centralibus obovatis, vix marginatis; acervulis gregariis, punctiformibus, amphigenis, hypodermicis, pustulatim prominulis,  $90-100 \mu$  latis; conidiis oblongatis, saepc inaequilateralibus,  $10-14 = 3-4 \mu$ ; basidiis simplicibus, cylindraceis,  $12-14 = 2-3 \mu$ .

Hab. in foliis Spiraeae opulifoliac prope „Prebischtbor“ Saxoniac.

*Myxosporium Mali* Bres. n. sp.

Acervulis primitus epidermide velatis, dein liberis, rotundatis vel oblongatis,  $\frac{1}{2}$ —1 mm latis, pallidis, siccitate nigris; conidiis oblongo-ovatis, hyalinis, 8—11 = 3—4  $\mu$ .

Hab. in ramis aridis Piri Mali prope „Königstein“.

*Ramularia Erodii* Bres. n. sp.

Maculis amphigenis, parvis, irregularibus, haud marginatis; caespitulis dense gregariis, albis, amphigenis; hyphis cylindraceo-tortuosis, septatis, simplicibus, 45—60 = 4  $\mu$ ; conidiis oblongis, 1—4 septatis, utrinque attenuatis, 24—55 = 2—3  $\mu$ .

Hab. in foliis Erodii cicutari prope „Königstein“.

*Macrosporium Kriegerianum* Bres. n. sp.

Caespitulis hypophyllis, dense gregariis, minimis, fuligineis; hyphis fasciculatis, interdum ramosis, septatis, olivaceis, 75—80 = 4—7  $\mu$ ; conidiis variantibus, 2—5-septato-muriformibus, 36—51 = 24—30  $\mu$ , olivaceo-fuligineis.

Hab. in foliis Mulgedii alpini „Arber im Böhmer-Walde“.

# ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Hedwigia](#)

Jahr/Year: 1897

Band/Volume: [36\\_1897](#)

Autor(en)/Author(s): Bresadola J.

Artikel/Article: [Fungi aliquot saxonici novi lecti a cl. W. Krieger. 381-382](#)