

## Investigations on secondary metabolites of lichens and lichen mycobionts in culture: production and biological activity

**Armin HAGER, Georg BRUNAUER &  
Elfie STOCKER-WÖRGÖTTER**

Lichen fungi produce a great variety of secondary metabolites, including aliphatic, cycloaliphatic, aromatic and terpenic substances. Lichen secondary metabolites, especially polymalonyl derived polyketides, have been screened for manifold biological activities, e.g. antiviral, antibiotic, antitumor and antiherbivore issues (HUNECK 1999, 2001; DAYAN & ROMAGNI 2001; MÜLLER 2001). They can also inhibit growth of higher plants. Culture experiments in recent years have shown that optimised culture conditions result in the formation of fats and fatty acids instead of secondary compounds. A sophisticated interaction of environmental factors, present nutrients and morphological complexity of the lichen influence the induction of secondary metabolite formation and therefore the production of particular lichen substances in culture. Mycobionts of various lichen species (e.g. *Heterodea muelleri*, *Lecanora rupicola*, *Xanthoria elegans*, *Cladonia leporina*) were isolated and cultured by means of single spore (AHMADJIAN 1973) and tissue culture (YAMAMOTO 1990) methods. Then, well grown subcultures were exposed to stress factors, e.g. shortwave UV and cold temperature. Lichen thalli and cultured mycobionts were analysed by means of HPLC and TLC to detect the secondary chemistry profile. DNA-analyses of the ITS region were performed to assure mycobiont authenticity. Exposing mycobiont cultures to the stress parameters induced the production of secondary compounds in higher quantities but in some cases different ratios. A further series of experiments focused on the effect of diffractaic and barbatic acids on the growth of the symbiotic photobiont *Trebouxia jamesii*; usnic acid was similarly tested. Pure substances were obtained from mycobiont cultures by performing a "preparative TLC". A determined quantity of algae was incubated on BBM plates containing three different concentrations of the pure lichen metabolites. The growth of the photobionts was monitored over a period of one month to evaluate the impact of each substance on the cultured algae. While diffractaic and usnic acid

had no noticeable effect; barbatic acid strongly inhibited algal growth and resulted in cell death.

**Literatur:**

- AHMADJIAN, V., 1973: Methods of isolating lichen symbionts and thalli. In: AHMADJIAN, V. & HALE, M.E. (eds.). *The Lichens*. Academic Press, New York: 653-659.
- DAYAN, F.E. & ROMAGNI, J.G., 2001: Lichens as a potential source of pesticides. *Pestic Outlook* **12**: 229-232.
- HUNECK, S., 1999: The significance of lichens and their metabolites. *Naturwissenschaften* **86**: 559-570.
- HUNECK, S., 2001: New Results on the Chemistry of Lichen Substances. In HERZ, W., FALK, H., KIRBY, G.W. & MOORE, R.E. (eds.). *Progress in the Chemistry of Organic Natural Products* **81**. Springer Verlag, Wien, New York.
- MÜLLER, K., 2001: Pharmaceutically relevant metabolites from lichens. *Applied Microbiology and Biotechnology* **56**: 9-16.
- YAMAMOTO, Y., 1990: Studies of cell aggregates and the production of natural pigments in plant cell culture. Nippon Paint Public.

**address:**

Armin HAGER, Georg BRUNAUER & Elfie STOCKER-WÖRGÖTTER  
Department of Organismic Biology  
University of Salzburg  
Hellbrunnerstr. 34  
A-5020 Salzburg,

**email:**

armin.hager@sbg.ac.at

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sauteria-Schriftenreihe f. systematische Botanik, Floristik u. Geobotanik](#)

Jahr/Year: 2008

Band/Volume: [16](#)

Autor(en)/Author(s): Hager Armin, Brunauer Georg, Stocker-Wörgötter Elfie

Artikel/Article: [Investigations on secondary metabolites of lichens and lichen mycobionts in culture: production and biological activity 323-324](#)