

## Diagnoses Fungorum novorum Agaricalium II\*).

Rolf Singer.

Instituto Miguel Lillo, Tucumán.

### **Polyporus princeps** Sing. nom. nov.

*Polyporus megaloporus* Mont., Ann. Sc. Nat. IV. 1: 124. 1854 non Pers., Mycol. Eur. 2: 88. 1825.

*Pleurotus* sect. **Calyptrati** Sing. sect. nov. Pileo velo pelliculari ornato; stipite fortiter reducto; tramate lamellarum irregulari, multis hyphis crassitunicatis. Typus sectionis: *P. calyptratus* (Lindb. in Fr.) Sacc.

*Pleurotus* sect. **Tuberregium** Sing. ex Sing. sect. nov. Sclerotio typi Pachymatis praesente. Typus sectionis: *P. tuber-regium* (Fr.) Sing.

### **Pleurotus subtilis** (Berk.) Sing. comb. nov.

*Lentinus subtilis* Berk., J. Linn. Soc., Bot. 15: 83. 1876 (published 1877).

### **Pleurotus fockei** (Miq.) Sing. comb. nov.

*Lentinus fockei* Miquel, Tijdsch. v. d. Wis. en Nytturk. Wetenschapen 5: 188. 1852.

### **Pleurotus geesterani** Sing. spec. nov.

*Pleurotus* manifeste annulatus, *P. sajor-caju* (Fr.) Sing. peraffinis sed sporis majoribus ( $7-9.2 \times 2.2-2.8 \mu$ , plerumque  $7.5 \mu$  longis) differt; cystidiis hyalinis vel flavis,  $18-27 \times 3-4 \mu$ , hyphis tramatis communicantibus, tenuitunicatis vel subcrassitunicatis; cystidiolis praesentibus; cheilocystidiis versiformibus, clavatis, fuscoideis, clavato-mucronatis, etc.,  $20-30 \times 4.5-8 \mu$ ; fasciculis hypharum conglomeratarum hymenialium projicientium numerosis; hyphis inamyloideis, fibulatis. Lamellis confertis sed haud confertissimis usque ad anulum decurrentibus. Totus fungus albus est. Ad truncum frondosum. South Africa, Cape Prov. Humansdorp Distr., 2-XII-1949. Leg. Maas Geesteranus no. 6736 (L), typus.

### **Pleurotus squarrosulus** (Mont.) Sing. comb. nov.

*Lentinus squarrosulus* Mont., Ann. Soc. Nat. 18: 257, n. 138, 1842.

### **Pleurotus javanicus** (Lév.) Sing. comb. nov.

*Lentinus javanicus* Lév., Ann. Sc. Nat. III. 5: 118, 1846.

### **Panus erinitus** var. **sehnyderi** (Speg.) Sing. comb. nov.

*Lentinus sehnyderi* Speg., An. Soc. Cient. Arg. 9: 165. 1880.

### **Panus badius** (Berk.) Sing. comb. nov.

*Lentinus badius* Berk., London J. Bot. 1: 148. 1842.

### **Panus brevipes** (Cooke) Sing. comb. nov.

*Lentinus brevipes* Cooke, Grevillea 14: 12. 1885.

\*) Prima series sub titulo aequali in Sydowia 2: 26-42. 1948 publicata.

**Panus inquinans** (Berk.) Sing. comb. nov.

*Lentinus inquinans* Berk., Hooker's J. Bot. 6: 132. 1954.

**Panus polychrous** (Lév.) Sing.

*Lentinus polychrous* Lév., Ann. Sc. Nat. III. 2: 175. 1844.

**Hygroaster nodulisporus** (Dennis) Sing. comb. nov.

*Hygrophorus nodulisporus* Dennis, Kew Bull. 1953 (no. 2): 259. 1953.

**Hygroaster asterosporus** (Lange) Sing. comb. nov.

*Omphalia asterospora* Lange, Dansk Bot. Ark. 6 (5): 10. 1930.

**Aeruginospora dennisiana** (Sing.) Sing. comb. nov.

*Hygrotrama dennisianum* Sing., Sydowia 12: 221. 1958.

**Aeruginospora hymenoccephala** (Sm. & Hesler) Sing. comb. nov.

*Hygrophorus hymenoccephalus* Sm. & Hesler, J. Elisha Mitch. Sc. Soc. 56: 311. 1940.

**Aeruginospora paupertina** (Sm. & Hesler) Sing. comb. nov.

*Hygrophorus paupertinus* Sm. & Hesler, Lloydia 5: 13. 1942.

**Aeruginospora microspora** (Sm. & Hesler) Sing. comb. nov.

*Hygrophorus microsporus* Sm. & Hesler, Lloydia 5: 11. 1942.

*Hygrocybe* sect. **Unguinosae** (Herink) Sing. = *Neohygrocybe* sect.

*Unguinosae* Herink, Act. Mus. Boh. Sept. Liberec. 1: 72. 1959.

**Neohygrophorus** Sing. ex Sing. gen. nov.

A *Camarophyllis* differt sporis amyloideis. Typus generis: *Hygrophorus angelesianus* Sm. & Hesler — Quamquam subgenus *Pseudohygrophorus*, quod est basonymon generis *Neohygrophori* Sing. Lilloa 22: 149. 1949 (publ. 1951), contineat unam solam speciem (typum) ita ut huius diagnosis latina ab auctoribus (Smith et Hesler) provista ut diagnosis generico-specifica interpretari possit, descriptionem latinam novam praesentamus ubi genus ut novum proponitur.

**Neohygrophorus angelesianus** (Sm. & Hesler) Sing. comb. nov.

*Hygrophorus angelesianus* Sm. & Hesler, Lloydia 5: 6. 1942.

(Tricholomataceae) tribus **Lyophylleae** Kühner ex Sing. trib. nov. Basidiis granulazione carminophila instructis. Typus tribus: *Lyophyllum* Karst.

*Calocybe* sect. **Heterosporae** Sing. ex Sing. sect. nov. (*Calocybe* sect. *Heterosporae* Sing. sine descriptionibus latinis, 1951). Pileo pigmento incrustante ornato; sporis basidiisque medioeriter voluminosis (sporis 6.7  $\mu$  cc. longis); illis echinatis, his pro ratione magnis; epicute haud cellulari; stipite velo destituto; in silvis tropicalibus. Typus sectionis: **Calocybe heterospora** Sing. spec. nov.

Pileo flavo-ochraceo ("yellow ocher", Ridgway), disco brunneo ("amber brown" Ridgway), margine extremo eburneo ("ivory yellow" Ridgway) e tegumento tenui sericeo, subopimo sed haud viscido, levi, glabro, umbilico minutissimo ornato, margine primum incurvato, ceterum convexo, 23 mm. lato; lamellis cremeis, angustis, confertissimis, adnatis; stipite ad apicem lamellis concolori et subfibrilloso,

ceterum sordido subfibrillosoque sed glabrescente, basin versus attenuato, 29 mm. longo, 6 mm. lato ad apicem sed usque ad 3 mm. diam. angustato ad basin; rhizomorphis albis inter folia extensis, ad basin radicelliformiter adhaesis; velo nullo; carne cremea, farinolente. Sporis (6)— $6.8 \times 3.5-4.2 \mu$ , hyalinis, ellipsoideis, fortiter asperulis, applanatione suprahilari praeditis, inamyloideis; basidiis  $28-39 \times 7.5-9.5 \mu$ , i. e. pro ratione longis latisque, sed haud Hygrophora-cearum modo cum manifeste granulatione carminophila impletae sint; cystidiis nullis visis; epicute ex hyphis repentibus levibus (pigmento excepto) efformata, tenuissima; hypodermio ex hyphis similibus crassioribus efformato; hyphis cuticularibus pigmento pallide roseolo intracellulari nec non pigmento atrocastaneo incrustante ornatis; hyphis in tramate hymenophorali regulari cylindraceis, densis, parallelis,  $2-12 \mu$  crassis hyalinis; elementis subhymenii ramosis, densis, exiguis; hyphis omnibus fibulatis, inamyloideis. In silva tropicali Floridae (U.S.A.), Matheson Hammock, 9—XI—1942. Leg. Singer no. F 1452 C (FH), typus.

**Calocybe constricta** (Fr.) Kühner comb. nov.

*Agaricus constrictus* Fr., Syst. Mycol. 1: 28. 1821.

**Calocybe pseudoflammula** (Lange) Morten Lange comb. nov.

*Tricholoma pseudoflammula* Lange, Dansk Bot. Ark. 8 (3): 24. 1933.

**Calocybe chrysentera** (Fr.) Sing. comb. nov.

*Agaricus chrysenteron* Bull.?, ex Fr., Syst. Mycol. 1: 126. 1821.

**Calocybe juncicola** (Heim) Sing. comb. nov.

*Tricholoma chrysenteron* var. *juncicola* Heim, Treb. Mus. Cienc. Nat. Barcelona, 15 (3): 101—102. 1934.

**Calocybe alpestris** (Britz.) Sing. comb. nov.

*Agaricus alpestris* Britz., Hym. Südb. 8: 4. 1891.

**Calocybe naucoria** (Murr.) Sing.

*Melanoleuca naucoria* Murr., North Amer. Flora 10: 15. 1914.

**Calocybe caucasica** Sing.

Pileo vitellino brunneo-tincto vel vitellino-alutaceo-brunneo, subglabro, sublevi, convexo, dein applanato, margine initio involuto, 10—35 mm. lato; lamellis concoloribus, ammoniaci ope sulphurescentibus, subconfertis, anguste adnexas, demum separantibus, interdum dente decurrentibus; sporis in massa albis; stipite pileo concolori vel dilutiore, glabro, subcylindraceo vel basin versus incrassato, farcto,  $22-40, \times 2-5$  mm.; carne pallide sulphureo vel superficiebus concolori; odore nullo; sapore miti; consistentia carnosa. Sporis  $3.3-4 \times 1.8-2.5 \mu$ , levibus, inamyloideis, ellipsoideis; basidiis brevibus; cystidiis nullis; epicute pilei ex hyphis  $3.3-6.6 \mu$  latis et sphaerocystibus globosis  $8-14 \mu$  diam. efformata; hyphis fibulatis. Ad terram et herbas putridas in pratis alpinis et subalpinis. U.R.S.S.,

Caucasus, Krasnodarsk. reg., Yatyrgvarta, 29—VII—1935, Leg. L. N. Vassiliev a (LE, KAZ), typus.

**Clitocybe carnea** (Bull. ex Fr.) Kühner comb. nov.

*Agaricus carneus* Bull. ex Fr., Syst. Mycol. 1: 130. 1821.

**Clitocybe persicolor** (Fr.) Sing. comb. nov.

*Agaricus ionides* var. *persicolor* Fr., Icon. Sel. 1: 35. 1870.

**Clitocybe glaucoalba** (Sing.) Sing. comb. nov.

*Tricholoma glaucoalbum* Sing., Sydowia 2: 28. 1948.

**Clitocybe ericeticola** Sing. nom. nov.

*Agaricus ericetorum* Bull. ?, ex Fr. Epicr. p. 73. 1838 non Fr. (1821).

*Clitocybe* sect. **Aberrantissimae** Sing. sect. nov. Sectioni Candidis analoga sectio sed stipite semper excentrico obliquoque (habitu pleuroidi), lignigeno, typice (in speciebus tropicalibus vel subtropicalibus) cystidiis praesentibus; tramate hymenophorali interdum non typi *Clitocybarum* sed magis irregulari aut hyphis subcrassitunicatis praesentibus. Typus sectionis: *C. aberrantissima* Sing.

*Clitocybe* sect. **Augustinenses**. Basi communi squamosa sub fasciulis carpophorum praesente; cystidiis nullis; ceterum sectionibus Candidis et Aberrantissimis analoga sectio. Typus sectionis: *C. augustinensis* Sing.

**Clitocybe decembris** Sing. nom. spec.

*Clitocybe bicolor* "(Pers. ex Pers.)" Lange non Pers.

*Clitocybe vibecina* (Fr.) Quél. sensu Konrad & Maublanc (sec. Kühner & Romagnesi).

Pileo griseofusco, margine subtiliter striatulo. demum infundibuliformi, 30—50 mm. lato; lamellis decurrentibus, pallidis vel griseolo-albidis; stipite magis sericeo quam farinoso-pruinoso ad apicem. Sporibus usque ad 8  $\mu$  longis. Ceterum *C. metachroae* (Fr.) Kummer et *C. favrei* Kühn. & Romagnesi simillima species. Typus est icon Langei, Flora Agaricina Danica, pl. 38, fig. B.

*Clitocybe* sect. **Farinolentes** (Sing.) Sing. = *Collybia* sect. *Farinolentes* Sing., Pap. Mich. Acad. Sc. 32: 122. 1946 (publ. 1948).

**Clitocybe vulgaris** Sing. spec. nov.

Pileo vix viscido, griseo-ochraceo; lamellis rotundato-adnexas vel adnato-decurrentibus, pallide cremeo-griseolis, subpallidis, latis, in junioribus irregulariter decurrentibus, mediocriter distantibus vel distantibus; stipite pileo pallidior, anguste tubulari, initio innate sericeo-fibrilloso (longitudinaliter) sed mox glabrescente; sporibus in massa, carne farinolente, mycelio — albis. Sporibus 6.5—8.2  $\times$  4.8—5.3  $\mu$ , membrana hyalina tenui homogenea, inamyloidea, levi instructis; basidiis tetrasporis; hyphis cuticularibus pilei pigmento-incrustatis, omnibus fibulatis; cystidiis et dermatocystidiis nullis. Ad terram prope domicilia plenumque sub arboribus. U.S.A., Mass., Cambridge; Leg. Singer (FH), typus. Haec species *Omphaliae barbularum* Romagnesi proxima est et praecipue habitu magis collybioide, pileo minus



gelatinisato in strato cuticulari differt. Descriptio amplior in Pap. Mich. Acad. Sc. 32: 121—122. 1946 (1948). Probabiliter *Omphalina mellea* Murr., Proc. Fla. Acad. Sc. 7: 112. 1945 et *Collybia ignobilis* auctorum Americanorum p. p. identicae sunt, sed ad Clitocybes transfertae homonymae evadunt.

*Tricholoma* subgen. **Pardinieutis** Sing. subgen. nov. Fibulis praesentibus; strato cuticulari pilei ex hyphis haud intertextis atque subparallelis efformato, pileo superficialiter tomentoso-subsquamoso, haud viscido; odore debili vel nullo; sporis pro ratione maiusculis ( $8-10 \times 5-7 \mu$ ); pigmento abundante prope superficiem pilei. Typus subgeneris:

*Tricholoma* (subgen. *Tricholoma*) sect. **Adusta** Sing. sect. nov. Pileo glabro, hygrophano, griseo, umbrino, atrofuligineo, opace fusco vel olivaceofusco; habitu carpophorum collybioide (vel subcollybioide). Typus sectionis: *T. adustum* (Murr.) Murr.

**Armillariella omnituens** (Berk.) Sing. comb. nov.

*Agaricus omnituens* Berk., Hookers J. Bot. & Kew Gard. Misc. 2: 46. 1850.

**Omphalina telmatiaea** (Berk. & Cooke) Dennis ex Sing. comb. nov.

*Agaricus telmatiaeus* Berk. & Cooke, in Cooke, Handb. Brit. Fung. 2nd ed., p. 93, 1885.

*Gerronema* subgen. **Omphaliopsis** (Pilát in Kaviná & Pilát emened. Jossierand) Sing. = *Pleurotus* sect. *Omphaliopsis* Pilát in Kaviná & Pilát (em. Joss.) Atlas Champ. Eur. 2: 135. 1935. Species minutiores stipite excentrico vel subcentrali, pigmento destitutae, fibuligerae; tramate hymenophorali ex hyphis filamentosis tenuibus efformato, hyphis omnibus tenuitunicatis, in tramate hymenophorali fortiter intertextis sed mediostrato interdum regulari intercalato. Frequenter inter muscos crescentes. Typus sectionis: *Pleurotus mutilus* (Fr.) Gillet (= *Gerronema jossierandii*, Sing.)

**Gerronema jossierandii** Sing. spec. nov.

Pileo albo, depresso; lamellis albis, ochrascentibus, decurrentibus; sporis in massa albis; sporis levibus,  $6.3-7.5 \times 3.5-4.5 \mu$  ellipsoideis, haud angularibus, inamyloideis; basidiis tetrasporis; hyphis epicutis pilei haud differentiatis, fibuligeris, inamyloideis; stipite praecipue adulatorum excentrico; tramate hyalino, ex hyphis irregularibus (cuticula pilei et mediostrato nonnumquam exceptis) filamentosis, haud gelatinosis efformato. Ad Muscos praesertim supra truncos. Hollandiae, Terschelling 16—VII—1955. Leg. P. B. Jansen, det Bas (L.), typus. Est *Pleurotus mutilus* (Fr, sensu Jossierand) vix Fr. **Gerronema albidum** (Fr.) Sing. comb. nov.

*Cantharellus albidus* Fr., Syst. Mycol. 1: 319. 1821.

*Gerronema* sect. **Xanthophylla** (Sing.) Sing. (= *Marasmiellus* sect. *Xanthophylli* Sing. nom. subnud.) sect. nov. Hyphis fibuligeris; carne subtenacella, elastica; pigmentis flavis vel aureis praesentibus prae-

sertim in lamellis, raro sine pigmento flavo. Typus sectionis: *G. icterinum* (Sing.) Sing.

**Gerronema strobodes** (Berk. & Mont.) Sing. comb. nov.

*Agaricus strobodes* Berk. & Mont., in Mont., Syll. Gen. spec. crypt. p. 112. 1856.

**Gerronema clavuligerum** (Sing.) Sing. comb. nov.

*Armillariella clavuligera* Sing. in Singer & Digilio, Lilloa 25: 70. 1951 (publ. 1952).

*Gerronema* subgenus **Quetria** Sing. subg. nov. Pigmento praesente at haud aurantiaco vel olivaceo vel rubido-castaneo; a subgenere *Gerronema* differt praesertim structura tramatis hymenophoralis regulari et contextu carnoso; cystidiis in tyo praesentibus et fibulis nullis. Typus subgeneris *G. minutum* (Sing. & Digilio in Sing.) Sing.

**Gerronema minutum** (Sing. & Digilio) Sing. comb. nov.

*Armillariella minuta* Sing. & Digilio in Sing., Sydowia 8: 106. 1954.

*Gerronema* subgen. **Romagnesia** (Sing.) Sing. = *Omphalina* subgen. *Romagnesia* Sing., Mycologia 39: 83. 1947.

*Gerronema* (subg. *Romagnesia*) sect. **Cyathiformia** Sing. esct. nov. Cystidiis nullis; fibulis praesentibus; pileo haud aurantiaco; contextu carnoso. Typus subgeneris: *G. cyathiforme* (Berk. : Curt.) Sing.

**Gerronema bryogeton** Sing. spec. nov.

Pileo olivaceo („tennis“ M&P), convexo-applanato, centro plano vel depresso circum discum papillatum vel subumbonatum, subglabro sed in siccis s. l. subtomentoso, opaco, levi, 10—12 mm. lato; lamellis 9—10, haud intervenosis, olivaceo-flavis („banana“ M&P), nonnullis furcatis, intermixtis, arcuatis, moderatim latis, decurrentibus, distantibus; stipite ad apicem lamellis concoloribus, umbrino vel brunneo basin versus, subglabro, subcurvatulo, aequali, 8.5—9.5×0.5—1 mm. Sporis 7.5—9.5—(11)×4—5.5—(6.2)  $\mu$ , hyalinis, vel pallide aurantiacis, levibus, inamyloideis, applanatione levi suprahilari provisis vel destitutis, ellipsoideis, membrana homogena instructis; basidiis aurantiacis (in alkali) 20—36×6—7  $\mu$ , tetrasporis, paucis 2—3-sporis, clavatis; cystidiis et cheilocystidiis nullis; epicute pilei ex hyphis irregulariter elongatis vel, filamentosis levissimis repentibus subhyalinis radialiter dispositis eformata; infra epicutem hyphis angustissimis vel admodum voluminosis repentibus intertextis; tramate hymenophorali regulari, subintertextis ex hyphis dense congregatis eformato; hyphis omnibus inamyloideis, nonnullis fibulatis, aliis defibulatis. Superficie pilei NH<sub>4</sub>OH et KOH ope castanescente. Ad thallum Musci supra lignum in silva tropicali-montana. Bolivia, Yungas, La Paz, Nor-Yungas, Carmen Pampa, I—II—1956. Leg. Singer no. B 811 (LIL), typus.

**Gerronema postii** (Fr.) Sing. comb. nov.

*Agaricus postii* Fr., Monogr. Hymenom. Suec. 2: 291. 1863.

**Gerronema macrosporum** (Sing.) Sing. comb. nov.

*Phaeomyceca macrospora* Sing. in Sing. & Digilio, Lilloa 25: 176. 1951 (publ. 1952).

**Gerronema venustissimum** (Fr.) Sing. comb. nov.

*Agaricus venustissimus* Fr., Monogr. Hymenom. Succ. 2: 289. 1863.

**Gerronema brevbasidiatum** (Sing.) Sing.

*Clitocybe brevbasidiata* Sing., Ann. Mycol. 41: 45-6. 1943.

*Gerronema* (subg. *Romagnesia*) sect. **Fibulae** (Sing.) Sing. = *Marasmiellus* sect. *Fibulae* Sing., Sydowia 2: 32. 1948.

**Gerronema setipes** (Fr.) Sing. comb. nov. (sensu Ricken!).

*Agaricus setipes* Fr., Syst. Mycol. 1: 159. 1821, esp. var. *acrocyanus* Fr., Icon. pl. 75, fig. 4. 1873.

**Gerronema alexandri** Sing. spec. nov.

A *Gerronema fibula* (Bull. ex Fr.) Sing. proxima absentia fibularum certe distinguitur. Inter Polytricha, gregatim. U. S. A., Michigan, prope Tahquamenon Falls, 23-VI-1953, leg. Singer no. N 127 (F). **Callistosporium chrysophorum** Sing. spec. nov.

Pileo dilute olivaceobrunneo vel alutaceobrunneo ("light brownish olive" vel "buffy brown" Ridgway), interdeum isabellino ("isabella color" Ridgway), dein cremealutaceo vel isabellino ("cream buff" vel "isabella color" Ridgway), dehydratione aut isabellino auf frequentius brunneolo-ochraceo ("colonial buff" Ridgway), levi, glabro, hygrophano, levissimo in humidis, usque ad 19 mm. lato; lamellis subconcoloribus, mediocriter latis (usque ad 3.5 mm.), confertis vel subconfertis, rotundatis vel emarginatis, anguste adnexis; stipite concolori, leniter longitudinaliter innate striatulo-fibrilloso, cavo, saepe canaliculato, aequali vel subtus vel ad apicem incrassato, 34 x 4 mm. cc.; mycelio tenui. albo ad basin; velo nullo; carne subconcolori sed interiore in parte cavitatis stipitis albida; odore iucundo; sapore amaro. Sporis 4.8-5.5 x 3.7-3.9  $\mu$ , hyalinis, dein pigmento amorpho haud dissoluto melleo-aureo impletis, levibus; basidiis 20-24 x 5-6.2  $\mu$ , plerumque tetrasporis, in multis basidiis corpusculum aureo-melleum praesens; cheilocystidiis nullis; hyphis hyalinis, defibulatis. Exsiccatione lamellae brunnescent (,vinaceous cinnamon" Ridgway); solutionibus alkalinis fortibus pileus cinnamomeus fit, sed etiam pallescit. Ad truncum *Pini* putridum. U.S.A., Fla., Alachua Co., south of Dayville, 8-VI-1943. Leg. Singer no F 2227 (F), typus. **Lactocollybia** sect. **Bertrandiella** (Heim) Sing. stat. nov.

*Bertrandiella* Heim (ut gen.) Rev. Mycol. 24: 192. 1959.

**Lactocollybia ianthina** (Heim) Sing. comb. nov.

*Bertrandiella ianthina* Heim, l. c.

**Lactocollybia microspora** Sing. spec. nov.

Habitu Marasmielli! Pileo albo, putredine tantum flavescente, levi, convexo, 5-9 mm. lato; lamellis albis, demum flavescentibus, moderatim confertis vel subconfertis, latis, angustato-adnato-sub-

decurrentibus; stipite concolori, subcentrali vel manifeste excentrico, curvato, glabro, levi, aequali, brevi, 3—5 × 0.5—0.8 mm.; carne alba, immutabili, inodora. Sporis parvis: 5.5—6.5 × 3.8—4.8  $\mu$ , hyalinis, levibus, ellipsoideis; basidiis clavatis, 23—24.5 × 6—6.5  $\mu$ , tetrasporis; cheilocystidiis versiformibus, 35—42 × 9—10  $\mu$ , plerumque fusoido-ventricosus vel subcylindraceis vel ampullaceis, interdum constrictis, hyalinis, contentu visibili in ammoniaco destitutis, levibus; gloeocystidiis numerosissimis ad latera aciemque lamellarum, e. gr. 24—32 × 6.5—8.5  $\mu$ , saepe gloeo-vasculis communicantibus, plerumque fusoides apice rotundatis, contentu laete aureo-luteo gaudentibus, intense azureis caerulei cresylicae ope; dermatopseudocystidiis gloeocystidiis hymenialibus homologis sed frequentius maioribus (e. gr. 80—85 × 10—15  $\mu$ ), gloeovasculis communicantibus; dermatocystidiis haud gloeocystidioideis rarioribus, cheilocystidiis analogis sed minoribus atque minus conspicuis, hyalinis, e. gr. 27—28 × 5—6  $\mu$ ; tramate hymenophorali pallide flavido, regulari, ex hyphis normalibus hyalinis nec non gloeovasculis efformato, omnibus parallelis vel subparallelis; tramate pilei laxo contexto cum spatiis aeriferis sed haud gelatinoso, irregulari, hyalino; hyphis omnibus hyalinis, levibus, inamyloideis, fibuligeris. Ad latus inferius trunci vetusti in silva subtropicali. Argentina, Misiones, Iguazú, Puerto Iguazú, 12—IV—1957. Leg. Singer no. M 950 (LIL), typus. — *L. cycadicolae* (Josserand) Sing. nec non facile etiam *Gymnopodi marasmioidei* Murr. affinis sed certe ab ambobus distincta species.

**Leptoglossum tremulum** (Schaeff. ex Fr.) Sing. comb. nov.

*Agaricus tremulus* Schaeff. ex Fr., Syst. Mycol. 1: 191. 1821.

*Tricholomataceae* tribus **Biannularieae** Sing. (1936) ex Sing. Tramate lamellarum laterali in juventute; sporis amyloideis; lamellis de-currentibus; velo praesente; sporis uni-nucleatis. Genus typicum: *Biannularia* G. Beck = *Catathelasma* Lovej.

**Asterotus bicolor** (Mont.) Sing. ex Sing. comb. nov.

*Panus bicolor* Mont., Crypt. Guy. no. 329. 1855.

**Stigmatolemma poriaeforme** (Pers. ex Mérat) Sing., comb. nov.

*Peziza anomala* var. *poriaeformis* Pers. Syn. 1: 656. 1801 ex Mérat.

**Pseudoclitocybe obbata** (Fr.) Sing. comb. nov.

*Agaricus obbatus* Fr., Epier., p. 74. 1838.

**Pseudoclitocybe oregonensis** (Murr.) Sing., comb. nov.

*Clitocybe oregonensis* Murr. Mycologia 5: 211. 1913.

**Pseudoclitocybe coprophila** (Speg.) Sing., comb. nov.

*Clitocybe coprophila* Speg., Bol. Ac. Nac. Cienc. Cordoba 11: 386. 1889.

**Pseudoomphalina felloides** (Kauffm.) Sing. comb. nov.

*Cantharellula felloides* (Kauffm.) Sing., Lilloa 22: 238. 1949 (publ. 1951).



**Pseudoomphalina intermedia** (Kauffm.) Sing. comb. nov.

*Cantharellula intermedia* (Kauffm.) Sing. l. c.

**Clitocybula abundans** (Peck) Sing. ex Sing. comb. nov.

*Agaricus abundans* Peck Ann. Rep. N. Y. St. Mus. 29: 38. 1878.

*Clitocybula* subgen. **Neocantharellula** (Sing.) Sing. n. c.

*Cantharellula* subgen. **Neocantharellula** Sing., Sydowia 6: 194. 1952.

**Clitocybula lacerata** (Lasch) Sing. comb. nov.

*Agaricus laceratus* Lasch apud Fr., Epicrisis p. 97. 1838 (syn. ex clusis!).

**Clitocybula aperta** (Peck) Sing. comb. nov.

*Agaricus apertus* Peck, Ann. Rep. N. Y. State Mus. 30: 38. 1878.

**Clitocybula atrialba** (Murr.) Sing. comb. nov.

*Clitocybe atrialba* Murr., Mycologia 5: 207. 1913.

**Clitocybula familia** (Peck) Sing. comb. nov.

*Agaricus familia* Peck, Ann. Rep. N. Y. State Cab. 23: 79. 1872.

**Clitocybula oculus** (Peck) Sing. comb. nov.

*Agaricus oculus* Peck, Ann. Rep. N. Y. State Cab. 23: 84. 1872.

**Clitocybula tilieti** (Sing.) Sing. comb. nov. 1)

*Fayodia tilieti* Sing., Ann. Mycol. 41: 63. 1943.

**Clitocybula dusenii** (Bres.) Sing. comb. nov.

*Panus dusenii* Bres., Overs. K. Vet-Akad. Förhandl. 1900 (2): 312. 1900.

*Porpoloma* subgenus **Pogonoloma** Sing. subgen. nov. Margine pilei piloso. Odor inocybis bongardii. Pigmento intracellulari. Cheilocystidiis paulum vel haud evolutis. Typus: *Tricholoma spinulosum* Kühn. & Romagnesi.

**Porpoloma spinulosum** (Kühner & Romagnesi) Sing. comb. nov.

*Tricholoma spinulosum* Kühn. & Romagnesi. Bull. Soc. L. Lyon 16: 134. 1947.

*Porpoloma* subgen. **Pseudotricholoma** (Sing.) Sing. = *Cantharellula* subgen. *Pseudotricholoma* Sing. Sydowia 2: 30. 1948.

**Porpoloma umbrosum** (A. H. Smith & Walters) Sing. comb. nov.

*Tricholoma umbrosum* A. H. Smith & Walters, Mycologia 35: 477. 1943.

**Leucopaxillus cerealis** (Lasch) Sing. comb. nov.

*Agaricus cerealis* Lasch, Linnaea 4: 526. 1829.

**Melanoleuca microcephala** (Karst.) Sing., comb. nov.

*Tricholoma microcephalum* Karst., Hedwigia 20: 177. 1881.

**Grifola sordulenta** (Mont.) Sing. comb. nov.

*Polyporus sordulentus* Mont., in Gay, Hist. Nat. Chile 7: 357. 1854.

**Collybia retiaria** (Berk.) Sing. comb. nov.

*Paxillus retiarius* Berk., Hook. J. Bot. & Gard. Misc. 8: 133. 1856.

1) Haec et 7 transferentiae anteriores e literis prop. cl. Deightonii Regn. Veg. 14: 39 (G/H). 1959 necessariae apparent sed cf. Sydowia 8: 110. 1954.

**Collybia kuehneriana** Sing. nom. nov.

*Marasmius bresadolae* Kühn. & Romagnesi, Flore anal. p. 88. 1953, non *Collybia bresadolae* Sacc & Sacc (1905).

*Marasmius erythropus* „Pers.“ sensu Bresadola pl. 496, non *Agaricus erythropus* Pers.

We refer to the original diagnose given by Bresadola (l. c.) and the material preserved under that name at the Riksherbarium in Stockholm.

**Collybia konradiana** Sing. nom. nov.

*Collybia fuscopurpurea* (Pers. ex Fr.) Kummer sensu Konrad & Maublanc, Kühner & Romagnesi (as *Marasmius*).

A *C. fuscopurpurea* (Pers. ex Fr.) sensu Persoon, Fries, (ut *Agaricus*) differt stipite brunneo-fulvo subrubescente sporisque paulum minoribus pro medio et praecipue granulis pigmenti extracellularis absentibus in carne tramateque lamellarum. Typus sub nom. *fuscopurpureae* in Herb. L. conservatur.

**Collybia neotropica** Sing. nom. nov.

Pileo cinnamomeo in umbone, pallidore in parte externa, albido in margine extremo, hygrophano, sulcato usque ad (sed umbone excepto) centrum, convexo, umbone lato obtuso praedito, 23 mm. lato. Lamellis sordide pallidis, subalbidis, moderate latis, subdistantibus. adnexas vel adnatis, haud intervenosis. Stipite concolori lamellis sed basin versus magis griseolo-cinnamomeo-pallido, subtiliter velutino, basin versus incrassato,  $33 \times 2.5$  „mm.; mycelio basali fibrilloso sordide pallido. Contextu albido, tenui; odore *Collybiae dryophilae*.

Sporis hyalinis, inamyloideis, levibus, ellispoideis vel oblongis,  $8.2-9.7 \times 4.3-4.8$   $\mu$ . Basidia clavatis tetrasporis, hyalinis,  $29-32 \times 6.8-9.7$   $\mu$ . Cystidiis nullis. Cheilocystidiis filamentosis, irregularibus, nonnumquam basin versus incrassatis vel ventricosus vel apicem versus incrassatis, plerumque basidiis angustioribus, plerumque hyalinis, saepe pluries constrictis,  $2-6.5$   $\mu$  latis. Tramate hyalino, regulari, hyphis filamentosis. Epicute ex hyphis tenuibus, filamentosis efformata; hyphis eis hyalinis vel dilute brunneolis, levibus vel raro subsinuose tunicatis, rarius ramulos terminales breves emittentibus qui pilositatis stipitis elementa in mentem revocant sed sunt leves, haud incrustati, Ramealium structuram haud imitant. Pilis stipitis erectis, filamentosis, nonnumquam incrassatis cheilocystidiaque in mentem revocantibus, usque ad 60  $\mu$  longis,  $2.5-8.3$   $\mu$  latis, hyalinis, haud incrustatis, multis apice capitato vel subcapitato gaudentibus, sed apice sat moniliformi in nonnullis pilis, capitulo semper exiguo; pilis illis totam longitudinem stipitis tegentibus. Hyphis omnibus hyalinis vel subhyalinis, inamyloideis, fibulatis, in pileo radiati, dispositis (epicute inclusa).

Ad humum inter detritum silestrem ad marginem umbrosorum platationum. BOLIVIA, La Paz, Coroico (Nor-Yungas), 15-II-1956.

**Leg.** Singer no. B 1173, typus, in Herbario Lilloano conservatur (LIL).

This species may be identical with one collected by Dennis and determined as *Marasmius longipes* Mont. The Australian type of *M. longipes* Mont. is also a *Collybia* of the *C. confluens* group but I am not certain whether it is the same species. At any rate, in *Collybia*, *M. longipes* would be a homonym. Dennis' material comes from Trinidad (no. 33, K).

**Collybia paurospora** (Sing.) Sing. comb. nov.

*Marasmiellus paurosporus* Sing. Sydowia 11: 228. 1958.

**Collybia fascicularis** (Rick ex Sing.) Sing. comb. nov.

*Marasmiellus fascicularis* Rick ex Sing., Lilloa 26: 112. 1953.

**Collybia urubambae** Sing.

A *Collybia* castaneoidisca et *C. fasciculari* differt absentia cystidiorum, a *C. subfumosa* differt sporis majoribus ( $9-12.5 \times 3.5-4.1 \mu$ ). Pileo 12-53 mm. lato; stipite  $23-90 \times 1.3-7$  mm. Ad detritum (folia, e. gr. Piptadeniae et ramulos) et ad humum in silva montana et subalpina tropicali. Typus ad fl. Rio Urubamba, Machu Pichu (Peru) leg. Singer (20-V-1958, no. B 3024, LIL) in altitudine 2000 m. s. n. m. Species identica a cl. Dennis sub nom. *Marasmiellus urubambae* in Sierra S. Domingo (Mérida, Venezuelae) lectus fuit (J. J. Buza & Dennis no. 1713, K) et mihi in prov. Tucumán (Quebrada de La Higuera, 25-I-1955) no. T 2097 (LIL) obvia

**Collybia distortiformis** (Sing.) Sing. comb. nov.

*Marasmiellus distortiformis* Sing. in Sing. & Digilio, Lilloa 25: 155. 1951. (publ. 1952).

**Collybia biformis** (Peck) Sing. comb. nov. **Coll**

**Collybia biformis** (Peck) Sing. comb. nov.

*Marasmiua biformis* Peck, Bull N. Y. St. Mus. 67: 25. 1903.

**Collybia domestica** (Murr.) Sing. comb. nov.

*Marasmius domesticus* Murr., Torr. Bot. Cl. Bull 66: 160. 1939.

**Collybia sacchari** (Sing.) Sing. comb. nov.

*Marasmiellus sacchari* Sing. in Sing. & Digilio, Lilloa 25: 170. 1951. (publ. 1952).

**Neoclitocybe** Sing. gen. nov.

*Marasmiellis* analogum et *Collybiis* interdum affine sed *Clitocybis* *Omphalinisque* plerumque habitu suo simillimum genus. Epicute structura Ramealium gaudente; lammellis decurrentibus; stipite centrali, mycelio basali plerumque praesente sed interdum sparso, sub lente haud insititio.

Species typica: *N. byssiseda* (Bres.) Sing. comb. nov. (*Omphalia byssiseda* Bres. in Rick, Fung. austroam. nol 47. 1906).

**Neoclitocybe subnimbata** (Rick) Sing. comb. nov.

*Clitocybe subnimbata* Rick, Lilloa 2: 259. 1938.

**Neoclitocybe irregularis** (Rick) Sing. comb. nov.

*Omphalia irregularis* Rick, Lilloa 2: 292. 1938.

**Neoclitocybe substenophylla** (Murr.) Sing. comb. nov.

*Marasmius substenophylla* Murr., Bull. Torr. Bot. Cl. 67: 153. 1940.

**Neoclitocybe nauseosa** (Rick) Sing. comb. nov.

*Clitocybe nauseosa* Rick, Broteria 17: 103. 1919.

**Neoclitocybe nivea** (Rick) Sing. comb. nov.

*Clitocybe nivea* Rick, Lilloa 2: 261. 1938.

**Neolitoocybe omphalina** (Sing.) Sing. comb. nov.

*Marasmiellus omphalina* (Sing., in Sing. & Digilio, Lilloa 25: 167. 1951. (publ. 1952).

**Neoclitocybe myceliosa** Sing. spec. nov.

Pileo brunneo. pellucide striato sulcatoque, umbilicato-infundibuliformi, membranaceo, 7—10 mm. lato. Lamellis dilute cremeo-grisellis, subangustis, subdistantibus, decurrentibus, Stipite vitreofusco, apice hyalino vel subhyalino, apicem versus incrassato, glabro vel subglabro, ad basin mycelio lanato griseolo-sordido, subtili ornato. Sporibus hyalinis, inamyloideis, oblongis vel late ellipsoideis, levibus, tenuitunicatis, 4—6.3 × 2.8—3.8 (4,5)  $\mu$ . Basidiis longis, tetrasporis, 30—40 × 5.7—7  $\mu$ . Cystidiis nullis. Cheilocystidiis irregularibus, haud conspicuis, exerescentiis praeditis vel destitutis, basidiis juxtapositis. Hyphis epicutis pilei filamentosis sed in maculis punctiformibus sub lente visibilibus congregatis et intertextis, exerescentiis et nodulis instructis, pigmento epicellulari incrustatis. Hyphis omnibus inamyloideis, fibulatis. N. omphalinae similis est. Ad lignum emortuum in silva tropicali montana. Specimen typicum (leg. Singer no. B 1211, Rio Yariza, Nor-Yungas, La Paz, Boliviae, 16-II-1956) in Herbario Lilloano conservatur.

**Neoclitocybe portentosa** Sing. spec. nov.

Pileo caeruleo-atro, convexo, centro fortiter depresso, in juvenilibus saepe umbilicato aut papillato, glabro, sicco, 14—72 mm. lato. Lamellis caerulescente-griseis, acie pallescente, 6 mm. latis, vel angustioribus, i. e. mediocriter latis, confertis, arcuato-decurrentibus, Stipite azureo dein pallide caeruleo, ad basin obscure luteo, glabro, solido, centrali vel saepius subexcentrico, erecto vel curvato, 16—17 × 2—5 mm. Basi sub lente manifeste strigoso, sparse mycelioso, albo. Sporibus hyalinis, membrana subtenui instructis, globosis vel breviter ellipsoideis, levibus, inamyloideis, 4.8—5.6 × 4—4.2  $\mu$ . Basidiis sublongis, 25—27 × 5—7  $\mu$ . Cystidiis nullis. Cheilocystidiis coralloideo-brassiciformibus Epicute structura Ramealium gaudente at debiliter evoluta, nodulis semilobatis dispersis vel congregatis ornata sed maxima e parte ex hyphis levibus vel sublevibus efformata. Tramate hymenophorali regulari vel subintertexto-subregulari; subhymenio bene evoluto. Hyphis tramatis pilei et tramatis hymenophoralis inamyloideis vel lenissime pseudoamyloideis, tenui- vel crassi-tuni-



catis, fibulatis. Pigmento in cuticula pilei (minus evoluto in tramate) nigrescenti-viridi vel viridulo ( $\text{NH}_4\text{OH}$ ,  $\text{KOH}$ ), amethystino in acidis ( $\text{HCl}$ ), membranas incrustante maculis vel lineis helicoideis, abundante. In palmis in silva tropicali pluviali. Bolivia, Riberalta, leg. Singer no. B 2368, 31-III-1956 (LIL), typus.

**Marasmiellus subingratus** (Dennis) Sing. comb. nov.

*Marasmius subingratus* Dennis, Trans. Brit. Mycol. Soc. **34**: 427. 1951.

**Marasmiellus cubensis** (Berk. & Curt.) Sing. comb. nov.

*Marasmius cubensis* Berk. & Curt. L. Linn. Soc. Bot. **10**: 296. 1868.

**Marasmiellus pulchellus** (Berk.) Sing. comb. nov.

*Marasmius pulchellus* Berk., Hook. J. Bot. & Kew Gard. Misc. **8**: 138. 1856.

**Marasmiellus eburneus** (Theissen) Sing. comb. nov.

*Marasmius eburneus* Theissen, Brot. **8**: 55. 1909.

**Marasmiellus synodicus** (Kunze in Fr.) Sing. comb. nov.

*Agaricus synodicus* Kunze in Fr., Linnæa **5**: 507. 1830.

**Marasmiellus subcoracinus** (Berk. & Curt.) Sing. comb. nov.

*Marasmius subcoracinus* Berk. & Curt., J. Linn. Soc. Bot. **10**: 294. 1869.

**Marasmiellus brasiliensis** (Berk. & Mont.) Sing. comb. nov.

*Marasmius brasiliensis* Berk. & Mont., Ann. Sc. Nat. III, **11**: 236. 1849.

**Marasmiellus rugulosus** (Berk. & Curt.) Sing. comb. nov.

*Marasmius rugulosus* Berk. & Curt., L. Linn. Soc. Bot. **10**: 294. 1869.

**Marasmiellus purpureoalbus** (Petch) Sing. comb. nov.

*Marasmius purpureoalbus* Petch, Trans. Brit. Mycol. Soc. **31**: 37. 1947.

**Marasmiellus purpureus** (Berk. & Curt.) Sing. comb. nov.

*Marasmius purpureus* Berk. & Curt., J. Linn. Soc. Bot. **10**: 299. 1869.

**Marasmiellus luteus** (Dennis) Sing. comb. nov.

*Micromphale luteum* Dennis, Trans. Brit. Mycol. Soc. **34**: 458. 1951.

**Marasmiellus epochnous** (Berk. & Curt.) Sing. comb. nov.

*Marasmius epochnous* Berk. & Curt., in Berk. & Br., J. Linn. Soc., Bot. **14**: 41. 1873.

**Marasmiellus concolor** (Berk. & Curt.) Sing. comb. nov.

*Marasmius concolor* Berk. & Curt., J. Linn. Soc. Bot. **10**: 299. 1869.

**Marasmiellus devenulatus** Sing. spec. nov.

Pileo albido vel arenicolori ("Cuban sand" M & P), profunde sulcato, glabriusculo vel paullum subtomentoso, ostreato, convexo, 5—132 mm. lato. Lamellis concoloribus ("polo tan" M & P) vel pallidioribus, distantibus vel distantissimis, haud intervenosis, latius-

culis, integris sed nonnullis interdum furcatis, intermixtis. — Stipite concolori, exiguo, curvato, primum centrali vel subcentrali dein excentrico vel laterali et inconspicuo, fibrilloso in juvenilibus, insititio,  $2 \times 1$  mm. — Cortexto albo, inodoro. — Sporibus oblongis subclaviformibus, levibus, hyalinis, inamyloideis,  $9.5-13.2 \times 4-4.8$   $\mu$ . Basidiis  $29 \times 5.5$   $\mu$  vel minoribus, Cystidiis in juvenilibus absentibus, vel inconspicuis, dein manifestis sed paucis sparsis praesentibus, ventricosis, integris, hyalinis, levibus, tenuitunicatis, facile collabentibus, haud inerustatis,  $33-41 \times 9-13$   $\mu$ . Cheilocystidiis brassiciformibus vel furcato-diverticulatis, stramineo-hyalinis. Elementis epicuticularibus pilei irregularibus, versiformibus, diverticulatis. Hyphis laxe dispositis in pileo, radiatim filamentosis et vere gelatinosis prope subhymenium ubi irregulares sunt atque totis tenuibus, hyalinis (sed in epicute stramineohyalinis), in subhymenio haud gelatinascentibus, fibulatis. Ad stipulos ligneos in silva pluviali gregatim. Bolivia, Beni, Vaca Diez, Riberalta, I-IV-1956. Leg. Singer no. B 2417 (LIL) typus. — *A. M. distantifolius* lamellis haud intervenosis differt.

**Marasmiellus gossypinulus** (Speg.) Sing. comb. nov.

*Agricus gossypinulus* Speg., Bol. Acad. Nac. Cienc. Cordoba 11: 9. 1887.

*Marasmiellus* sectio **Distantifolii** Sing. sect. nov. Tramate gelatinoso  
Species typica: *M. distantifolius* (Murr.)

**Marasmiellus distantifolius** (Murr.) Sing. comb. nov.

*Scytinotus distantifolius* Murr., N. Amer. Flora 9: 239. 1915.

*Marasmiellus* sect. **Candidi** (Bat.) Sing. stat. nov., comb. nov.

*Marasmius*, subseries **Candidi** Bat., Mar. Eur. p. 27. 1919.

**Marasmius dealbatus** (Berk. & Curt.) Sing. comb. nov.

*Marasmius dealbatus* Berk. & Curt., J. Linn. Soc. Bot. 10: 294. 1869.

**Marasmiellus sprucei** (Berk.) Sing. comb. nov.

*Helionyces sprucei* Berk., Hook. J. Bot. & Kew Misc. 8: 134. 1856.

**Marasmiellus tetrachrous** (Sing. in Sing. & Digilio) Sing. comb. nov.

*Collybia tetrachroa* Sing. in Sing. & Digilio, Lilloa 25: 52. 1951 (publ. 1952).

**Marasmiellus saccharophilus** (Speg.) Sing. comb. nov.

*Omphalia saccharophila* Speg., Bol. Acad. Nac. Cordoba 28: 287. 1926.

**Marasmiellus stenophyllus** (Speg.) Sing. comb. nov.

*Marasmius stenophyllus* Mont. Ann. Sc. Nat. IV, 1: 116. 1854.

**Marasmiellus pruinosulus** (Murr.) Sing. comb. nov.

*Marasmius pruinosulus* Murr., North Amer. Flora 9: 258. 1915.  
Subtribus **Oudemansiellinae** Sing. subtrib. nov. (*Tricholomataceae*, trib. *Marasmiaceae* Fayod). Sporibus hyphisque inamyloideis; elementis epicutis inamyloideis, trichodermium vel stratum hymeniforme formatibus, rarius diverticulatis et hyphosis et repentibus; sporibus

plerumque subsodiametricis vel globosis, appendice hilari saepe majusculo; cystidiis plerumque manifestis ad latera lamellarum. Stipite haud manifeste insititio sed frequenter radicante. Genus typicum: *Oudemansiella* Speg.

*Oudemansiella* subgenus *Xerula* (Maire) Sing. stat. nov.

*Xerula* Maire (ut genus), Publ. Junta Cienc. Nat. Barcelona, p. 66. 1933.

*Oudemansiella pilosa* (Rick) Sing. comb. nov.

*Lentinus pilosus* Rick, Lilloa 2: 310. 1938.

*Oudemansiella macracantha* Sing. spec. nov. Ab *Oudemansiella steffenii* sporis echinatis, spinis 4—5.5  $\mu$  longis differt. Bolivia, Pando, Madre de Dios, Las Piedras, leg. Singer no. B 2525 (ILL), 5-IV-1956, typus.

*Oudemansiella kuehneri* (Romagnesi) Sing. comb. nov.

*Mycenella kuehneri* Romagnesi, Bull. Soc. Mycol. Fr. 56: 65. 1940.

*Xerulina* Sing. gen nov. (Tricholomataceae). Epicute pilei floccoso-diffracta, trichodermica, subpulverulenta, aurantiaco-, flavo-, vel olivaceopigmentata. Ceterum cum genere *Oudemansiella* subgen. *Xerula* congruens. Species typica: *Xerula chrysopepla* (Berk. & Curt.) Sing.

*Xerulina chrysopepla* (Berk. & Curt.) Sing. comb. nov.

*Lentinus chrysopeplus* Berk. & Curt., J. Linn. Soc. Bot. 10: 301. 1869.

*Xerulina lachnocephala* (Pat.) Sing. comb. nov.

*Collybia lachnocephala* Pat., Bull. Soc. Mycol. Fr. 18: 299 bis. 1902.

*Xerulina verruculosa* (Sing.) Sing. comb. nov.

*Xerula verruculosa* Sing., Rev. Myc. 18: 6. 1953.

*Mycenella margaritispota* (Lange) Sing. comb. nov.

*Mycena margaritispota* Lange, Dansk Bot. Ark. 5: 37. 1914.

*Pseudohiatula callistosporoides* Sing. spec. nov.

Pileo flavo vel aureo, dein fulvo-maculato, 3—35 m. lato; lamellis flavidis, subconfertis, latiusculis, haud ventricosis, rotundato-adnexis vel subliberis; stipite lamellis concolore, demum fulvescente, 27—29  $\times$  1.5—4 mm., mycelio basali pallido; carne subconcolori, carnosa vel carnosio-subgelatinosa, haud revivescente, immutabili, inodora. Sporis 6.8—8  $\times$  3.5—4.2  $\mu$ , hyalinis, dein rubido-fulvescentibus in membranibus aut in parte interna aut in ambobus, levibus, inamyloideis, depressione suprahilari destitutis; basidiis tetrasporis, 20—21  $\times$  4.5—5  $\mu$ ; cystidiis numerosis eis epicutis pilei analogis; epicute pilei hymeniformi, ex elementis clavato-subvesiculososis (20—21  $\times$  9—11  $\mu$ ) et dermatocystidiis ventricosis, ampullaceis, plerumque rubidofulvescentibus in KOH et in ammoniaco sed flavidis in acidis (48—94  $\times$  11—17.5  $\mu$ ) efformata; hypodermio gelatinoso hyalino, intertexto tramate hymenophorali laterali; hyphis carnis pilei partim subgelatinosis (praecipue prope lamellas); elementis cystidiformibus ad apicem stipitis hyalinis vel fulvescentibus, 40—75  $\times$  13—14  $\mu$  vel

angustioribus; hyphis in basi subpseudoamyloideis fibulatis. Ad truncum *Alni jorullensis* var. *spachii* in silva montana. San Agustin, Taficillo, Sierra de San Javier, Tucumán, Argentina. Leg. Singer no. T 3196 (LIL), 7-I-1959, typus.

Subtribus *Marasmiinae* Sing. (tribus *Marasmiaceae* Fayod, *Tricholomataceae*). Elementis epicuticularibus, sporis, hyphis haud omnibus inamyloideis formis reductis (hymenophoro carentibus) exceptis. Genus typicum: *Marasmius* Fr.

**Crinipellis micropilus** (Reichardt) Sing. comb. nov.

*Marasmius micropilus* Reichardt, Verh. Zool.-Bot. Ges. Wien 16: 375. 1856.

**Crinipellis pseudoarachnoides** (Dennis) Sing. comb. nov.

*Marasmius pseudoarachnoides* Dennis, Trans. Brit. Myc. Soc. 34: 425. 1951.

**Crinipellis sarmentosus** (Berk.) Sing. comb. nov.

*Marasmius sarmentosus* Berk., Hook. J. Bot. 5: 2. 1846.

**Crinipellis actinophorus** (B. & Br.) Sing. comb. nov.

*Marasmius actinophorus* Berk. & Br., J. Linn. Soc., Bot. 14: 39. 1873.

**Chaetocalathus hispidus** (Mass.) Sing. comb. nov.

*Trogia hispida* Mass., Kew Bull. f. 1901: 162. 1901.

**Lachnella pulchra** (Berk. & Br.) Sing. comb. nov.

*Cyphella pulchra* Berk. & Br., J. Linn. Soc., Bot. 14: 74. 1875.

**Palaeocephala** Sing. gen. nov.

Characteribus eis Gloiocephalae gaudentibus, sed cystidiis hymenialibus destitutae et crinibus carentes fructificationes quae exiguae, foliicolae, palaeotropicae sunt. Species typica: **Palaeocephala cymatelloides** (Dennis & Reid) Sing. comb. nov. = *Marasmius cymatelloides* Dennis & Reid, Kew Bull. f. 1957: 292. 1957.

Subtribus **Myceninae** Sing. subtrib. nov. (*Marasmiaceae* Fayod, *Tricholomataceae*). Sporibus amyloideis vel rarius inamyloideis; hyphis pseudoamyloideis vel inamyloideis; cystidiis praesentibus vel absentibus; semper inamyloideis; epicute numquam pseudoamyloidea; hyphis in formis heterothallicis fibuligeris; tramate haud revivescente; hymenophoro plerumque bene evoluto et lamelloso vel tubuloso. Genus typicum: *Mycena* (Pers. ex Fr.) S. F. Gray.

**Hemimycena rickenii** (A. H. Smith) Sing. comb. nov.

*Mycena rickenii* A. H. Smith, North Amer. Spec. Mycena, p. 158. 1947.

**Hemimycena delicatella** (Peck) Sing. comb. nov.

*Agaricus delicatellus* Peck, Ann. Rep. N. Y. State Mus. 30: 39. 1878.

**Hemimycena angusta** Sing. spec. nov.

Pilei cylindrico albo, 3 mm. alto, 1.5 mm. lato, acute papillato; lamellis minime evolutis; hymenio sublevi, albo; stipite albo, subtili-



ter piloso sub lente ad basin et pilis aliis substrato appressis, haud viscido, filiformi,  $12 \times 0.1$  mm. ad apicem, 0.2 mm. lato ad basin; carne subnulla, inodora. Sporis  $8-8.5 \times 3.3-3.7$   $\mu$ , fusioideis, levibus, inamyloideis, contentu oleaginoso impletis, tenuitunicatis; basidiis clavatis, hyalinis, atrolilaceis in caeruleo cresylico, plerumque tetrasporis,  $16-25 \times 4.8-6.2$   $\mu$ ; cystidiis nullis visis. Tramate hymenophorali subnullo; tramate pilei ex hyphis tenuissimis vel sublatusculis (usque as 12  $\mu$  diam.) densissimis et compacte conglomeratis sed parallelis vel subparallelis, inamyloideis, fibulatis efformato; epicute pilei cutem formante, hyphis 1.5-2  $\mu$  latis, e papilla ad marginem verticaliter subparalleleque decurrentibus, diverticulatis (diverticulis  $1.2-2.8 \times 0.3-0.5$  (1.3)  $\mu$ ), inamyloideis. Ad folia arboris dicotyledonum, probabiliter Ilicis. Bolivia, Yungas, La Paz, Nor-Yungas, Cotapata, 3000 m. alt. 28-I-1956, leg. Singer nol B 631 (LIL) typus. — Haec species inter affines (*H. crispata* etc.) unica insignisque forma pilei et habitatione.

**Hemimycena epibiotica** Sing. spec. nov.

Pileo alutaceo (M & P 11-K-7), subaurantiaco-ochraceo-tincto, campanulato, papillato, glabro. pellucide striato sulcatoque,  $6 \times 6$  mm.; lamellis alutaceis, latioribus quam longis. arcuatis, profunde decurrentibus, distantibus (9-10); stipite alutaceo, glabro, filamentoso, aequali, haud viscido, anguste cavo in maturitate,  $26 \times 0.5-0.8$  mm., ad basin frustulis minimis incrustato sed vix fibrilloso; carne exigua, fragili, inodora. Sporis  $6-8.2 \times 4-5.5$   $\mu$ , ellispoideis, hyalinis, contentu refringente repletis, inamyloideis; basidiis  $25-26 \times 7-7.5$   $\mu$ , tetrasporis; cystidiis cheilocystidiisque nullis. Epicute pilei ex elementis hyphosis elongatis repentibus efformata, appendicibus irregularibus diverticulatis et interdum appendicibus diverticulatione secundaria ornatis; hyphis superficialibus stipitis item ornamentatis vel elementis exiguis ramosis ornatis sed pilis destitutis; hyphis stipitis ab eis pilei haud differentibus, parallelis, interioribus subochraceis (KOH), in pileo latioribus quam in stipite; hypodermio cellulari absente; tramate hymenophorali flavido angustissimo, regulari, ex hyphis mediocriter latis efformato; hyphis omnibus haud gelatinosis, fibulatis, subinamyloideis vel minime pseudoamyloideis. Ad lignum detritum et excrementa insectorum lignivororum in cavitatibus ligni trunci in aquam delapsi. Bolivia, Yungas, La Paz, Nor-Yungas, Charobamba, 13-II-1956. Leg. Singer no. B 1146 (LIL), typus.

**Hemimycena albida** (Murr.) Sing. comb. nov.

*Gymnopus albidus* Murr., Bull. Torr. Bot. Cl. 66: 32. 1939.

**Hemimycena subnigricans** (Murr.) Sing. comb. nov.

*Marasmius subnigricans* Murr., Bull. Torr. Bot. Cl. 67: 152. 1940.

**Hemimycena longicystis** Heim spec. nov.

Pileo albo, semiglobose-campanulato, 6-20 mm. lato. Hymenophoro separabili, lamellis distantibus (e. gr. 12), undulatis, venoso-

anastomosantibus, adnato-decurrentibus; stipite longo. Sporis 12.5—15.5 × 4.5  $\mu$ , lacrimiformibus-elongatis, irregularibus; basidiis tetrasporis; cystidiis usque ad 100  $\mu$  longis, apice longo cylindraceo angusto praeditis; cuticula pilei paullum differentiata. In olla bambusae et ad terram circumdantem Leg. Stehle no. 5 (P), 28-V-1936. Etiam a cl. Dennis in ins. Trinidad nec non a me in Bolivia collectus fuit. Ab auctore descriptione gallica sub *Mycena* (*Hemimycena*) publicata et Dennisio *Collybia*! Typus est collectio boliviana (Charobamba, Nor-Yungas, 30-I-1956, Singer B 707 (LIL)).

**Hemimycena truncicola** Sing. spec. nov.

Pileo albo, rugoso-sulcato, umbilicato vel subumbilicato, campanulato-subpapillato in juvenilibus, dein convexulo, glabro; lamellis bene evolutis, albis, subdecurrentibus, distantibus; stipite albo, fuscescente et umbrino basin versus, levi, minute pubescente, glabrescente, ad basin stellatim fibrilloso, 10—40 × 0.2—0.3 mm. Carne exigua, inodora. Sporis 7—9.5 × 3.5—5 (5.5)  $\mu$ , hyalinis, levibus, contentu oleaginoso abundante impletis, inamyloideis; basidiis tetrasporis, 18—28 × 7—7.5  $\mu$ ; cystidiis nullis; cheilocystidiis numerosis, cylindraceo-filamentosis vel irregulariter tumescentibus, excrescentiis praeditis vel diverticulatis vel furcatis, interdum latera lamellarum invadentibus; pubescentia stipitis pilis erectis consistente; pilis 27—63 × 4—8.5  $\mu$ , plerumque subulatis, subacutis vel acutis, interdum subdiverticulatis. Hyphis pilei hymenophorice inamyloideis, stipitis partim subrubidis iodi ope (subpseudoamyloideis), constanter fibulatis. Elementis epicutis pilei diverticulatis. Ad lignum truncorum Dicotyledonum partim adustum. Argentina, Tucumán, Ciudad Universitaria (Sierra de San Javier), 12-I-1957. Leg. Singer no. T 2872 (LIL), typus.

**Hemimycena angustispora** (Josserand ex Orton) Sing.

*Mycena angustispora* Josserand ex Orton, Trans. Brit. Myc. Soc. 43: 305. 1960.

**Mycena sotae** Sing. spec. nov.

Pileo albo vel griseo-fusco, partim caeruleo, convexo, dein centro appanato, papillato vel depapillato, sed papilla semper paullum manifesta, subglabro, margine striato, 6—7.5 mm. lato; lamellis albis vel albidis, confertiusculis, liberis; stipite pileo concolori, pubescente vel piloso debiliter ad apicem, fortiter ad basin, insititio ad discum latum sulcatum pulverulentum griseum basalem; carne tenui, inodora, concolori superficiebus. Sporis 6.8—8.2 × 5—5.5  $\mu$ , ellipsoideis, raro oblongis, levibus, tarde leviterque amyloideis, hyalinis in KOH; basidiis 15—17 × 6—6.5  $\mu$ , tetrasporis; cystidiis nullis; cheilocystidiis 23—35 × 15—18  $\mu$ , echinulatis, vesiculososis (setulis 1—1.5  $\mu$  altis); epicute pilei consistente sphaerocystibus quae saepe separantur cellulis hypodermii atque inter se, demum multis liberis, echinulatis ut cheilocystidia, globosis vel rotundato-triangularibus, succo fusco

impletis; pilis stipitis ventricosis in tertia inferiore, ampullaceis, ad apicem longe cylindraceis. hyalinis vel fuscis, echinulatis,  $55-310 \times 6-20 \mu$ , apex  $3.5-7 \mu$  diam.; hyphis haud gelatinosis, fibuligeris. Ad lignum sub arbustis Dicotyledonibus. Bolivia, Yungas, La Paz, Nor-Yungas. Las Ciénagas, 10-II-1956. Leg. Singer & E. R. de la Sota no B 1073 (LIL), typus. — Species insignis correlatione characterum *M. osmundicolae* et *M. discopodae*; in exsiccatore brunnescit! Species collectorum et amico Elias Ramón de la Sota, botanico tucumanensi dedicata.

***Mycena squamulosa*** Sing. spec. nov.

Pileo cylindrico-oblongo ut in *Coprinis*, dein subexpanso, campanulato papillatoque, glabro sed disco furfuraceo Coprinorum modo, longe subtiliter pectinato, mox rimoso Inocybarum modo, margin frequenter crenato, sicco, brunneo (inter „Cookie“ et „Malay“ M & P),  $15-20$  mm. lato; lamellis albis, ascendentibus vel subascendentibus, confertis dein subconfertis, liberis; stipite albo, pruinato, aequali,  $20-30 \times 1.5-2$  mm.; carne alba, tenuissima in pileo, inodora. Sporibus  $7.5-9 \times 3.7-4.5 \mu$ , caeruleo-amyloideis, hyalinis in KOH, cylindraceis vel ellipsoideo-oblongis vel ellipsoideis, interdum subcurvulis, tenuitunicatis, levibus, guttula una vel guttulis duobus impletis; basidiis  $15-18.7 \times 5.5-7.5 \mu$ , exiguis angustisque, setrigmatibus uno vel duobus vel tribus ornatis, ad basin fibuligeris vel defibulatis; cystidiis nullis, sed cheilocystidiis numerosis (et uno alioque ad latera lamellarum scandente, ventricosus in parte inferiore, apice appendice longo filamentoso acuto vel obtuso ornatis, numerosis vel dense congregatis, hyalinis,  $27-65 \times 5-7 \mu$ ; subhymenio ex elementis minutissimis consistente; tramate hymenophorali regulari, ex elementis voluminosis longis efformato; epicute pilei ex elementis repentibus catenatis (cellulis longis cylindraceis et aliis subsodiametricis) levibus consistente, cellulis terminalibus plerumque breviter cylindraceis, ampullaceis, capitatis, clavatis etc. et ibidem parietibus interdum crassitunicatis observatis; hypodermio ex elementis repentibus sat voluminosis elongatis efformato; hyphis omnibus carnis et hymenophori pseudoamyloideis, fibulis sparsis munitis. Ad *Tipuanae* corticem vivae gregatim. Argentina, Tucumán, Capital, 4-I-1959. Leg. Singer no. T 3175 (LIL), typus. — An cum *Leptomycece squamulosa* Mont. identica species? Huius typus in Herbario P haud conservatur, sed facile hic spectat.

***Mycena alexandri*** Sing. spec. nov.

*Mycena psammicola* (Berk. & Broome) Sacc. sensu Alexander H. Smith non Berk. & Br.

Carpophoris galerinoideis (colore *Galerinae leavis*); pseudo-cystidiis destitutis, fragilibus; cheilocystidiis echinulatis. Ad acus pini. USA, Michigan, 23-IX-1911 leg. Kauffman (MICH) est typus (*M. psammicola*, det. Smith).

***Mycena palmivora* Sing. spec. nov.**

Pileo centro avellaneo, margine roseolo, umbilicato, convexo, margine substriato, neque hygrophano nec viscido, pubescente in juvenilibus, glabrescente in maturis, margine stricto in juvenilibus, 22 mm. lato vel minore; lamellis candidis, subventricosis, sinuatis, subdistantibus, adnexed, medioeriter latis vel latiusculis; stipite albo, pubescente praesertim prope basin, ad basin albo-strigoso, aequali,  $30 \times 2.2$  mm. Sporis  $6-7.8 \times 4.3-6$   $\mu$ , levibus, ellipsoideis, amyloideis; basidiis tetrasporis;  $38-42 \times 7.8-8.3$   $\mu$ ; pro ratione longis; cystidiis fusiformi-ventricosis, rarius late cylindraceis vel ampullaceis, hyalinis,  $47-68 \times 9-14.3$   $\mu$ ; epicute pilei e dermatocystidiis densissimis sed haud stratum hymeniforme formantibus efformata; dermatocystidiis pilei cylindraceis, rarius ampullaceis, hyalinis, levibus, raro basi ventricosa praeditis,  $8-47 \times 3.5-5$   $\mu$ ; hypodermio cutem efformante, ex hyphis levibus elongatis usque ad 15  $\mu$  latis efformatam; hyphis omnibus pseudoamyloideis, fibulatis; hymenio in solutione ammoniaci crystalli Conocybis tenerae modo formante. Ad folia delapsa Sabali palmetto in silva. — USA, Florida, Highlands Hammock State Park, 11-VIII-1942. Leg. Singer no. F 158 (F sub nom. *Mycena sabali* Murr.), typus. — Siccus in herbario totus cinereus, pruinosis.

***Mycena viridula* (Berk. & Cooke) Sing. comb. nov.**

*Laschia viridula* Berk. & Cooke, Grevillea 19: 105. 1891.

***Mycena holoporphyra* (Berk. & Curt.) Sing. comb. nov.**

*Agaricus holoporphyrus* Berk. & Curt., J. Linn. Soc., Bot. 10: 284. 1869.

***Mycena griseoreta* Sing. spc. nov.**

Pileo atropurpureo („Maracaibo“ M&P), conico-expanso, interdum subumbonato, subumbilicato vel truncato in centro, longe radiatim pellucide striato, glabro, 10–17 mm. lato; lamellis griseis („Kaffa“ M&P), sed albis in vicinitate acieis, subangustis, confertis vel subconfertis, dense intervenosis, adnatis, dente decurrente; stipite lamellis concolori, obscuriore ad apicem, ad basin albido-pubescente,  $48-51 \times 1-1.5$  mm. sed ad apicem usque ad 3.5 mm. incrassato; carne odore raphanaceo praedita. Sporis  $7-8.5 \times 4.5-4.8$   $\mu$ , hyalinis, levibus, tarde subtiliter amyloideis; basidiis  $23-33 \times 7-8.3$   $\mu$ , tetrasporis; cystidiis nullis; cheilocystidiis  $30-45 \times 9-13$   $\mu$ , ventricosis, ampullaceis, ampullaceo-subcapitatis vel gradatim apicem versus attenuatis, obtusis, raro furcatis; epicute ex hyphis hyalinis levibus, filamentosis, repentibus efformata, tenui; hyphis hypodermii in ammoniaci lilaceo-amethystinis, voluminoso-inflatis, frequenter brevissimis, cutem formantibus; hyphis omnibus fortiter pseudoamyloideis, fibulatis. Ad folia dejecta in plantatione Coffeae arabicae. Bolivia, Yungas, La Paz, Nor-Yungas, Coroico, 27-I-1956. Leg. Singer no. B 566 (LIL), typus.



**Mycena coccinea** (Murr.) Sing. comb. nov.

*Omphalina coccinea* Murr., North Amer. Flora 9: 350. 1916.

**Mycena drepanocladi** (Sing.) Sing. comb. nov.

*Hemimycena drepanocladii* Sing., Ann. Mycol. 41: 123. 1943.

*Mycena* sectio **Sacchariferae** Kühner sect. nov. Species disco basali destitutae aut praeditae; pileo haud gelatinoso in cuticula, epicute cum multis cellulis inflato-vesiculosus diverticulatis et frequenter acanthophysoideis. Typus: *M. tenerrima* (Berk.) Sacc.

*Mycena* sectio **Citricolors** Sing. sect. nov. Stipite disco basali destituto, insititio vel subsinititio; elementis epicutis pilei diverticulatis; tramate pilei gelatinoso cheilocystidiis echinulatis (acanthophysoideis), hyphis gelatinosis inamyloideis; hyphis non gelatinosis fortiter pseudoamyloideis; carpophoris omphalioideis; sporis ellipsoideis vel oblongis, minoribus, inamyloideis. Generatio asexualis ad folia vigentia ubi stilboideae fructificationes (propagula) formantur. Typus: *M. citricolor* (Berk. & Curt.) Sacc.

*Mycena* sectio **Radiatae** Sing. sect. nov. Stipite disco basali destituto, insititio vel subsinititio, pigmento caeruleo nullo; epicute pilis longis ornata vel catenulis elementorum breviusculorum consistente; cheilocystidiis absentibus vel haud echinulatis; hyphis saepe fibulis inconstantibus praeditis; carpophoris mycenoideis vel mycenoideo-pluteoideis; hyphis pseudoamyloideis vel inamyloideis; sporis amyloideis, ellipsoideis vel cylindraceis. Ad ligna vel humum lignosum in tropicis vel subtropicis. Typus: *M. radiata* (Dennis) Sing.

*Mycena* subgen. *Glutinipes* subsectio **Fuliginellae** A. H. Smith (primitus ut sectio sed sine diagnose). Pileo viscido; species solitariae vel gregariae humicolae. Typus subsectionis: *M. vulgaris* (Pers. ex Fr.) Kummer.

*Mycena* subgen. *Glutinipes* subsect. **Caespitosae** A. H. Smith (primitus ut sectio sed sine diagnose) Pileo viscido; species caespitosae vel dense gregariae, lignicolae. Typus subsectionis: *M. texensis* A. H. Smith.

*Mycena* sect. **Aeiculae** Kühner (ut subsectio sed sine diagnose latina). Stipite structura sua a tramate pilei zona heterogenea separato; carpophoris mycenoideis; lamellis primum plus minusve ascendentibus, adnatis vel dente decurrentibus, haud ventricosus; pigmento microscopice flavo, macroscopice aurantiaco vel rubro; cheilocystidiis et plerumque etiam plerocystidiis praesentibus; hyphis epicuticularibus pilei diverticulatis; tramate hymenophorali subregulari; hyphis oleiferis laticiferis similibus nullis vel paucis et exiguis. Typus sectionis: *M. aeicula* (Schaeff. ex Fr.) Kummer.

*Mycena* sect. **Hiemales** subsect. **Insititopedes** Sing. subsect. nov. A stirpibus sectionis Hiemalium ceteris differt stipite insititio radicellis destituto nec pseudorhiza praedito. Typus subsectionis: *M. tucumanensis* Sing.

*Hydropus* sectio **Mycenoides** Sing. sect. nov. Sporibus inamyloideis a

sectione *Hydropode* differt. Typus sectionis: *H. mycenoides* (Dennis) Sing.

***Hydropus mycenoides*** (Dennis) Sing. comb. nov.

*Collybia mycenoides* Dennis, Trans. Brit. Mycol. Soc. **34**: 439. 1951.

***Hydropus aristoteliae*** (Sing.) Sing. comb. nov.

*Mycenella aristoteliae* Sing., Arkiv för Botanik 2. **4**: 381. 1959.

*Hydropus* sect. **Floccipedes** Kühn. (ut subsectio *Mycenae* sine diagnose latina) ex Sing. sect. nov. Sporis amyloideis vel inamyloideis; hyphis debiliter pseudoamyloideis; hyphis pilei epicutis levibus, elongatis, pigmento frequenter intracellulari praesente in epicute; pleurocystidiis et cheilocystidiis praesentibus. Typus sectionis: *M. floccipes* (Fr.) Kühner.

***Hydropus scabripes*** (Murr.) Sing. comb. nov.

*Prunulus scabripes* Murr., North American Flora **9**: 331. 1916.

*Hydropus* sect. **Floccipedes** subsectio **Spuriae** Kühn. (ut *Mycena* sect. *Mycenopsis* subsect. *Spuriae*, Enc. Mycol. **10**: 515. 1938) Characteribus sectionis **Floccipedes**; sporis amyloideis. Typus subsectionis: *H. scabripes* (Murr.) Sing.

*Hydropus* sect. **Floccipedes** subsectio **Floccipedes** Kühn. (ut *Mycena* subgen. *Paramycena-Hemimycena* sect. *Genuinae* subsectio **Floccipedes**, l. c. p. 538) Characteribus sectionis **Floccipedes**; sporis inamyloideis. Typus subsectionis: *H. floccipes* (Fr.) Sing.

***Hydropus floccipes*** (Fr.) Sing. comb. nov.

*Agaricus floccipes* Fr., Epicrisis p. 87. 1838 (sensu Kauffman).

***Hydropus subalpina*** (Höhnelt) Sing. comb. nov.

*Mycena subalpina* Höhnelt, Fragm. XV, p. 275. 1913 (sensu Kühner).

*Fayodia* subgenus **Myxomphalia** Kühn. (ut divisio *Spuriarum* (*Mycena*) sine hierarchia definita, Enc. Mycol. **10**: 535. 1938) ex Sing. Pleurocystidiis frequenter praesentibus; cheilocystidiis numerosis; sporis subcrassitunicatis sed apparenter membrana homogenea praeditis, levibus vel subsperulis; lamellis decurrentibus. Typus subgeneris *F. maura* (Fr.) Sing.

*Fayodia* subgenus **Heterosporula** Sing. subgen. nov. Sporis membrana heterogenea vel haud levi praeditis, inamyloideis. Typus subgeneris: *F. pseudoclusilis* (Josserand & Konrad) Sing.

***Fayodia pseudoclusilis*** (Josserand & Konrad) Sing. comb. nov.

*Collybia pseudoclusilis* Josserand & Konrad, Bull. Soc. Linn. Lyon 1931, p. 4 (repr.).

***Baeospora pleurotoides*** (Dennis) Sing. comb. nov.

*Collybia pleurotoides* Dennis, Trans. Brit. Myc. Soc. **34**: 455. 1951.

***Baeospora brunneipes*** (Sing.) Sing. comb. nov.

*Heimiomyces brunneipes* Sing. in Sing. & Digilio, Lilloa **25**: 242. 1951 (publ. 1952).

**Baeospora pruinatipes** (Sing.) Sing. comb. nov.

*Heimiomyces pruinatipes* Sing. in Sing. & Digilio, l. c. p. 243.

*Amanita* sect. **Ovigerae** Sing. sect. nov. A sectione *Vaginatae* differt sporis ovoideis vel ellipsoideis nec globosis nec subglobosis. Typus sectionis: *A. biovigera* Sing.

*Amanita* sect. **Roanokenses** Sing. sect. nov. Pileo stipiteque pulvere detersili obtectis. Typus sectionis: *A. roanokensis* Coker.

**Volvariella fibrillosa** (Bres. ex Rick, Broteria **17**: 117, 1919; Lilloa **3**: 451, 1938) Sing. comb. nov.

**Volvariella pseudovolva** (Berk. & Br.) Sing. co, b. nov.

*Agaricus pseudovolva* Berk. & Br., J. Linn. Soc., Bot. **11**: 530. 1871.

**Volvariella villosovolva** (Lloyd) Sing.

*Volvaria villosovolva* Lloyd, Mycol. Notes 31. 1899.

**Volvariella cyenopotami(a)** (Berk.) Sing.

*Agaricus cyenopotami(a)* J. Linn. Soc., Bot. **18**: 389. 1881.

**Volvariella californica** (Earle) Sing.

*Locellina californica* Earle, Bull. N. Y. Bot. Gard. 1904: 299. 1904.

**Volvariella arenaria** (Pat.) Sing.

*Pluteus arenarius* Pat., Rev. Mycol. **13**: 136. 1891.

**Macrolepiota zeyheri** (Fr.) Sing. comb. nov.

*Agaricus zeyheri* Fr., Fungi natalenses, p. 2. 1848.

**Leucocoprinus birnbaumii** (Corda) Sing. comb. nov.

*Agaricus birnbaumii* Corda, Icon. Fung. **3**: 48. 1839.

*Coprinus* sect. *Coprinus* subsect. **Alachuani** Sing. ex Sing. sect. nov. Hyphis epicuticularibus pilei ramoso-dichophysatoideis. Typus sub-sectionis *Coprinus alachuanus* Murr.

Subfam. **Psathyrelloideae** Sing. Lamellis in sectione transversali subulatis. Superficie pilei haud pectinato-fissa; sporis in acido sulphurico concentrato pallide lividis. Genus typicum: *Psathyrella* (Fr.) Quél.

*Psathyrella* subgen. **Psalliotina** (Velen.) Sing. stat. nov.

*Psalliotina* Velen. (ut genus), Nov. Mycol. p. 155. 1939.

*Psathyrella* sect. **Candolleanae** Romagnesi sect. nov. Sporis minoribus 10  $\mu$ ; pleurocystidiis nullis; tramate hymenophorali hyalino; carpophoris carnosio-fragilibus, velo abundante, frequenter caespitose crescentes. Species typica: *P. candolleana* (Fr.) Maire in Maire & Werner.

*Psathyrella* sect. **Fatae** Romagnesi sect. nov. Sporis minoribus 11  $\mu$ ; pleurocystidiis abundantibus, raro mucronatis; carne demum hyalina. Species typica: *P. pygmaea* (Quél.) Sing.

**Psathyrella muricellata** (Sing.) Sing. comb. nov.

*Psathyra muricellata* Sing., Notul. Syst. Inst. Bot. USSR **10**—**12**: 13. 1938.

*Psathyrella* sect. **Typhicolae** Romagnesi sect. nov. Sporis 10  $\mu$  maiori-

bus; pleurocystidiis nulis; tramate hymenophorali colorato pigmento membranali; lamellis subanguste adnexis. Species typica: *P. typhae* (Kalchbr.) Pearson & Dennis.

*Psathyrella* sect. **Hydrophilae** Romagnesi sect. nov. Sporis 11.5  $\mu$  minoribus frequenter sat minusculis; cystidiis utriformibus; tramate colorato. Species typica: *P. hydrophila* (Bull. ex Mérat) Maire in Maire & Werner.

*Psathyrella* sect. **Frustulentae** sect. nov. (Romagnesi ut subsectio). Sporis 6—11.5  $\mu$  longis; pleurocystidiis numerosis; carpophoris minoribus et velo fugaci. Species typica: *P. frustulenta* (Fr.) A. H. Smith sensu A. H. Smith.

*Psathyrella* sect. **Microrhizae** Romagnesi sect. nov. Characteribus subgeneris *Psathyrella*; tramate hymenophorali colorato; stipite pseudorhiza praedito; velo plus minusve evoluto. Species typica: *P. microrhiza* (Lasch) Konrad & Maublanc.

*Psathyrella* sect. **Atomatae** Romagnesi sect. nov. Characteribus subgeneris *Psathyrellae*; differt a sectione praecedente absentia pseudorhizae. Species typica: *P. prona* (Fr.) Gillet sensu Ricken.

*Psathyrella* sect. **Subatratae** Romagnesi sect. nov. Characteribus *Psathyrellae* (subgeneris) sed velo nullo. Species typica: *P. subatrata* (Batsch ex Fr.) Quéf.

*Psathyrella* subgenus **Cystopsathyra** Sing. subgen. nov. Vestimento velari pilei et stipitis pulverulento-farinoso, consistente e sphaerocystibus; pleurocystidiis praesentibus, late rotundatis, non muricatis. Species typica: *P. kellermanii* (Peck) Sing.

Subfamilia **Panaeoloideae** Sing. subfam. nov. Cortinacearum (Ann. Myc. 34: 339. 1936 sine diagnose latina). Ab aliis subfamiliis Coprinacearum differt sporis atris vel fuscis in acido sulphurico haud pallescentibus Genus typicum: *Panaeolus* (Fr.) Quéf.

**Conocybe siennophylla** (Berk. & Br.) Sing. comb. nov.

*Agaricus siennophyllus* Berk. & Br., J. Linn. Soc., Bot. 11: 545. 1871.

**Conocybe martiana** (Berk. & Curt.) Sing. comb. nov.

*Agaricus martianus* Berk. & Curt., J. Linn. Soc., Bot. 10: 291. 1869.

**Conocybe umbellula** (Mont.) Sing. comb. nov.

*Agaricus umbellula* Mont., Ann. Sc. Nat. IV. 1: 100. 1854.

*Conocybe* sect. **Mixtae** Kühn. (primitus subsectio!) ex Sing. sect. nov. Dermato-cystidiis stipitis et capitatis et non capitatis praesentibus; pilis stipitis etiam numerosis. Typus sectionis: *C. cryptocystis* (Atk.) Sing.

*Conocybe* sect. **Pilosellae** Kühn (primitus ut subsectio!) ex Sing. sect. nov. Stipite haud toto albo; hyphis pilosis numerosis; dermatocystidiis capitatis in stipite nullis. Typus sectionis: *C. pilosella* (Pers. ex Fr) Kühner.



*Conocybe* sect. **Candidae** Kühner (primitus subsectio!). Stipite candido-dermatocystidiis stipitis nullis capitatis; hyphis piliformibus numerosis; pileo frequenter striato; pseudoparaphysibus frequenter numerosis. Typus sectionis: *C. lateritia* (Fr.) Kühner sensu Kühner.

*Pholiotina* sect. **Piliferae** Kühner (ut sectio generis *Conocybe* sed sine diagnose latina). A *Pholiotina* sectione *Pholiotina* differt absentia veli. Typus sectionis: *P. pygmaeoaffinis* (Fr. sensu Kühner) Sing.

**Bolbitius tucumanensis** Sing. spec. nov.

Pileo atrobrunneo dein griseolocastaneo („broncho“ dein „auburn“ M&P), disco albicante, glutinoso, longe denseque pectinato Coprinorum modo, campanulato margine stricto et stipite applicato, dein expanso, demum applanato, glabro, 36—44 mm. lato; lamellis brunneis (inter „Cochin“ et „burnt umber“ M&P), angustis, adnexas vel liberis, confertis vel confertissimis; sporis in massa brunneo-ferrugineis („cocoa“, in massa densa copiosa inter „Mandalay“ et „burnt umber“ M&P); stipite albo, floccoso, ad apicem attenuato, 75—96 × 4—5 mm.; carne fragili, alba, tñui, inodora. Sporis 18—20.7 × 7—8.7 μ, ferrugineis, levibus, crassitunicatis, membrana complexa et poro germinativo praeditis, sporis *Conocybis* analogis; basidiis bi-vel tetrasporis, 18—21.5 × 14 μ; pseudoparaphysibus 13.7—18.8 × 12—14.5 μ, vesiculoso-clavatis, numerosis; cheilocystidiis e. gr. 25 × 12.7 μ, ventricosis; hyphis defibulatis. Ad fimum equinum gregatim. Argentina, Tucuman, Quebrada de la Higuera, 25-I-1955, leg. Singer no. T 2081a (LIL), typus.

**Agrocybe underwoodii** (Murr.) Sing. comb. nov.

*Naucoria underwoodii* Murr., Mycologia 4: 80. 1912.

*Agrocybe* sect. **Microspora** Sing. (1936) ex Sing. sect. nov. Velo nullo, cystidiis ad latera lamellarum praesentibus aut sporis 10 μ minoribus. Typus sectionis: *A. tuberosa* (Henn.) Sing.

*Agrocybe* subgen **Aporus** Sing. (1936) ex Sing. subg. nov. Poro germinativo frequenter indistincto vel nullo aut sporis, si porus praesens, haud manifeste truncatis. Typus subgeneric: *Agrocybe aegerita* (Brig.) Sing. Subfamilia **Stropharioideae** Sing. (fam. *Strophariaceae*). Sporis in massa atrolilaceis, fusco-sepiaceis; sporis siccis s. l. saepe olivaceis vel linea annulari endosporiali castaneo-ruba ornatis, interdum lentiformibus. Genus typicum: *Stropharia* (Fr.) Quél.

*Stropharia* sect. **Stercophila** (Romagnesi ex) Sing. (primitus ut genus sed sine diagnose). Pileo stipiteque glutinosis e velo toto glutinoso-sporis grandibus. Typus sectionis *S. semiglobata* (Batsch ex Fr.) Quél.

**Stropharia pendolae** (Speg.) Sing. comb. nov.

*Pilosace pendolae* Speg. Bol. Acad. Cienc. Cord. 28: 332. 1926.

*Ptilocybe* sect. **Deconica** (W. G. Smith) Sing. stat. nov.

*Deconica* W. G. Smith (ut subgenus *Agarici*), Clavis Agaricorum p. 23. 1870.

**Psilocybe crobula** (Fr.) Morten Lange ex Sing. comb. nov.

*Agaricus crobulus* Fr., Epir. p. 199. 1838.

**Psilocybe goniospora** (Berk. & Br.) Sing.

*Agaricus goniosporus* Berk. & Br., J. Linn. Soc. Bot. 11: 541. 1871.

**Melanotus ridleyi** (Mass.) Sing.

*Crepidotus ridleyi* Mass., Bull. Misc. Inf. 1899: 169. 1899.

**Melanotus cassiaecolor** (Berk.) Sing. comb. nov.

*Agaricus cassiaecolor* Berk., Hooker, Fl. Tasmania 2: 246. 1860.

*Pholiota* subgenus **Hemipholiota** Sing. ex Sing. subgenus nov. Pleurocystidiis nullis vel sparsis; chrysocystidiis nullis; typus subgeneris: *Pholiota destruens* (Brond.) Gillet.

*Pholiota* sect. **Alboerenulatae** Sing. ex Sing. sect. nov. Characteribus subgeneris *Hemipholiotae*; pileo viscido, squamoso. Typus sectionis: *Pholiota alboerenulata* (Peck) Sacc.

**Pholiota heliocaes** (Berk. & Br.) Sing. comb. nov.

*Agaricus heliocaes* Berk. & Br., J. Linn. Soc., Bot. 11: 544. 1871.

*Pholiota* sect. **Sordidae** Sing. sect. nov. Characteribus *Hemipholiotae*; pileo primum pellicula gelatinosa tecto, hygrophano; sporis poro minuto germinativo instructis sed apice rotundatis. Typus sectionis: *Flammula sordida* Kühn. in Kühn. & Romagnesi nom. nud. = *Pholiota oedipus* (Cooke) Orton.

*Pholiota* sectio **Subflammantes** Sing. sect. nov. Pileo superficie subviscida vel lubrica gaudente, floccis magnis mollibus squarroso, sed floccis pallide coloratis fugacibus ita ut pileus facile denudetur. Typus sectionis: *Pholiota subflammans* (Speg.) Sacc.

*Pholiota* sect. **Sericellae** (Fr.) Sing.

*Agaricus* tribus *Flammula* sect. *Sericellae* Fr., Hymen. Europ. p. 252. 1874.

Typus sectionis: *Agaricus scambus* Fr.

*Pholiota* sect. **Privignae** Sing. sect. nov. Pileo subviscido vel viscido; pellicula gelatinosa crassa praedito; pleurocystidiis ad typum metuloideum vergentibus sed interdum contentu chrysocystidiali impletis. Typus sectionis: *Pholiota privigna* (Speg.) Sing.

**Pholiota anomala** (A. H. Smith) Sing. comb. nov.

*Hypholoma anomalum* A. H. Smith. Mycologia 33: 2, 1941.

**Pleuroflammula hibernianus** (Pears. & Dennis in Pearson) Sing.

*Crepidotus hibernianus* Pearson & Dennis in Pearson, Trans. Brit. Mycol. Cl. 32: 268. 1949.

*Inocybe* sect. **Cervicolores** Kühner & Romagnesi ex Sing. Sporibus levibus; metuloideis nullis; pileo squamoso vel tomentososo; carne rubescente; odore aromatico vel fructuum vel terreo. Species typica: *Inocybe cervicolor* (Pers.) Quéf.

**Alnicola mirabilis** (Atk.) Sing. comb. nov. 1).

*Galerula mirabilis* Atk., Proceedings Amer. Phil. Soc. 57: 3. 1918.

**Alnicola inculta** (Peck) Sing. comb. nov.<sup>1)</sup>

*Galera inculta* Peck, Ann. Rep. N. Y. St. Mus. 41:69. 1888.  
*Alnicola* sect. **Submelinoideae** Sing. ex Sing. sect. nov. Cheilocystidiis plerumque crassiusculis haud attenuatis vel acutis ad apicem. Typus sectionis: *A. submelinoides* Kühner.

**Alnicola velutina** (Murr.) Sing. comb. nov.

*Naucoria velutina* Murr. North Amer. Flora 10: 177. 1917.

**Simocybe tiliophila** (Peck) Sing. comb. nov.

*Agaricus tiliophilus* Peck, Ann. Rep. N. Y. State Mus. 35: 133. 1884.

**Simocybe reductoaffinis** Sing. spec. nov.

Pileo pallide sordideque brunneolo, pellucide striato, hygrophano, convexo, obtuso, 12 mm. lato; lamellis pileo concoloribus, subventricosis, confertiusculis, adnatis vel subsinuato-adnatis; stipite pallido, glabro, aequali, 19×1.5 mm.; carne pallida vel subconcolori super-

<sup>1)</sup> I cannot understand the statement by Dennis, Orton & Hora (Trans. Brit. Myc. Soc., Suppl. p. 200 under "*mirabilis*" and Trans. Brit. Myc. Soc. 43: 319. 1960, by Orton) in which it is said that "Singer ... maintains that *G. mirabilis* is the same as *N. alnetorum*; I think this unlikely, however, on account of ..." etc. This equation is not only unlikely but erroneous and, what is more, has never been "maintained" by me. I identified (Singer in Sydowia 9: 406) *Alnicola mirabilis* (Atk.) Sing. with *A. bohémica* (Velen.) Sing. It is *G. inculta* Peck which I identified (l. c.) with *A. alnetorum* (Maire in Kühner) Romagnesi. These two statements were obviously confused by Orton. It is quite another question whether within the species earlier defined by Kühner and keyed out by Romagnesi as *A. bohémica* respectively *N. bohémica*, there are one, two or three species hidden. In view of the latest notes published by Kühner & Romagnesi (Bull. Soc. Nat. Oyon. 10—11: 9 11. 1957) one would be inclined to agree with Orton that the *A. bohémica* in its original sense should be separated from the forms with broadly clavate-capitate cheilocystidia. I do not wish to enter this discussion because I have never studied the form with the narrow cheilocystidia. However, if the forms now distinguished from *A. bohémica* as *Naucoria langei* forma *tetraspora* Kühner in Kühner & Romagnesi and as *Naucoria salicis* Orton (apparently both once described by Lange under the common denomination *Naucoria macrospora* Lange non Patouillard & Doassans) are actually specifically different as suggested by Orton (not by Kühner & Romagnesi, l. c.), *Naucoria salicis* Orton must be called *Alnicola mirabilis* (Atk.) Sing. If they are identical and merely represent a two and a four-spored race of the same species the species must be known as *Alnicola mirabilis* (Atk.) Sing. whereby the two-spored race is the type form. I know the four-spored form only from one American collection (received from A. H. Smith under the name *Naucoria hamadryas* and commented on in Rev. Myc. 5: 11. 1940) which is evidently quite identical with *N. langei* Kühner. The other (numerous) collections I have made or received are bisporous and obviously the same as *N. salicis* Orton; with the exception of one collection from USSR, Tataria, Kazan, leg. Vasilieva (LE) which agreed in all regards with A. H. Smith's specimens. My two-spored material: Holland: Reijnders no. 2254 (LIL, L); Tristan da Cunha (K); U.S.A., N. Y. (type of *G. mirabilis*); Mich.: Singer N 744 (F) and N 948 (F). It should be noted that N 744 was collected definitely under *Alnus*. It appears therefore that *A. mirabilis*, two-spored form, not grow exclusively under *Salix* as suggested by Orton.

ficiebus, subinodora. Sporis  $6.8-7.8 \times 5-5.5 \mu$ , levibus, sublato ellipsoideis, haud reniformibus vel phaseoliformibus, poro germinativo destitutis; basidiis tetrasporis,  $15.5-19.5 \times 7.2 \mu$ ; cystidiis nullis; cheilocystidiis cc.  $32 \mu$  longis, versiformibus sed semper rotundatis in apice, frequenter ventricosis in parte superiore aut capitatis aut subcapitatis, interdum semel bisve constrictis vel breviter mucronatulis,  $7.8-12 \mu$  latis, pedicellatis; epicute e dermatocystidiis haud in hymenium condensatis consistente, e cellula subsodiometrica ortis e qua frequenter nascuntur sine septa limitante, interdum basi incrassata sed generatum filiformibus vel filiformi-subcapitatis,  $25-37.5 \times 7-9.3 \mu$ , cellulis subjacentibus e. gr.  $14-20 \mu$  in diam. et si cum dermatocystidiis confluent, cellula integra  $35-60 \mu$  longa; hyphis fibuligeris. Ad lignum putridissimum in silva subtropicali. U.S.A., Florida, Dade Co., Brickel's Hammock, leg. Singer no. F 1618 (F), Typus.

**Simocybe rubi** (Berk.) Sing. comb. nov.

*Agaricus rubi* Berk., in Hooker, English Flora 5 (2): 102. 1836.

**Simocybe junquillea** Sing. spec. nov.

Pileo luteo ("jonquil" + "deep chrome" M & P), dein pallidior aurantiaco-flavido, levi, subglabro, dein sicco irregulariter punctato, subpruinato in juvenilibus, convexo dein applatato, interdum paullum concavo in vetustis,  $17-22$  mm lato; lamellis ochreo-grisoleis (inter "Longbeach" et "fallow" M & P), demum brunneolis, subdistantibus, sublatis, adnatis; stipite albo, pubescente vel hirsuto e tomento basali pruinato ad apicem extremum, in zona intermedia glabro, et ad basin ipsam tomentoso-hirsuto; curvato, excentrico vel sublaterali sed haud continuo cum margine pilei,  $5-10 \times 2-4$  mm.; mycelio basali albo; velo nullo; carne alba, miti, inodora. Sporis  $6-8.5 \times 4.7-6 \mu$ , levibus, obovoideis, poro germinativo destitutis, melleis vel dilute ochraceo-brunneis, applanatione depressioneque suprahilari destitutis; basidiis  $26.5 \times 6.7 \mu$ , 2-4-sporis, plerumque tetrasporis; cystidiis nullis; cheilocystidiis  $29.5-43 \times 6.7-10.7 \mu$ , hyalinis vel pallide aureis, clavatis vel capitatis, numerosis, ex eis acie lamellarum heteromorpha; epicute ex hyphis filamentosis consistente quae elementa terminalia cystidiformia exhibunt; dermatocystidiis hyalinis vel aureis, plenumque hyalinis, ascendentibus, e. gr.  $26-36 \times 4.7-6.8 \mu$ , clavatis vel subcapitatis, rarius erectis, clavatis vel capitatis; tramate hymeniophorali ex hyphis hyalinis parallelo-subintertextis consistente; regulari; hyphis hypodermii frequenter incrustationibus aureo-ochraceis gaudentibus, sed haud abundanter incrustatis, pigmento membranali. Ad truncum arboris Dicotyledonum in silva subtropicali gregatim. Argentina, Tucumán, Sierra de San Javier, Parque Aconquija, 17-II-1955, leg. Singer no. T 2176 (LIL), typus.

**Simocybe tucumana** Sing. spec. nov.

Pileo pallide fusco vel fusco ("yellow beige", 14-I-9, "cowboy",



M & P), margine frequenter pallidiore, hygrophano, dehydratione usque ad eburneum vel brunneolum pallescente (siccio "ivory", "putty", "chipmunk"), subglabro vel subtomentoso, pallidiore etiam in siccis, aut pallide alutaceo in toto pileo siccio, circumferentia circulari vel oblonga (longitudinaliter), integra vel lobata, margine stricto in juvenilibus, convexo, papillato aut obtuso, levi nec transparentia striato, 7—29 mm. lato; lamellis pileo subconcolori dein sporis in massa concoloribus, mediocriter confertis, mediocriter latis, adnexis vel adnatis; sporis in massa brunneolis ("butterscotch" M & P); stipite pileo subconcolori sed tomento ascendente a basi albo, centrali vel sublaterali, apicem versus subattenuato vel subaequali, obliquo et stricto aut curvato, 3—10 × 1.5—3 mm.; velo nullo; carne exigua, superficiebus subconcolori, subinodora. Sporis 6—7.5 × 4.7—5.5  $\mu$ , levibus. membrana firma sed non crassa ochraceo-brunneola (ex endo- et episporio consistente) instructis, ovalibus (sed basi latioribus et lateraliter visis uno in latere minus convexis quam in altero in latere interno applanatis, rarissime subdepressis in latere interno), apice continuis haud papillatis nec porigeris, appendice hilarum exiguo munitis; basidiis 14—20 × 5.2—7.4  $\mu$ , tetrasporis; cystidiis nullis sed interdum basidiolis pigmentatis visis; cheilocystidiis 16—41 × 6.7—14  $\mu$ , versiformibus, plerumque clavatis, paucis ascendentibus, plurimis erectis, hyalinis, rarius ochraceo-brunneolis, levibus, densis (ex eis acie heteromorpha); epicute pilei hymeniformi sed hymenio pilei interrupto irregularique vel in fascicula hymeniformia disrupta, dermatocystidiis hyalinis vel brunneolis, cheilocystidiis simillimis sed interdum oblique appendiculatis, raro subdiametricis, aut applicatis, aut geniculatis, 13—40 × 6.5—12  $\mu$ ; infra dermatocystidia hyphis plus minusve intertextis subinaequalibus, interdum membrana subincrassata praeditis vel melleo-tunicatis, paucis pigmento melleo incrustatis; tramate hymenophorali regulariter dispositis, hyalinis, dein e pigmento intracellularem stramineis, haud incrustatis, crassitudine variabilibus (2.5—16.5  $\mu$  in diam.) vel tenuibus; hyphis ceteris hyalinis vel subhyalinis, fibuligeris. Ad superficiem stipitis dermatocystidiis in apice tantum praesentibus, fasciculatis et eis pilei simillimis. Ad lignum arborum Dicotyledonum in silva subtropicali gregatim, Argentina, Tucumán, Sierra de San Javier, Parque Aconquija, 17-II-1955 et 26-II-1961, leg. Singer no. T 2181 (LIL, paratypus); T 3661 (LIL, holotypus).

*Simocybe curvipes* (Sing.) Sing. comb. nov.

*Naucoria curvipes* Sing., Sydowia 8: 135. 1954.

*Simocybe ovalis* Sing. spec. nov.

Pileo pallidissime ochraceo, macroscopice subglabro, asperulato-velutino sub lente in siccis, levi, circumferentia ovali, 5—6 mm. lato; lamellis albis, dein brunneolis acie albidis, latis, subdistantibus, interdum distantibus (11 lamellis integris!), anguste adnexis vel

sublate adnatis; stipite albo vel pallide flavido, fibrilloso-vel flocculoso-scabro sub lente, brevi, prope latus latius pilei affixo, curvato, sublateriali,  $1 \times 0.6$  mm.; mycelio basali sparso vel abundante, albo, Sporis  $7-8.2 \times 5.8-6.3$   $\mu$ , levibus, ellispoideis vel breviter ellipsoideis, latus ad interius haud applanatis nec applanatione aut depressione suprahilari gaudentibus, numquam reniformibus, apice rotundatis, poro germinativo destitutis sed rarius interruptione membranae ad apicem notata (callo vel poro vis evoluta); basidiis  $22 \times 8.2$   $\mu$ , tetrasporis, pro ratione brevibus et latis, hyalinis; cystidiis nullis; cheilocystidiis  $31-40 \times 4.3-6$   $\mu$ , filamentosis vel e basi ventricosa filamentosis, apice claviculatis vel subclaviculatis et saepe sub apice constrictis, usque ad  $5.5$   $\mu$  latis, constrictione  $2.7-3.5$   $\mu$  diam. hyalinis vel pallidissimime flavidis, densis (ex eis acie heteromorpha); epicute subpalsadica, ex elementis filamentosis vel filamentoso-capitatis pallide melleis vel flavis ( $\text{NH}_4\text{OH}$  aut  $\text{KOH}$ ) subtus magis pigmentatis et ibidem frequenter pigmento epicellulari flavido adspersis, elementis eis (dermatocystidiis)  $18-41 \times 3.5-4.8$   $\mu$ ; hyphis hypodermii partim flavidis, ceterum hyphis hyalinis, omnibus fibulatis. Ad ramulos delapsos Dicotyledonum gregatim in Horto Bogotensi Venezuelae. Leg. Dennis no. 1072 (LLL, K), typus.

**Simocybe sumptuosa** (Orton) Sing. comb. nov.

*Naucoria sumptuosa* Orton, Trans. Brit. Myc. Soc. **43**: 324. 1960.

**Simocybe olivaceonana** (Sing.) Sing. comb. nov.

*Naucoria olivaceonana* Sing., Sydowia **8**: 136. 1954.

**Simocybe pantelaecodes** (Sing.) Sing. comb. nov.

*Naucoria pantelaecodes* Sing., Sydowia **8**: 138, 1954.

**Simocybe melleiceps** (Murr.) Sing. comb. nov.

*Galerula melleiceps* Murr., Proc. Fla. Acad. Sc. **7**: 119. 1944 (publ. 1945).

**Simocybe citrinipes** (Murr.) Sing. comb. nov.

*Naucoria citrinipes* Murr., l. c. p. 124.

**Simocybe olivaceiceps** (Sing.) Sing. comb. nov.

*Naucoria olivaceiceps* Sing., Sydowia **8**: 137. 1954.

**Simocybe subolivacea** (Murr.) Sing. comb. nov.

*Naucoria subolivacea* Murr., North Amer. Flora **10**: 182. 1917.

**Simocybe praeandina** (Sing.) Sing. comb. nov.

*Naucoria praeandina* Sing., Sydowia **4**: 149. 1950.

**Simocybe polytropa** (Sing.) Sing. comb. nov.

*Naucoria polytropa* Sing., Sydowia **7**: 237. 1953.

**Simocybe polytropoides** Sing. spec. nov.

Pileo dilute brunneo ("Aztec" M & P); lamellis subconfertis vel subdistantibus, latis; stipite curvato excentricoque; sporis minoribus ( $6.8-7.7 \times 4.2-4.8$   $\mu$ , haud  $6.8-8.5 \times 4.8-5.5$   $\mu$ ); ceterum *N. polytropae* Sing. holotypo, simillima species. Ad ramum delapsam. Argen-

tina, Tucumán, Playa Grande, Rio de los Sosas, 800 m. altitudinis. Leg. Singer no. T 1120 (LIL), typus.

**Simocybe atomacea** (Murr.) Sing. comb. nov.

*Atylospora atomacea* Murr., Lloydia 5: 152. 1942.

**Simocybe subvelutina** (Sing.) Sing. comb. nov.

*Naucoria subvelutina* Sing., Sydowia 4: 149. 1950.

**Simocybe serrulata** (Murr.) Sing. comb. nov.

*Naucoria serrulata* Murr., North Amer. Flora 10: 178. 1917.

**Simocybe pallidissima** (Sing.) Sing. comb. nov.

*Naucoria pallidissima* Sing., in Sing. & Digilio, Lilloa 25: 366. 1951 (publ. 1952).

**Simocybe alachuana** (Murr.) Sing. comb. nov.

*Naucoria alachuana* Murr., Lloydia 5: 149. 1942.

**Simocybe tepeitensis** (Murr.) Sing. comb. nov.

*Naucoria tepeitensis* Murr., Mycologia 4: 79. 1912.

**Simocybe unicolor** (Peck) Sing. comb. nov.

*Naucoria unicolor* Peck, Ann. Rep. N. Y. State Mus. 41: 68. 1888.

**Phaeomarasmium limulatus** (Fr. ex Weinm.) Sing. comb. nov.

*Agaricus limulatus* Fr. ex Weinm., Hymen. Ross. p. 201, 1836.

**Phaeomarasmium nebularum** Sing. spec. nov.

Pileo dilute ochraceobrunneo (12—J—8 M&P), opaco, sub lente tenuiter tomentoso, levi, convexo, papilla minutissima instructo, 7 mm. lato; lamellis pileo concoloribus, intermixtis, distantibus (13 lamellae integrae!), adnatis, latis; stipite intense saturateque castaneo, floccoso vel subglabro, subaequali, 8×0.5 mm. Sporis 9.5—13×7.2—7.7 μ, facile collabentibus, dilute ochraceo-brunneis, membrana tenui duplici instructis, late ellipsoideis vel ellipsoideis, levibus, poro germinativo destitutis, applanatione depressioneque suprahilari destitutis; basidiis 15—18×(7)—8.3—11.7 μ, tetrasporis, paucis bisporis; cystidiis nullis; cheilocystidiis (20)—55—60×8.5—11 μ, prope basin ventricosis, longe rostratis in parte apiculari (rostro 44—40 μ longo, 2—2.8 μ lato), hyalinis; epicute ex elementis elongatis (cylindraceis incrustatis, latis, e. gr. 9 μ diam.) apice late rotundatis, numquam isodiametricis (raro uno duobusve membris subsodiametricis praesentibus), catenulatis, interdum nonnullis membris catenulae ventricosis, tenuiparietalibus; hyphis fibulatis. Ad stipitem ligneum exiguum, in humo profundo putrescentem. Bolivia, Yungas, La Paz, Nor-Yungas, Cotapata, 3000 m. altitudinis, 28-I-1956. Leg. Singer no. B 637 (LIL), typus.

**Phaeomarasmium sect. Confragosi** Sing. sect. nov. Pileo epicute tecto quae ex elementis catenulatis (cellulis) elongatis efformata est. Membrana cellularum epicuticularium pigmento incrustata. Species typica sectionis: *P. confragosa* (Fr.) Sing.

**Phaeomarasmius muricatus** (Fr. ex Fr.) Sing. in Sing. & Digilio ex Sing. comb. nov.

*Agaricus muricatus* Fr. ex Fr., Syst. Mycol. 1: 244. 1821.

*Gymnopilus* sect. **Annulati** Romagnesi ex Sing. Annulo praesente. Typus sectionis: *G. spectabilis* (Fr.) Sing.

**Gymnopilus oxylepis** (Berk. & Br.) Sing. comb. nov.

*Agaricus oxylepis* Berk. & Br., J. Linn. Soc., Bot. 11: 541. 1871.

**Gymnopilus filiceus** (Cooke) Sing.

*Agaricus filiceus* Cooke, Seeman's J. Bot. 1863: 65.

**Gymnopilus traillii** (Berk. & Cooke) Sing. comb. nov.

*Agaricus traillii* Berk. & Cooke, J. Linn. Soc. 15: 368. 1876.

**Gymnopilus crocias** (Berk. & Br.) Sing. comb. nov.

*Agaricus crocias* Berk. & Br., J. Linn. Soc., Bot. 11: 542. 1871.

**Gymnopilus chrysites** (Berk.) Sing. comb. nov.

*Paxillus chrysites* Berk., Hooker's J. Bot. & Kew Gard. Misc. 3: 41. 1851.

**Gymnopilus marasmioides** (Berk.) Sing. comb. nov.

*Agaricus marasmioides* Berk. Hooker's J. Bot. & Kew Gard. Misc. 8: 133. 1856.

**Gymnopilus humicola** P. R. Harding ex Sing. spec. nov.

Pileo intense ferrugineo-brunneo vel obtuse brunneo vel obscure melleo, juventute et marginem versus interdum dilutius colorato, siccio, subumbonato vel obtuso, disco interdum olivaceo-umbrino, squamuloso, 22—34 mm. lato; — Lamellis siennicoloribus ("raw sienna" M & P), juventute flavis, marginem pilei versus etiam magis flavidis, aciem versus flavidis, angustis vel subangustis, confertis vel confertissimis, frequenter brunneo-maculatis, adnatis; stipite flavo, dein flavido basique fusca vel olivaceo-umbrino obscurato, fibrilloso, solido, dein farcto, aequali, 16—42 × 3—6 mm.; velo nullo; carne ochracea, e basi brunnescente, in apice pilei flava, in pileo dein alutacea; odore subaromatico, sed debili, Naematoloma fasciculare in mentem revocante, sapore submiti vel subamariusculo. Sporis 7.3—9 × 4.5—6  $\mu$ , verruculosi, area suprahilari levi destituti; basidiis 24—30 × 6.5—7  $\mu$ , tetrasporis; cystidiis nullis; cheilocystidiis 19.5—24 × 5.2—6.8  $\mu$ , aureo-ferrugineis vel hyalinis, ampullaceis, apice capitatis; tramate hymenophorali hyalino, regulari vel subregulari; epicute e catenulis elementorum elongatorum pigmento incrustatoeum 4—13.5  $\mu$  diam., efformata; hyphis omnibus (exceptis septis secundariis interdum numerosis) fibuligeris. Ad humum silvestrem nec nō ad terram arenosam in silva mixta. Reactiones chemicas: NH<sub>4</sub>OH ope superficiebus atroferrugineo-maculantibus et KOH ope superficiebus nigrescentibus. U.S.A., Michigan, Emmet Co., Colonial Pt., 4-VII-1953. Leg. Singer no. N 383 (F), paratypus — Cheboygan Co., east shore of Burt Lake, 3-VII-1953 (F), typus. — Praeterea collectiones in MICH conservatae a cl. Harding determinatae.



*Pyrrhoglossum holocrocinum* (Berk.) Sing. comb. nov.

*Agaricus holocrocinus* Berk., Hook. J. Bot. 5: 485. 1846.

*Pyrrhoglossum lilaceipes* Sing. spec. nov.

Pileo ferrugineo (inter "alamo" et "agate" M&P), marginem versus aureobrunneo ("Titian gold" M&P), hygrophano, in siccis vivis usque ad stramineum pallescente, glabro sed in parte posteriore tomentoso, margine lobato, convexulo sed in siccis parte posteriore umbonato, usque ad 50 mm. lato, circumferentia elliptica; lamellis ferrugineo-brunneolis ("Alamo" M&P), angustis (3 mm. latis), confertis; subdecurrentibus vel decurrentibus; stipite lilaceo ("orchis" vel "windflower" M&P), velutino-tomentoso, usque ad 11×3 mm.; carne tenuissima, inodora. Sporis 5.4—5.6×4.2—4.4 μ, intense ferrugineo-verrucosis ornamentatione exosporiali supra episporio flavido, typo ornamentationis VI, raro VI—II—IV, verrucis 0.5 μ altis; basidiis 18—18.5×6—6.2 μ, tetrasporis; cystidiis nullis; cheilocystidiis zonam citrinam formantibus, 20—27.5×5.5—7.5 μ, versiformibus, plerumque capitatis basi ventricosis, interdum ampullaceo-cylindraceis, capitulo 2.8—7 μ diam., irregulariter dispositis; tramate hymenophorali citrino, regulari, membranis hypharum usque ad 1 μ crassis; subhymenio citrino, ex elementis exiguis efformato; tramate pilei ex hyphis frequenter crassitunicatis et magnis composito; apicute ex hyphis tenuioribus quam in carne et tenuitunicatis efformata, earum membris terminalibus interdum liberis et piliformibus vel dermatocystidioideis, tenuitunicatis, citrinis, subulosis vel cylindraceis, 26—26.5×5.5 μ (apice interdum usque ad 2.7—3.5 μ attenuatis). Hyphis omnibus fibuligeris. Ad lignum putridum in silva montano-tropicali. Bolivia, Yungas, La Paz, Nor-Yungas, Carmen Pampa, 17-II-1956. Leg. Singer no. B 1225 (LIL), typus.

*Phaeocollybia amazonica* Sing. spec. nov.

Pileo brunneo ("burnt umber" M&P) et partim atrobrunneo ("leafmold" M&P) hygrophano, desiccatione multum pallidiore, per quartam partem radii pellucide striato, 40—45 mm. lato, convexo sed maturitate circum umbonem depresso; lamellis ochraceo-brunneis ("Sudan brown" M&P), demum intensius ferrugineis e massis sporarum depositis ("alamo" M&P), confertis, latis, liberis; stipite dilute alutaceo-cinnamoneo vel cinnamomeo ("dorado" vel "cinnamon" M&P), glabro, levi, cavo, ad basin incrassato sed basi pseudorhiza prolongata, parte epigea 40—45×8 mm.; ad apicem 5 mm. diam., pseudorhiza acute longeque attenuata; carne odore saporeque raphanaceo insigni. Sporis 8.2×4.8 μ; levibus sed subtiliter nebuloso-marmoratis (immersiois ope in KOH visibili ornamentatione exosporiali), fusiformibus, depressione haud levi sed paullum manifesta suprahilari praeditis, membrana manifeste duplici crassiuscula, saturate fulvo-ferruginea instructis, apice callum exhibentibus sed poro germinativo genuino destitutis; basidiis 15—20×6—8.3 μ,

tetrasporis; cystidiis nullis; cheilocystidiis  $33-47 \times 6.8-8.2 \mu$ , subcylindraceis, sed subirregularibus, subflexuosis, subcapitatis vel subampullaceis vel filamentoso-curvatis, hyalinis; ad aciem lamellarum etiam pseudoparphysibus sparsis, e. gr.  $25 \times 14 \mu$  interdum praesentibus, vesiculoso-clavatis, contenu granuloso brunneolo repletis; hyphis tramatis hymenophoralis regularis et epicutis filamentosis, subparallelis vel subintertextis, hyalinis, fibulatis; epicute cutem representante, dermatocystidiis nullis visis; hypodermio fortiter pigmentato sed haud fortiter incrustato; zonis gelatinosis haud praesentibus. Ad humum terramque (apparenter in ligno immerso) in silva hylaea pluviali solitario. Bolivia, Oriente, Beni, Vaca Diez, Riberalta, 31-III-1956. Leg. Singer no. B 2372 (LIL), typus.

**Phaeocollybia subattenuata** Sing. spec. nov.

Pileo brunneo (14 E 11 M & P) dehydratione hygrophanica usque ad ochraceum pallescente umbone brunneolo ("Ponce de Leon" M & P), leniter subviscido-subopimo, glabro, radiatim arato inter zonam marginalem, ubi striatellus est per radii partem minutissimam et umbonem qui levis est, campanulato, 40 mm. cc. lato et usque ad 20 mm. alto; lamellis subargillaceo-brunneis ("Aztec" M & P), confertis, subluberis; stipite pallido, subtus ferrugineo-maculato, glabro, levi, sicco, cavo, cepiformi ad basin, pseudorhiza subtus attenuata breviuscula ornato, apicem versus subaequali vel lenissime attenuato, parte epigaea plus minusve 100 mm. longo, ad apicem 5, ad basin 11 mm. diam.; carne odore raphani praedita. Sporis  $7.5-8.2 \times 4.5-5.5 \mu$ , late ellipsoideo-fusiformibus, circumferentia levibus sed subtiliter punctulatis (ut in *P. radicata*, sed magis visibiliter ornamentatis quam in *P. amazonica*); basidiis  $18.5-27.5 \times 7.5-8.2 \mu$ , hyalinis, clavatis, tetrasporis; cystidiis nullis; cheilocystidiis numerosis  $20-28 \times 3.5-5.5 \mu$ , versiformibus, plerumque anguste filamentoso-subclavatis et interdum omnino filamentosis vel apicem versus subattenuatis, frequenter irregulariter furcatis vel excrescentiis ornatis vel geniculatis, sed frequentius simplicibus erectisque; tramate hymenophorali regulari; subhymenio hyalino, manifesto, ex elementis exiguis efformato; epicute subindifferentiata ab hypodermio, ex hyphis pigmentatis, angustis, parallelis, haud gelatinisatis, inamyloideis formata; hyphis omnibus inamyloideis, fibulatis. Ad terram humosam in silva pluviali tropicali solitario. Bolivia, Oriente, Pando, Madre de Dios, Las Piedras, 5-IV-1957. Leg. Singer no. B 2510 (LIL), typus. — Primae species tropicales huius generis cognitae sunt *P. amazonica* et *P. subattenuata*. Cheilocystidiis versiformibus nec non praesentia fibularum et tenuissima sporarum ornamentatione insignes et probabiliter primitivae sunt.

**Crepidotus velutinoaffinis** Sing. spec. nov.

Pileo albido-tomentoso supra epicutem brunneolam, levi sed subtiliter ruguloso in siccis, 2-4 mm. lato.; lamellis in aciebus pallidis,

ceterum sporis brunnescentibus, angustis, distantibus, subobtusis; stipite brunneo, albo-pruinoso, sublaterali, curvato,  $1.8 \times 0.4$  mm. Sporis  $7.5-9.7 \times 7.5-9$   $\mu$ , subglobosis vel globosis, crude verrucosis vel breviter cristatis et ornamentatione vix immersa, asperis circumferentia, episporio bene pigmentato, ornamentatione haud obscurius colorata quam episporium; basidiis  $26-29.5 \times 8.8-9.5$   $\mu$ , clavatis, hyalinis, tetrasporis; pleurocystidiis nullis; cheilocystidiis numerosis, hyalinis, paucis ochraceis,  $40-61 \times 3.8-7$   $\mu$ , cylindraceis vel subventricosis, rarius aequalibus quam apice subcapitatis; hyphis omnibus fibulatis. In muscis humidis, Venezuela, Mérida, Sierra de Santo Domingo, Laguna Negra, 31-VII-1958. Leg. Dennis no. 1801 (K, LIL). Typus in Herbario Kewensi conservatur.

*Crepidotus serotinus* Sing. spec. nov.

Pileo brunneo, hygrophano, dehydratione in vivis usque ad flavido-album pallescente, striatulo in humidis, levi in siccis, sub lente subfibrilloso fimbriatoque, 8 mm. lato.; lamellis brunneis, latis, medioeriter confertis, concurrentibus; stipite nullo. Sporis  $9.3-12.7 \times 7-8.7$   $\mu$ , ellipsoideis, multis submucronatis in apice, ochraceo-brunneis, callo munitis sed poro germinativo destitutis, basidiis  $24-33 \times 8-13$   $\mu$ , tetrasporis; cheilocystidiis  $40-56 \times 6-8$   $\mu$ , non capitatis, sed versiformibus; ampullaceis, cylindraceis, interdum ad apicem bifurcatis, hyalinis; cystidiis nullis; tramate pilei et lamellarum haud gelatinoso et zonis gelatinosis destituto vel paulum subgelatinoso (ut in *C. antillarum*), ex hyphis tenuibus fibulatis, filamentosis efformato. Ad ramum delapsam Dicotyledonum in silva semisicca serotinus (tarde autumnus!). Argentina, Tucumán, Arroyo Mista, 15-VI-1955. Leg. Singer no. T 2782 (LIL), typus. — Mycelium basale abundans, subpelliculosus, albidus vel albus. *C. parlatorei* Sing. affinis.

*Crepidotus polylepidae* Sing. spec. nov.

Pileo intense aureo-flavo, omnino pubescente, levi, convexo conchato, 5 mm. cc. lato; lamellis dilute flavis, dein brunneolis acie flavis latissimis, medioeriter latis, concurrentibus; stipite rudimentario, haud manifesto, velutino. Sporis  $6.2-7.5$   $\mu$  diam. globosis, spinulis episporio concoloribus cylindricis in episporio immersis ornatis, paullum punctatis (ornamentatione typi *C. applanati*); basidiis  $25 \times 7$   $\mu$ , tetrasporis; cystidiis nullis; cheilocystidiis  $33-81 \times 5.5-8.2$   $\mu$ , hyalinis, subulatis apice subacuto angustato praeditis vel filamentosis basi dilatata; epicute pilei palisata trichodermialiter formata elementorum dermatocystidioeorum et eorum continuationum hyphalium, intus succo auro-flavido impletorum, elementis vesiculososis inter ea admodum sparsis et  $16-21 \times 11$   $\mu$  cc. metientibus; hyphis fibulatis. Venezuela, Mérida, Sierra de Santo Domingo, 3600 m. altitudinis, 24-VII-1958. Leg. Dennis no. 1726 (K), typus. — Ab

affini *C. crocophyllo* *carpophoris* minoribus et forma *cheilocystidi-  
orum* differt.

***Clitopilus rhodophyllus*** (bres.) Sing. comb. nov.

*Pleurotus rhodophyllus* Bres., Ann. Mycol. 3: 159. 1905.

***Rhodocybe lignicola*** Sing. spec. nov.

Pileo rubescenti-brunneo ("Arab" M & P), hygrophano, longe pellucide striato, 18 mm. lato, convexo, umbilicato; lamellis carnis, confertis at haud confertissimis, adnatis; stipite pileo concolori, apice intensius rubrobrunneo („Mohawk" M & P), subaequali vel apicem versus subattenuato, 28×2 mm. (ad basin usque ad 3 mm. lato). Sporis a polo visis angularibus, asperulatis, 6.2—7×4.2 μ; basidiis tetrasporis, 24.5—25×8—8.5 μ cystidiis nullis; tramate hymenophorali regulari, hyalino, fibulato. Ad ligna emortua in zona superiore silvarum tropicali-montanarum. Bolivia, Yungas, La Paz, Nor-Yungas, Carmen Pampa, 26-II-1956. Leg. Singer no. B 1495 (LIL), typus.

***Rhodocybe laeta*** Sing. spec. nov.

Pileo ferrugineo-aurantiaco ("Mosque" M & P) vel intense ferrugineo ("Kobe" M & P), margine aurantiaco-salmoneo ("honeydew" M & P), zonis concentricis sordidioribus vel aquosis vel obscurioribus adornato, umbilicato, irregulariter convexo vel centro applanato vel subdepresso, umbilicato, 52 mm. lato; lamellis obtuse ochraceis (11 D 6 M & P), confertis, in siccis haud distantioribus quam eae *R. russulariae* aut *R. collybioideae*, arcuato-decurrentibus; stipite pallidiore sordidioreque pileo, ad basin incrassato, 40 mm. cc. longo, 5 mm. lato ad apicem, basi 10 mm. diam.; carne miti. Sporis a polo visis angularibus, asperulatis, 4.8—6.2(7)×4—4.8(5.5) μ; basidiis (2)—4-sporis, 18—25×4.8—5.5 μ, clavatis, hyalinis; cystidiis paucis et sparsis, cylindraceo-subulatis vel subulatis, acuminatis vel subacutis vel acutis, supra basidia non vel vix projicientibus, inconspicuis, 3.5—4.5 μ diam., hyalinis. Ad humum silvestrem lignosum in hylaea. Bolivia, Oriente, Beni, Vaca Diez, Guayaramerin, 14-III-1956. Leg. Singer no. B 1951 (LIL), typus. — *A. R. collybioidea* coloribus nec non consistentia rigidiore fragili differt.

***Rhodocybe rickii*** Sing. nom. nov.

*Clitocybe gilvoidea* Rick, Broteria 24: 101. 1930.

***Rhodocybe rickii*** var. *convexa* (Sing.) Sing.

*Rhodocybe gilvoidea* var. *convexa* Sing., Sydowia 12: 242. 1958.

***Rhodocybe russularia*** Sing. spec. nov.

Pileo dilute cinnamomeo, subaurantio-cinnamomeo in statu sicco, (13-C-9/11, 11-D-8; sicco 13-B/C-9/10, 11-R-7, 13-A-9 "creole", M & P), hygrophano. subopimo sed haud viscido, glabro et levi, sed dense pellucide striato in zona marginali vel estriato, convexo, dein applanato, frequenter umbilicato, 38—51 mm. lato; lamellis ochraceo-gilvis, vel sordide gilvis ("lariat" M & P), confertissimis, angustissimis



vel angustis (usque ad 4 mm. latis), adnatis vel adnato-decurrentibus; sporis in massa pallide carneolis (2 A 8 M & P); stipite aut pileo aut lamellis concolori, glabro vel initio plus minusve innato-sericeo-striatulo apice pruinoso vel, subpruinoso, cavo, apicem vel basin versus attenuato, 46—68×4.5—7 mm., plus minusve subradicante; carne in humidis alba vel superficiebus plus minusve subconcolori sed tunc pallidiore in siccis, odore Collybiae dryophilae vel subfarinaceo; sapore miti vel farinaceo. Sporis polo visis angularibus, asperulatis, stramineo-carneolis, sed admodum pallidis, 4.8—6.5×3—4.3  $\mu$ ; basidiis 22×6.5  $\mu$  tetrasporis; cystidiis contentu heterogeno visibili destitutis, 23—49×5.5—6.3  $\mu$ , tenuitunicatis, hyalinis, levibus, ventricosis, apice ampullaceo-extensis, filamentosis, pedicello longo instructis; apicute ex hyphis repentibus parallelis filamentosis efformata; tramate hymenophorali regulari, hyalino. Hyphis omnibus defibulatis. Ad terram arenosam vel ad detritum hylaeae. Bolivia, Oriente. Beni, Vaca Diez, Guayaramerin. Leg. Singer no. B 1709 (typus, LIL), B 2015 (LIL, paratypus). Color habitusque Lactarii cuiusdam e sectione Russulariarum.

**Rhodocybe galerinoides** Sing. spec. nov.

Pileo obtuse brunneo ("bure", "butterscotch" M & P), hygrophano, dehydratione flavo (10-E-5 M & P), haud profunde subsulcato, (glabro, conico, papillato, 16 mm. lato, 7 mm. alto; lamellis ochraceis 10/11-F-6 M & P), subdistantibus vel distantibus, profunde sinuatis, latuscusculis, haud intervenosis, ventricosis; stipite aurantiaco vel aurantiaco-cinnamomeo, (12-G-11, 13-B-11 M & P), cylindraceo, siccio, glabro, 25×1.5 mm.; carne inodora. Sporis polo visis subangularibus, sublevibus vel crude asperulatis, 4.8—5.5×3.8—4.2  $\mu$ ; basidiis 22×4.5—7  $\mu$ ; cystidiis nullis, sed pseudocystidiis conspicuis, prope aciem mediocriter numerosis, cum hyphis oleiferis connexis, haud incrustatis, longis, ventricosis vel subcylindraceis, 6.8—9  $\mu$  latis; tramate lamellarum regulari, hyalino; strato cuticulari pilei ex hyphis melleis, pigmento membranali destitutis filamentosis repentibus efformata; hyphis omnibus defibulatis. Ad lignum putridum in silva pluviali tropicali. Bolivia, Oriente, Beni, Vaca Diez, Guayaramerin, 14-III-1956. Leg. Singer no. B 1973 (LIL), typus.

**Rhodocybe lactariiformis** Sing. spec. nov.

Pileo brunneo vel badio ("kiskilim" vel "montella" M & P), leniter pellucide striatulo, subapplanato, acute papillato, 29 mm. lato; lamellis cinnamomeis, confertis, adnatis; stipite cinnamomeo, glabro, apicem versus attenuato, 50×3.5 mm., ad basin 6 mm. diam. Sporis polo visis subangulatis vel angulatis, asperulatis, 4.8—6.2 (7.5×4—4.8(5.5)  $\mu$ ; basidiis 25×7  $\mu$ , hyalinis, clavatis, tetrasporis; pseudocystidiis contentu melleo guttiformi impleto, cum hyphis oleiferis concurrentibus, quae item contentu melleo guttato gaudent, haud incrustatis, 62 cc.  $\mu$  longis et 4.8—8.2  $\mu$  latis, cylindraceo-vermiformi-

bus, frequenter particus ventricosus instructis vel apice acuminato gaudentibus, sed constanter ad ipsum apicem rotundatis, longitudine variabilibus, contentu iodi ope castanescente (ut in *R. caelata*); tramate hymenophorali regulari, ex hyphis parallelis filiformibus hyalinis efformato; epicute ex hyphis repentibus, cutem formantibus efformata; hyphis omnibus defibulatis. Ad lignum Dicotyledonum in hylaea. Bolivia, Oriente, Beni, Vaca Diez, Guayaramerin, 17-III-1956. Leg. Singer no. B 2070 (LIL), typus. — Species *R. naucorioide* maior; lamellis confertis, cystidiis haud incrustatis recendes. A *R. galerinoide* forma et magnitudine pilei, coloribus obscurioribus et superficie haud hygrophana (vel vix hygrophana), contentu magis denso pseudocystidiorum differt. Habitus Lactarii coloresque eiusdem generis exhibit.

*Paxillus* sect. **Parapaxillus** (Sing. ex) Sing. sect. nov. Fibulis praesentibus; velo praesente. Typus sectionis: *Paxillus argentinus* Speg.

*Paxillus* sect. **Defibulati** Sing. sect. nov. Fibulis nullis; velo nullo. Typus sectionis: *Paxillus defibulatus* Sing.

*Paxillus* sect. **Veluticipites** Sing. sect. nov. Carpophoris evelatis; sporis elongatis (13—16  $\mu$ ); cystidiis nullis; fibulis praesentibus. Typus sectionis: *Paxillus veluticeps* (Cooke & Mass.) Sing.

**Paxillus veluticeps** (Cooke & Mass.) Sing. comb. nov.

*Agaricus veluticeps* Cooke & Mass., *Grevillea* 18: 4. 1889.

**Gomphidiaceae** Maire ex Sing. fam. nov. Habitus clitocyboideus; lamellae ceraceae; leptocystidia praesentia; hyphae defibulatae; trama hymenophorale bilaterale, sporae in massa subatrae, s. l. griseae, leves; fungi mycorrhizici, ectotrophici. Genus typicum: *Gomphidius* Fr.

**Suillus nueschii** Sing. nom. nov.

*Boletus flavus* With. ex Fr. sensu Bres., *Icon. Mycol.* 19: 904 pl. 904. 1931.

**Suillus nueschii** var. **caerulescens** Sing. var. nov.

A *Suillo nueschii* var. *nueschii* carne autoxydatione caerulescente et KOH ope viridescente differt. Sub Laricibus sibiricis in zona montana et subalpina Montium Altaicorum, U.S.S.R., prope Kurai frequens. Leg. Singer & Vasilieva aestate 1937, typus in Herb. LE conservatus est.

(*Ixocomus flavus* f. *caerulescens* Sing. *Rev. Myc.* 3: 42. 1938).

**Suillus bovinus** (L. ex Fr.) Kuntze var. **viridicaerulescens** (Pearson) Sing. comb. nov.

**Phylloporus hyperion** (Cooke & Mass.) Sing. comb. nov.

*Agaricus hyperion* Cooke & Mass., *Grevillea* 16: 72. 1888.

*Pulveroboletus* sect. **Sulphurei** (Sing.) Sing. comb. nov.

*Phlebopus* sect. *Sulphurei* Sing., *Am. Midl. Natur.* 37: 2. 1947.

**Pulveroboletus hemichrysus** (Berk. & Curt.) Sing. comb. nov.

*Boletus hemichrysus* Berk. & Curt., *Ann. Mag. Nat. Hist.* II. 12: 429. 1853.

*Boletus sulphureus* Fr., *Epicrisis*, p.413. 1838, non Bull. ex Mérat 1821.

**Boletellus verrucarius** (Berk.) Sing. comb. nov.

*Boletus verrucarius* Berk., Hooker's J. Bot. & Kew Gard. Misc. 6: 135. 1854.

**Russula picearum** Sing. nom. nov.

*Russula rubra* Fr. sensu Bresadola, Fungi Trid. 2: 91, pl. 203. 1887.

*Lactarius* sect. **Lactariopsidei** Sing. ex Sing. Velo praesente aut tramite hymenophorali heteromero insignes. Typus sectionis: *Lactarius zenkeri* (Henn.) Sing.

**Lactarius reticulatus** (Berk.) Sing. comb. nov.

*Panus reticulatus* Berk., Hooker's J. Bot. & Kew Gard. Misc. 8: 142. 1856.

**Lactarius alni** Sing. spec. nov.

A *L. obscurato* et *L. tabido* sporarum massae colore candido et associatione mycorrhizica cum *Alno jorullensi* differt. Argentina, Tucumán, Sierra de Sans Javier, San Agustin, 7-I-1959, 1500 m. altitudinis. Leg. Singer no. T 3181 (LIL), typus. In Alnetis austro-americanis frequens est (cf. Singer & Digilio in Lilloa 25: 452. 1951, publ. 1952 sub nom. *L. cupularis* (Bull. ex Fr.) Qué. sensu Bres.)

**Galeropsis madagascariensis** (Pat. ex) Sing. spec. nov.

*Galera besseyi* var. *madagascariensis* Pat. ined. ex Heim, Rev. Mycol. 15: 11, 13. 1950.

Sporis minoribus, pileo saepe obtuso a *G. besseyi* differt. Madagascar (P).

**Galeropsis angusticeps** (Peck) Sing. comb. nov.

*Galera angusticeps* Peck, Bull. Torr. Bot. Cl. 24: 143. 1897.

**Xerotinus erubescens** (Berk.) Sing. comb. nov.

*Daedalis erubescens* Berk., Notice on some fungi collected by C. Darwin Esqu., during the Expedition of H. M. Ship Beagle, Ann. Mag. Nat. Hist. 4: 292. 1840.

*Lenzites erubescens* (Berk.) Sacc., Syll. 5: 645. 1887.

*Xerotus erubescens* (Berk.) Sing. apud Martin, Contr. Science, Los Angeles Co. Mus. 24: 6. 1958.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1961/1962

Band/Volume: [15](#)

Autor(en)/Author(s): Singer Rolf

Artikel/Article: [Diagnoses Fungorum novorum Agaricalium II. 45-83](#)