

## Some interesting Micro-Fungi from flood affected Substrata in Poona, India.

By S. Anantha Narayanan.

(M.A.C.S. Laboratory, Law College Buildings, Poona 4, India).

With 6 Textfig.

The unprecedented floods that visited Poona on the 12th of July, 1961 left behind a large number and varieties of substrata as well as consumers goods completely soaked in the waters and mud brought by them. These substrata soon started developing many types of moulds and microfungi. A survey and detailed study were undertaken of these micro-fungi as, such a study was always helpful in determining the occurrence, distribution and the sources of pathogenic fungi. The total number of fungal genera collected by the writer is over 25 comprising mostly of *Hyphomycetes*, and a few of *Zygomycetes*. It was interesting to note that different substrata showed




Fig. 1. *Sporodesmium backeri* Syd. — A. Conidiophores with spores  $\times 440$ . — B. Developmental stages of Conidia  $\times 440$ . — C. Conidia  $\times 950$ .




Fig. 2. *Stachybotrys lobulata* Berk. — A. Branched conidiophores with conidia  $\times 264$ . — B. Conidiophore  $\times 440$ . — C. Conidia  $\times 950$ .

growth of certain specific types of micro-fungi only and no other, probably indicating biochemical relationship.

The substrata like different types of paper, leather, wooden articles, grains, fabrics of cotton, silk and wool, plastics, soil, gunny bags etc. showing mould infection were collected and kept under moist chambers for varying lengths of time to stimulate proper sporulation and formation of developmental stages. Semi-permanent slides were then immediately prepared from the colonies with a view to prevent contamination from the rapidly growing moulds like *Rhizopus* and *Aspergillus*; the fungi were then mounted in lacto-phenol and stained in Cotton Blue for final observation.

This paper gives an account of six micro-fungi, all belonging to the *Hyphomycetes*, two of which are considered new to science, three new records for India and one, *Microsporium gypseum* a dermatophyte and was obtained from animal hair.

1. *Sporodesmium bakeri* Syd. (Fig. 1). — Sydow, P. A. Ann. Mycol. Berl. XII, pp. 204, 1914. — Hughes, S. J., C. M. I. paper 50: pp. 66, 1953.

Colony black, irregular in outline, scattered, raised 0.5–1.5 mm. in diam. Conidiophores crowded, erect, continuous, short, cylindrical, persistent, hyaline to pale yellow,  $6.5\text{--}13.00\ \mu \times 2.15\text{--}5.25\ \mu$ . Conidia fusoid to cylindric oval and muriform, broad at apex, tapering at base, thick walled, 3–4 septate, generally 3-septate, constricted at septa, septa transverse, longitudinal as well as oblique, formed

singly, dark, brown, finely echinulate,  $19.5\text{--}28.0\ \mu \times 9.5\text{--}10.75\ \mu$ , detached from the conidiophore invariably bear a basal hyaline frill.

On plywood, collected by S. Anantha Narayanan, 22, VII, 1961, Poona, India.

Remarks: Hughes (1953) has collected this fungus from dead leaves of *Zea mays* and *Bridelia ferruginea*.

This is a new genus record and an addition to the fungi of India.

2. *Stachybotrys lobulata* Berk. (Fig. 2). — Saccardo, P. A. Sylloge Fungorum 4: pp. 269–270, 1886. 10: pp. 577 (1892); 11: pp. 612 (1895); 14: pp. 1071 (1899) and 22: pp. 1352 (1913). — Bisby, G. R. Trans. Brit. Mycol. Soc. 26: pp. 133–143 (1943); 28: pp. 11–12 (1945). — Subramanian, C. V. Proc. Ind. Aca. Sci. 36: pp. 43–53 (1953).

Colony, black, raised, in irregular patches, 1–1.5 mm. diam. Mycelium pale brown, branched, septate,  $3\text{--}4.5\ \mu$  thick. simple to branched, sparingly septate (1–2), dark brown at apex and paler below, broader at base, thin-walled,  $30\text{--}68.8 \times 3.2\text{--}4.2\ \mu$ , bearing




Fig. 3. *Microsporium gypseum* (Bodin) Gniart & Grigork. — A. Macroconidiophores with macro conidia attached  $\times 440$ . — B. Macroconidiophore enlarged  $\times 950$ . — C. Macroconidia  $\times 1710$ . — D. Mycelia with Microconidia  $\times 950$ . — E. Microconidia  $\times 950$ .




Fig. 4. *Septonema punctiforme* B et C. — A. Conidiophore with attached conidia  $\times 264$ . — B. Part of the Conidiophore enlarged  $\times 792$ . — C. Conidia  $\times 792$ .

a whorl of phialides. Phialides short simple, 1-celled, pale brown, thin-walled, oblong,  $8.5-13.0 \times 4.5-6.5 \mu$ . Conidia 1-celled, ovoid to ellipsoid, dark brown, distinctly verrucose,  $6.5-10.75 \times 4.3-6.5 \mu$ .

Habit: On plywood, cotton fabrics and paper, collected by S. Anantha Narayanan, 24, VII, 1961, Poona, India.

Remarks: The species collected on cotton fabric and paper showed much branched and longer conidiophores ( $150.5$  to  $155 \mu$ ), than the species collected on plywood.

The fungus genus is a new record and addition to the fungi of India.

3. *Microsporium gypseum* (Bodin) Griart & Grigork. (Fig. 3). — Ajello, *Mycologia*, 51: pp. 69-76 (1959). — Cooke, W. B. *Mycologia* 44: pp. 247-248 (1952). — Dey, N. C. & L. M. Ghosh, *Ind. Med. Gaz.* 79: pp. 51 (1944). — Fuentes, F. A., *Mycologia* 48: pp. 614 (1956). — Padhye, A. A., Prof. Agharkar Commemoration Volume pp. 117-119 (1961). — Randhwa, A. N. et. al, *Sci. Cult. (India)* 25: pp. 326-327 (1959).

Colony thick, raised, shining, cream white, irregularly spreading on the substrata. — Mycelium hyaline, thin walled, much branched,

septate 2.15—3.25  $\mu$  in breadth. — Conidiophores erect, rather short, slender, simple, 1—2 septate, hyaline, 10.75—47  $\mu \times$  2.15—3.25  $\mu$ , rarely longer (82.0  $\times$  2.5  $\mu$ ). — Micro-conidia on lateral sides of mycelium, minutely stalked, 1-celled, hyaline, globular, thin-walled, 2.3—2.75  $\mu \times$  2.2  $\mu$ . — Macroconidia fusoid, 4—7 celled, double-walled, distinctly warty, hyaline to tan coloured formed singly, 21.5—43  $\mu \times$  8.5—13.0  $\mu$ .

Habit: On animal hair, collected by S. Anantha Narayanan, 20. VII, 1961, Poona, India.

Remarks: The fungus is a soil-dwelling dermatophyte and is known to incite scalp infections. This is the 3rd record from India, the two previous ones being by Randhwa et al. (1959) from Delhi, India and Padhye (1961) from Poona Soils.

4. *Septonema punctiformae* B. et. c (Fig. 4). — Saccardo, P. A. Sylloge Fungorum 4: 397—401 (1886). — Vol. 10: pp. 609 (1892); Vol. 11: pp. 622 (1895); Vol. 14: pp. 1084 (1899); Vol. 16: pp. 1062 (1902); Vol. 18: pp. 583 (1906); Vol. 22: pp. 1381 (1913). — Cash & Watson, Mycologia 47: pp. 745 (1955).

Colony deep brown, effuse, scattered, 1.5—2 mm. Mycelium pale yellow, septate, much branched, thin-walled, 4—6  $\mu$  in width. Conidiophores branched, erect, thin-walled, subhyaline to pale yellow, septate, slightly constricted at septa, individual branches short 1—2 celled, slightly bulged at apex, 13.0—30.0  $\mu \times$  4.3—5.3  $\mu$ . Conidia in acropetal chains, profusely branched, deep yellow to brown, 2—4 celled, rarely 5-celled, deeply constricted at septa, thick-walled (elliptical to obovate), cylindrical to fusoid, smooth-walled, 13.0—21.5  $\times$  4.3—8.5  $\mu$ .

Habit: on old jute bag, plywood & bark, Poona, India, 24. VII. 1961, collected by S. Anantha Narayanan.

Remarks: This is a new record and addition to the fungi of India. Subramanian (1956) has recorded *S. bombyence* Subr. from Bombay, which however, is distinct from the present species in having 1-septate, non-constricted and verrucose nature of conidia.

#### 5. *Cephaliphora uniformis* S. A. Narayanan sp. nov. (Fig. 5).

Coloniae albae vel alutaceae, patentees 2—2.5 mm. diam. Mycelium alutaceum, tenue, ramosissimum, septatum, 6.4—8.5  $\mu$  latum. Conidiophora typice curvata gibbo ad latus posito, qui evolvitur in capitulum apicale et fertile, septata, angusta ad basin (6.4—8.6  $\mu$ ), latiora ad apicem (10.75—19.4  $\mu$ ), tenuibus parietibus praedita, pallide brunnea, 64.5—99  $\mu$  longa. Capitula fertilia pallide brunnea ex ovoideo globosa, 17.2—23.5  $\times$  21.5—30  $\mu$ , cellulis sporogenis radialiter dispositis hyalinis cuneiformibus ornata. Conidia cellulis sporogenis, singulariter insidentia, semper unicellularia et obovata, levia, parietibus crassis




Fig. 5. *Cephalophora uniformis* S. A. Narayanan sp. nov. — A. Conidiophore with conidia.  $\times 440$ . — A<sub>1</sub> Developing conidiophore  $\times 440$ . — C. Conidia  $\times 792$ . — F. Chlamydospore  $\times 440$ .

praedita, pallide brunnea.  $15-30 \mu \times 10.75-17.2 \mu$ . Chlamydosporae sphaericae, parietibus crassis praeditae, plerumque intercalares, obscure brunneae,  $21.5-23.65 \mu$  diam.

In saccis e fibris *Corchori* factis, ad Poona, India 18. VII. 1961, leg. S. Anantha Narayanan, M. AC. S. Herb. No. 107.

Remarks: The genus was established by Thaxter (1903) for two coprophilous fungi with *C. tropica* Thaxt. as type and the second one as *C. irregularis* Thaxt. The Poona fungus appears to be quite distinct in the morphological characters of the conidiophore and conidia, their shape as well as septation from *C. irregularis* Thaxt. to which it bears resemblance. The conidia are characteristically uniform both in respect of shape and septation. The Poona fungus is therefore, described as a new species on the basis of these distinct characters with Latin diagnosis.

5. *Oedocephalum indicum* S. A. Narayanan sp. nov. (Fig. 6).

Coloniae albae, dispersae vel gregariae, pulverulentae ad maturitatem,  $0.5-0.7$  mm. diam. Mycelium hyalinum, ex hyphis breviter articularis, tenuiter tunicatis ad septa plus minusve dilatatis,  $8.6-10.6 \mu$  latis compositum. Conidiophora erecta, et angulum acutum




Fig. 6. *Oedocephalum indicum* S. A. Narayanan sp. nov. — A<sub>1</sub>—A<sub>7</sub>, Developing stages of conidiophore × 264. — B, Head or vesicle × 792. — C, Conidia × 792.

formantia, tenuiter tunicata 4—6-septata, ad septa tumescencia geniculata, ad basin usque ad 13  $\mu$  lata, apicem versus angustata, in vesiculam fertile formantia 130—163.5  $\mu$  longa. Capitula fertilia globosa, hyalina, per totam superficiem sterigmatibus arte acutatis dense dispositis ornatis, 19.5—28.0  $\mu$ . Conidia singulatim sterigmatibus insidentia, hyalina, unicellularia, globoso-ovoidea, tenuiter tunicata, levia, 6.45—8.6  $\times$  4.3  $\mu$ .

In saccis e fibris *Corchori* factis, leg. S. Anantha Narayanan, ad Poona, India 18. VII. 1961, M. A. C. S. Herb. No. 108.

Remarks: Saccardo (1886—1906) lists 25 species of this genus of which three species occur in India. *O. liniatum* Bakshi was recorded by Bakhsi (1950) from Great Britain associated with Ambrosia beetles and possessing non-septate conidiophores and thick-walled conidia. The Poona species has some resemblance to *O. fimitarium* (Riess.) Sacc. (Syll. Fung. 4: 47—50) but is distinct from it in possessing conidiophores characteristically bulged at septa, densely arranged sterigmata on the fertile vesicle and ellipsoid to oblong shape of conidia. The Poona fungus is therefore, considered new to science and presented as new species with Latin diagnosis. The fungus genus is an addition to fungus of Bombay, Maharashtra.

### Summary.

This paper gives an account of six micro-fungi, all belonging to the *Hyphomycetes*, two of which are considered new to Science, three new records for India and one *Microsporium gypseum* is a dermatophyte and was obtained from animal hair.

### Acknowledgement.

The writer is grateful to Prof. M. N. Kamat for his keen interest, encouragement and guidance both during investigations and in preparation of the manuscript, to the Director, M. A. C. S., Poona-4 for the laboratory facilities, to Prof. H. Santapau for Latin rendering of the two new species and to the Secretary, University Grants Commission, New Delhi for the award of the Scholarship.

### References.

- Agnihorthrudu, V. et. al. 1955. Fungi Associated with Rhizosphere. I., Proc. Ind. Aca. Sci., 42-B, pp. 101-102.
- Bakshi, B. K. 1950. Fungi associated with *Ambrosia beetles* in Great Britain: Trans. Brit. Mycol. Soc. 33 pp. 113-116.
- Butler, E. J. & Bisby, G. R. 1931. The Fungi of India. Sci. Monogr. Coun. Agri. Res. India, 1 pp. 148.
- Mundkur, B. B. 1938. Fungi of India, Supplement I. Sci. Monogr. Coun. Agri. Res. India, 12, pp. 35.
- Srinivasan et al. 1961. *Cephalophora tropica* Thaxt. & *C. irregularis* from Poona in Prof. Agharkar Commemoration Vol. pp. 79-88.
- Subramanian, C. V. 1953. Fungi Imp. from Madras IV, Proc. Ind. Aca. Sci. B-37: pp. 96-105.
- 1952. Fungi Imp. from Madras I, Proc. Ind. Aca. Sci. 36, pp. 43-53.
  - 1955. Fungi Imp. from Madras, VII, Proc. Ind. Aca. Sci. 42: pp. 283-292.
  - 1956. Hyphomycetes II. J. I. Bot. Soc. 35, pp. 446-494.
  - 1957. Hyphomycetes IV. Proc. Ind. Acad. Sci. B-46 pp. 331.
  - 1958. Hyphomycetes V. J. I. Bot. Soc. 37, pp. 47-64.
- Uppal, B. N., Patel, M. K. and Kamat, M. N. 1935. The fungi of Bombay. VIII.



# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1962/1963

Band/Volume: [16](#)

Autor(en)/Author(s): Narayanan S. Anantha

Artikel/Article: [Some interesting Micro-Fungi from flood affected Substrata in Poona, Indien. 77-84](#)