

The Genus *Valsaria* in India

By S. B. Kale

(Botany Department Marathwada University, Aurangabad, India)

With 2 figures

The genus *Valsaria* was originally erected by Cesati & de Notaris (1863) with *V. insitiva* as the type species. However, it remained unrepresented in the Indian flora until Mundkur and S. Ahmad (1946) reported two species i. e. *V. salvadora* Mundkur and S. Ahmad on *Salvadora oleoides* Dcne. and *V. tamaricis* Mundkur and S. Ahmad on *Tamarix articulata* Vahl. Recently Kapoor and Gill (1961) reported *Valsaria bambusae* Kapoor and Gill on *Bambusa* sp.

The genus is characterised by the black valsoid or pulvinate stroma in which perithecia are clustered, ascii cylindrical, 8-spored, ascospores brown to almost black, 1-septate, uniseriate to biseriate.

During the investigation of different genera of ascomycetes from India the authors collected dried stems of *Acacia melanoxylon* Br. and *Bambusa* sp., heavily infected with some black fungus. A careful study revealed that these belong to the genus *Valsaria* Ces. & De Not. The fungus on *Bambusa* sp. was found to be *Valsaria bambusae* Kapoor & Gill. and is also compared with the type material. The fungus on *Acacia melanoxylon* Br., however, has been described as a new species on the basis of comparative morphology, besides being reported on a hitherto unreported host.

1. *Valsaria indica* sp. nov.

Stroma nigrum, valsoideum, disco obscure griseo vel nigrescente, ostioli minute papillato erumpens, $675-1200 \times 1350 \mu$; perithecia densiuscula stipata, in ostiola cylindracea, intus dense periphysata paucithecata, attenuata, $180-375 \times 180-225 \mu$; ascii cylindracei, breviter stipitati, $88-96 \times 8-8.8 \mu$, paraphysisibus filiformibus superati; sporae monostichiae, ellipsoideae, medio septatae et valde constrictae, brunneae vel obscure brunneae, $13.6-15.2 \times 5.6-7.2 \mu$.

Stroma black, valsoid, erumpent by a dark grey or black disc on which the clusters of ostioles appear as minute papillae, measuring from $675-1200 \times 1350 \mu$, perithecia black-shaped with long necks, black, measuring from $180-375 \times 180-225 \mu$, in small groups clustered within a valsoid black stroma, with a thin limiting layer; ascii cylindrical, short-stalked, measuring from $88-96 \times 8-8.8 \mu$, 8-spored, paraphysate,




Fig. 1: *Valsaria indica* sp. nov. A. Section through the ascocarp. — B. Ascus and ascospores

Comparative table of the Indian species of *Valsaria*

Species	Perithecia	Asci	Ascospores	Remarks
1. <i>V. indica</i> sp. nov. on <i>Acacia melanoxylon</i> Br.	180—375 = 88—96 ×	8—8.8 μ	13.6—15.2 × 5.6—7.6 μ	No longitudinal striations, broadly elliptical
2. <i>V. bambusae</i> K. & G. on <i>Bambusa</i> sp.	—	119—144 × 14 μ	22—22 × 10—11 μ	Longitudinal striations present, smoky brown
3. <i>V. salvadora</i> Mundkur & S. Ahmad on <i>Salvadore oleoides</i> Dcne.	350—450 μ	140 μ	16.5—24.5 × 9—11 μ	Biseriate to irregularly biseriate
4. <i>V. tamaricis</i> Mundkur & S. Ahmad on <i>Tamarix articulata</i> Vahl.	—	87.5 × 9.5 μ	12.5—14.5 × 6—7 μ	Not deeply constricted, uniseriate

paraphysis filiform, longer than the ascii; ascospores uniseriate, broadly elliptical, 2-celled, strongly constricted at the septum, brown to dark brown, measuring from $13.6-15.2 \times 5.6-7.2 \mu$; periphyses present in the neck.

Collected on the dead stem of *Acacia melanoxylon* Br. at Kadai Kanal in the month of October 1964. Leg. Tilak S. T. & S. B. Kale and deposited in the herbarium of Marathwada University, Aurangabad.

2. *Valsaria bambusae* Kapoor & Gill. (1961). Ind. Phytopath. 14: 149-153.

On the stem of *Bambusa* sp. from Dharwar. The type material from I. A. R. I. New Delhi as well as our own collections on the same host collected at Aurangabad in the month of November 1964 and January 1968, have been critically studied. Both the materials showed the typical




Fig. 2: *Valsaria bambusae* Kapoor & Gill. — Ascospore, showing the longitudinal striations

ascospores which are brown to dark brown but with longitudinal striations on the wall and this character has not been mentioned earlier.

3. *Valsaria salvadora* Mundkur & S. Ahmad (1946). Mycol.

Pap. Imp. Mycol. Inst. 18: 1-11.

on *Salvadora oleoides* DCNE.

4. *Valsaria tamaricis* Mundkur & S. Ahmad (1946). Mycol.

Pap. Imp. Mycol. Inst. 18: 1-11.

On *Tamarix articulata* Vahl.

Key to the Indian species of *Valsaria*.

- A. Stroma pulvinate wide spread initially innate, but later on superficial, black to grey.

1. Ascospores with longitudinal striations

V. bambusae.

2. Ascospores without longitudinal striations.

 - i) Ascospores uniseriate *V. tamaricis.*
 - ii) Ascospores biseriate to irregularly biseriate *V. salvadora.*

B. Stroma innate-erumpent by a black to grey disc by rupturing the bark valsoid. Ascospores broadly elliptical, strongly constricted in the middle *V. indica.*

The type materials have been deposited in the herbarium of Marathwada University, Aurangabad and Cryptogamie Indiae Orientalis, New Delhi (India).

Acknowledgements

Grateful thanks are due to Dr. S. T. Tilak, Botany Dept. Marathwada University, for his valuable guidance, to Marathwada University, Aurangabad, for the laboratory facilities, to University Grants Commission, New Delhi, for the award of Research Fellowship, to Dr. S. P. Raychaudhuri, Head Division of Mycology and Plant Pathology, I. A. R. I. New Delhi for the supply of some of the materials of *Valsaria* Ces. & de Not., and to Dr. F. Petrák for the Latin diagnosis of a new species.

Literature cited

- Butler, E. J. & G. R. Bisby revised by Vasudeva, R. S., 1960: The fungi of India. I. C. A. R. New Delhi. p. 67.

Clements, F. E. & C. L. Shear, 1931: Genera of fungi. Hafner Publ. co. New York. p. 70.

Dennis, R. W. G., 1960: British cup-fungi and their allies, Ray Society London. p. 235.

Kapoor, J. N. & H. S. Gill, 1961: Notes on Indian ascomycetes I. Ind. Phytopath. 14: 149—153.

Mundkur, B. B. & S. Ahmad, 1946: Revision of additions to Indian fungi II. Mycol. Pap. Imp. Mycol. Inst. 18: 1—11.

Saccardo, P. A., 1882: Sylloge Fungorum Vol. I: 741—753.

Tandon, R. N. & Sudhir, Chandra, 1963—1964: Supplement to the list of Indian fungi (1957—1962), p. 166. Allahabad University.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1969/1970

Band/Volume: [23](#)

Autor(en)/Author(s): Kale S. B.

Artikel/Article: [The Genus Valsaria in India. 194-197](#)