

## Notes on Taxonomy and Biogeography of *Rozites* KARSTEN

E. HORAK

Geobotanical Institute, ETHZ, CH-8092 Zürich, Switzerland

**Summary.** — Two new species of *Rozites* KARSTEN (1879; ep. HORAK, 1968) are reported (*R. similis* from Papua New Guinea, and *R. neocaledonica* from New Caledonia). All 14 taxa now referred to *Rozites* are keyed out and their respective spores are illustrated. The biogeography of *Rozites* is discussed and the area of distribution of the species is mapped.

### Introduction

The original circumscription of *Rozites* KARSTEN (1879) is exclusively based upon *Rozites caperata* (Fr.) KARSTEN which is restricted to the temperate regions of Eurasia and North America. Subsequently the genus remained monotypic until CLELAND & CHEEL (1918) described a second species allotted to *Rozites* (*R. australiensis*; see below).

After a revision of several taxa described under *Rozites* MOSER (1953) introduced 2 further species in this genus, that is 4 species have been recognized. Upon critical re-examination one of those taxa had to be transferred to *Descolea* [*D. flavoannulata* (VASILIEVA) HORAK, 1971]. In addition the macroscopic and microscopic data reported and found on *R. australiensis* CLELAND & CHEEL (1918) indicate that this agaric rather belongs to *Cortinarius* (*Phlegmacium*) than to *Rozites*. Finally SINGER (1975) recognized only *Rozites caperata* (Fr.) and *R. emodensis* (Berk.) and the results of the present study fully support SINGER's opinion.

Recently several contributions on *Rozites* in *Nothofagus* forests of the southern hemisphere have been published. To present knowledge 5 taxa occur in southern South America (MOSER & HORAK, 1975; HORAK, 1979) and 5 species are recorded from New Zealand (HORAK & TAYLOR, 1981).

The present pattern of distribution (Fig. 6) supports the hypothesis that *Rozites* apparently evolved on the southern hemisphere (Argentina—Chile: 5 spp., New Zealand: 5 spp., New Caledonia 1 sp., New Guinea 1 sp.). The two taxa so far recorded from the northern hemisphere (*R. caperata*, *R. emodensis*) — or their ancestors — seem to have migrated from the Gondwanian territories to the North via the Indomalayan archipelago. In this connection *R. similis* (the name

emphasises its relationship to *R. caperata*) probably plays a key role in the relationships between the northern and southern taxa. In New Guinea *R. similis* enters obviously ectotrophic mycorrhiza with *Castanopsis* and *Lithocarpus* (both belonging to the Fagaceae). Since the two host trees (of northern origin) reach their southern limit in New Guinea the occurrence of *R. similis* can safely be predicted elsewhere in the Indomalayan region and in the Far East where *Castanopsis* and *Lithocarpus* locally predominate in deciduous forests. In New Guinean *Nothofagus* forests no species of *Rozites* has been recorded yet but *R. neocaledonica* gathered under *Nothofagus* on neighbouring New Caledonia is clearly a close relative of *R. similis*. Further field-work will demonstrate whether or not *R. neocaledonica* is also present in the *Nothofagus* woods of New Guinea which — needless to say — here and there are in direct contact with oak forests supporting *R. similis*.

Type material of the new species is kept in ZT. If not otherwise stated the magnifications of the figures are: carpophores (nat. size), spores ( $\times 2000$ ), and basidia ( $\times 1000$ ).

#### Key to Species of *Rozites* KARSTEN

1. Carpophores (pileus, young lamellae, context of stipe and/or veil remnants) with lilac to blue colours, at least in young specimens . . . . . 2
- 1\*. Carpophores without conspicuous lilac or blue colours (cp. *R. caperata* with pale lilac veil remnants) . . . . . 8
2. Pileus dry, wrinkled in aged specimens (cp. also *R. caperata*, *R. rugosiceps*); stipe cylindrical-equal to subbulbous . . . . . 3
- 2\*. Pileus glutinous or viscid . . . . . 4
3. Pileus — 65 mm, ochre-brown to orange-brown, with scattered concolorous or ochre squamules or patches from universal veil; lamellae pale blue turning rust brown; stipe — 110  $\times$  — 12 mm, pale blue (at least at apex), whitish to ochre-brown in mature specimens; annulus white to pale ochre, membranous, substriate; several incomplete, submembranous zones of universal veil towards base; context distinctly blue to lilac in apex of stipe; spores 11—14.5  $\times$  7—8  $\mu\text{m}$ . On soil under *Nothofagus* spp. New Caledonia . . . . . 4. *R. neocaledonica*
- 3\*. Pileus — 80 mm, ochre-brown with distinct red-brown tinge over disc, densely covered with whitish, small squamules from universal veil; lamellae pale argillaceous (sometimes with pale lilac tinge in young specimens) becoming pale rust brown; stipe — 110  $\times$  — 15 mm, white; annulus white, striate; several incomplete submembranous zones or belts from universal veil towards base; context pale lilac in apex of stipe; spores 11—

- 13×7—8 μm. On soil under *Castanopsis* spp. and *Lithocarpus* spp. Papua New Guinea ..... 3. *R. similis*
4. Stipe —150×—20 (—45) mm, often robust, with rooting base, rarely cylindrical, whitish ochre below blue apex; anulus membranous striate, white; towards base with several, incomplete, submembranous, white to ochraceous belts from universal veil; pileus —120 mm, red brown or liver brown, margin with ochraceous remnants of veil; lamellae blue turning rust brown; context lilac to pale blue; spores 11—13×5—7 μm. On soil under *Nothofagus* spp. Chile, Argentina .. 10. *R. collarata*
- 4\*. Stipe cylindrical to subbulbous, base not rooting ..... 5
5. Young lamellae argillaceous ..... 6
- 5\*. Young lamellae lilac to blue ..... 7
6. Pileus —80 mm, argillaceous to ochre, disc often reddish, fibrillose or appendiculate, lilac veil remnants towards or on margin; stipe —110×—12 (—25, at base) mm, lilac to pale blue changing to whitish with age; anulus fibrillose to membranous, lilac, persistent, towards base with 1 to 3 incomplete, lilac belts of universal veil; odour none; spores 10—12.5×6—7 μm. On soil under *Nothofagus* spp. Chile, Argentina... .. 11. *R. ochraceoazurea*
- 6\*. Pileus —100 mm, lilac to purple changing to hazel brown or brown, margin appendiculate from whitish to pale lilac veil remnants; stipe —100×—20 mm, white to pale brown; anulus white to pale lilac, membranous, not striate, subpersistent, towards base with several incomplete, concolorous, appressed or submembranous belts from universal veil; odour strong, ranging from sweet to unpleasant (like coal gas); spores 9—11×5—5.5 μm; in subhymenium with conspicuous, yellow-brown (KOH) oleiferous hyphae. On soil under *Nothofagus* spp. Chile, Argentina ..... 12. *R. violacea*
7. Pileus —85 mm, grey to pale grey-brown with lilac tint, densely covered with white to pale yellow-brown, coarse, fibrillose squamules from universal veil; stipe —120×—20 mm, apex lilac, pale brown towards base; anulus pale ochre, striate, membranous, persistent; towards base with numerous, ochre, fibrillose or recurved scales from universal veil; context lilac in upper portion of stipe; spores 10—12.5×7.5—9.5 μm. On soil under *Nothofagus* spp. New Zealand ..... 5. *R. meleagris*
- 7\*. Pileus —90 mm, brown, with white, floccose veil remnants; stipe —190×—30 mm, pale brown; anulus whitish, membranous, substriate, persistent; spores 13—16×8—9.5 μm. On soil under *Abies webbiana*. India (Sikkim)... 2. *R. emodensis*
- 8 (2\*). Stipe —150×—14 mm, fusoid with rooting base, white changing to pale yellow-brown; anulus white, evanescent;

- towards base with inconspicuous, white, fibrillose zones from universal veil; spores  $11-15 \times 6-7.5 \mu\text{m}$ . On soil under *Nothofagus* spp. New Zealand ..... 6. *R. fusipes*
- 8\*. Stipe cylindric, base equal, bulbous or submarginate ..... 9
9. Pileus —90 mm, centre conspicuously wrinkled or veined, yellow-brown to red-brown; stipe —80×—16 mm, whitish turning pale brown; annulus whitish, striate, membranous, persistent; towards base with whitish to reddish brown fibrils or scales from universal veil; spores  $11-15 \times 7-8 \mu\text{m}$ , strongly warted and warts often confluent to short crests. On soil under *Nothofagus* spp. New Zealand ..... 7. *R. rugosiceps*
- 9\*. Pileus not conspicuously wrinkled (except *R. caperata*), spores covered by small to minute, isolated warts ..... 10
10. Pileus —55 mm, whitish, grey or pale argillaceous, margin translucently striate, white squamules of universal veil “swimming” in gluten; stipe —80×—15 mm, white; annulus white, membranous, striate, persistent; towards base with several, inconspicuous, white zones of universal veil; spores  $9.5-12 \times 6-7.5 \mu\text{m}$ . On soil under *Nothofagus* spp. New Zealand ..... 8. *R. pallida*
- 10\*. Pileus yellowish, yellow-brown, orange, or date brown, margin not (or indistinctly only) striate ..... 11
11. Pileus argillaceous, yellowish or pale ochre-brown; cuticle viscid ..... 12
- 11\*. Pileus rich orange-brown, date brown or fuscous, often with red-brown tinge; cuticle glutinous..... 13
12. Pileus —60 mm, scattered white veil remnants on striate margin; stipe —65×—16 mm, white; annulus white, membranous, striate, subsistent; towards base with few, white, incomplete zones of universal veil; spores  $11-14 \times 6.5-8 \mu\text{m}$ . On soil under *Nothofagus* spp. Argentina, Chile. . 13. *R. sarmienti*
- 12\*. Pileus —120 mm, centre and/or margin hoary from lilac veil remnants, margin often wrinkled; stipe —120×—25 mm, pale ochraceous; annulus whitish, submembranous, smooth or substrate, often dehiscent; towards base with fugaceous, fibrillose, lilac to whitish veil remnants; spores  $10-13 \times 6.5-8 \mu\text{m}$ . On acid soil in coniferous and deciduous forests. Temperate regions of northern hemisphere ..... 1. *R. caperata*
13. Pileus —45 mm, conspicuous universal veil remnants absent; stipe —60×—8 mm, yellowish; annulus yellowish to ochre, striate, membranous, persistent; stipe —60×—8 mm, yellow; annulus yellow, membranous, striate, persistent; towards base with several, yellow-brown, fibrillose to submembranaceous squamules or belts from universal veil; spores  $9.5-12.5 \times$

- 5.5–7  $\mu\text{m}$ . On soil under *Nothofagus* spp. Argentina . . . . .  
 . . . . . 14. *R. gamundiae*  
 13\*. Pileus —60 mm often covered with conspicuous, whitish to reddish brown lumps of veil; stipe —80 $\times$ —15 mm, pale ochre-brown; annulus membranous, pale ochre-brown, striate, persistent; spores 9–12 $\times$ 5–7  $\mu\text{m}$ . On soil under *Nothofagus* spp. New Zealand. . . . . 9. *R. castanella*

### Descriptions and Annotations to Species

1. *Rozites caperata* (FRIES) KARSTEN 1879 — Fig. 1, 1  
 Bidr. Känn. Finl. Nat. Folk 32: 290.  
 Bas. *Agaricus caperatus* FRIES 1821: Syst. Myc. 1: 241.  
 Syn. Cp. MOSER (1953: 167).  
 Illustrations. — HORAK (1968).  
 Habitat and distribution. — On acid soil in coniferous (*Picea*) and deciduous (*Fagus*, *Quercus*) forests, often associated with *Vaccinium* spp. — Temperate region of northern hemisphere.
  
2. *Rozites emodensis* (BERKELEY) MOSER 1963 — Fig. 2  
 Schw. Zeitschr. Pilzkunde 31: 169.  
 Bas. *Cortinarius (Myxaciium) emodensis* BERKELEY 1852: Hook. J. Bot. 4: 132.  
 Illustrations. — MOSER (1953: 171).  
 Habitat and distribution. — On soil under *Abies webbiana* (about 10.000 feet a. s. l.). — Sikkim.  
 Material. — India: Sikkim, Lachen, 31. V. 1849, leg. HOOKER (K, holotype).
  
3. *Rozites similis* HORAK sp. n. — Fig. 3  
 Pileus —80 mm, ex convexo umbonatus, ochraceo-brunneus castaneo tinctu, siccus vel subviscidus, marginem versus subrugosus, squamis albis minutisque e velo dense obtectus. Lamellae sublilacinae dein argillaceae, ex adnato emarginatae, crenulatae. Stipes —110 $\times$ —15 mm, cylindricus vel subclavatus, albus, annulus albus, striatus, membranaceus, persistens, basim versus zonis albis cingulatus. Caro lilacina. Sporae 11–13 $\times$ 7–y  $\mu\text{m}$ , amygdaliformes, verrucosae. Septa fibulata. Ad terram in silvis fagineis. Nova Guinea. Typus, ZT, 72/645.  
 Pileus —80 mm, hemispheric to convex when young, margin strongly incurved, becoming umbonate-expanded; orange-brown with distinct red-brown tinge (especially over disc), towards margin yellow-brown in aged specimens; subviscid when moist, soon dry, margin occasionally striate but always wrinkled; all over covered with small, fibrillose, white or grey squamules of the universal veil, margin sometimes appendiculate with lumps of veil. Lamellae 26–36, —7, crowded, adnate to emarginate; very pale lilac in young and fresh




Fig. 1. Spores: 1. *Rozites caperata* (FR.) KARSTEN (ZT, 79/349). — 2. *Rozites ochraceoazurea* (HORAK) HORAK (type). — 3. *Rozites fusipes* HORAK & TAYLOR (type). — 4. *Rozites meleagris* HORAK & TAYLOR (type). — 5. *Rozites rugosiceps* HORAK & TAYLOR (type). — 6. *Rozites castanella* HORAK & TAYLOR (type). — 7. *Rozites sarmienti* (SPEGAZZINI) HORAK (type). — 8. *Rozites pallida* HORAK & TAYLOR (type). — 9. *Rozites collarata* (HORAK & MOSER) HORAK (type). — 10. *Rozites gamundiae* HORAK (type).

specimens, soon turning argillaceous to pale ferruginous, edge paler, crenulate. Stipe —  $110 \times 15$  mm, cylindric, base swollen or bulbous; pale lilac at apex turning white; strongly fibrillose, dry, solid, single, in groups; annulus white, striate, submembranous often dehiscent or incomplete, towards base with several, white, incomplete zones or squamules of universal veil. Odour and taste not distinctive.




Fig. 2. *Rozites emodensis* (BERKELEY) MOSER (type): carpophores ( $\frac{1}{2}$  nat. size), spores

Context orange beneath cuticle of pileus, lilac in apex of stipe, ochraceous in base of stipe. Chemical reactions on pileus: KOH, HCl and  $\text{NH}_3$  — negative.

Spores  $11-13 \times 7-8 \mu\text{m}$ , amygdaliform to pip-shaped, rust brown, verrucose, plage and perispore absent. Basidia  $35-45 \times 9-12 \mu\text{m}$ , 4-spored. Cheilo- and pleurocystidia absent. Cuticle a cutis of repent, cylindric hyphae ( $3-8 \mu\text{m}$  diam.), subcutis composed of ovoid to globose cells, strongly encrusted by brown (KOH) pigment. Clamp connections present.

**Habitat and distribution.** — On soil in forests dominated by *Castanopsis* spp. and *Lithocarpus* spp., 1000–1400 m a. s. l. — Papua New Guinea.

**Material.** — Papua New Guinea: Morobe district: Bulolo, Manki, 20. XI. 1972, leg. HORÁK (ZT, 72/645, holotype). — Bulolo, Heads Hump, 4. XI. 1971, leg. HORÁK (ZT, 71/252).




Fig. 3. *Rozites similis* HORÁK (type): carpophores, spores

4. *Rozites neocaledonica* HORÁK sp. n. — Fig. 4

Pileus — 65 mm, e convexo umbonatus, laete ochraceus, siccus, marginem versus subrugosus, interdum squamis subochraceis e velo obtectus. Lamellae lilacinae dein argillaceo-ferrugineae, ex adnato-emarginatae, crenulatae. Stipes — 110 × — 12 mm, cylindricus, aequalis, primo lilacinus dein ochraceo-brunneus, anulus albus vel pallide ochraceus, striatus, membranaceus, persistens, basim versus zonis ochraceo-albis nonnullis cingulatus. Caro lilacina. Sporae 11—



14,5 × 7–8  $\mu$ m, amygdaliformes, verrucosae. Ad terram in silvis nothofagineis. Nova Caledonia. Typus, ZT, 77/34.

Pileus — 65 mm, hemispheric or convex with incurved margin when young becoming umbonate-expanded, occasionally margin up-turned; ochre to ochre-brown, red-brown tints absent; dry, margin wrinkled; surface covered with silvery or pale ochraceous squamules or lumps from universal veil, remnants fugaceous hence absent in




Fig. 4. *Rozites neocaledonica* HORÁK (type): carpophores, spores

aged carpophores. Lamellae crowded, adnate to emarginate; at first lilac to pale blue turning argillaceous or ferruginous; edge paler, crenulate; up to 5 mm wide. Stipe — 110 × 12 mm, cylindric, equal; pale blue or lilac all over when young changing to pale brown or pale ochre-brown (especially towards base); dry, coarsely fibrillose, solid becoming fistulose with age; annulus white or pale ochraceous, membranous, striate, persistent; towards base with several, incon-

spicuous, white to ochre, incomplete zones or belts of universal veil; single, in groups. Odour and taste not distinctive. Context white in pileus, lilac in upper portion of stipe, pale orange in base of stipe. Chemical reactions on pileus: KOH — brown.

Spores  $11-14.5 \times 7-8 \mu\text{m}$ , amygdaliform (to elliptic), rust brown, verrucose, plage and perispore absent. Basidia  $25-35 \times 10-12 \mu\text{m}$ , 4-spored. Cheilocystidia  $20-65 \times 4-10 \mu\text{m}$ , fusoid with tapering neck, membrane hyaline, pigment absent, often scattered on edge. Pleurocystidia none. Cuticle a cutis of repent cylindrical hyphae ( $2-8 \mu\text{m}$  diam.), subcutis of ovoid cells, strongly encrusted with red-brown (KOH) pigment. Septa with clamp connections.

Habitat and distribution. — On soil among litter in forests dominated by *Nothofagus* spp., 1000—1200 m a. s. l. — New Caledonia.

Material. — New Caledonia: Paita, Mt. Mou, 22. II. 1977, leg. HORAK (ZT, 77/34, holotype). — Same locality, 20. II. 1977, leg. HORAK (ZT, 77/10).

5. *Rozites meleagris* HORAK & TAYLOR 1981 — Fig. 1, 4

New Zealand J. Botany (in press).

Illustrations. — HORAK & TAYLOR (1981: l. c.).

Habitat and distribution. — On soil under *Nothofagus* spp. — New Zealand.

Material. — New Zealand: Canterbury, E of Lewis Pass, 23. III. 1968, leg. HORAK (PDD, 27164, holotype). — For further collections studied cp. HORAK & TAYLOR (1981).

6. *Rozites fusipes* HORAK & TAYLOR 1981 — Fig. 1, 3

New Zealand J. Botany (in press).

Illustrations. — HORAK & TAYLOR (1981: l. c.).

Habitat and distribution. — On soil under *Nothofagus* spp. — New Zealand.

Material. — New Zealand: Nelson, Lake Rotoiti, Mt. Robert, 2. V. 1968, leg. HORAK (PDD, 27162, holotype). — For further collections examined cp. HORAK & TAYLOR (1981).

7. *Rozites rugosiceps* HORAK & TAYLOR 1981 — Fig. 1, 5

New Zealand J. Botany (in press)

Illustrations. — HORAK & TAYLOR (1981: l. c.).

Habitat and distribution. — On soil under *Nothofagus* spp. — New Zealand.

Material. — New Zealand: Otago, Haast Pass, Makarora, 16. IV. 1965, leg. TAYLOR (Herb. TAYLOR, 240, holotype). — For further collections examined cp. HORAK & TAYLOR (1981).

8. *Rozites pallida* HORAK & TAYLOR 1981 — Fig. 1, 8

New Zealand J. Botany (in press)

Illustrations. — HORAK & TAYLOR (1981: l. c.).

Habitat and distribution. — On soil under *Nothofagus* spp. — New Zealand.

Material. — New Zealand: Nelson, Lake Rotoiti, St. Arnaud Range, 23. V. 1968, leg. HORAK (PDD, 27163, holotype). — For further collections examined cp. HORAK & TAYLOR (1981).

9. *Rozites castanella* HORAK & TAYLOR 1981 — Fig. 1, 6

New Zealand J. Botany (in press).

Illustrations. — HORAK & TAYLOR (in press).

Habitat and distribution. — On soil under *Nothofagus* spp. — New Zealand.

Material. — New Zealand: Gisborne, Urewera N.P., Ngamoko, 27. VI. 1968, leg. HORAK (PDD, 27161, holotype). — For further collections examined cp. HORAK & TAYLOR (1981).

10. *Rozites collariata* (HORAK & MOSER) HORAK 1979 — Fig. 1, 9

Fl. Criptogamica Tierra del Fuego 11: 345.

Bas. *Cortinarius collariatus* HORAK & MOSER ap. MOSER & HORAK 1975: Nova Hedwigia, Beih. 52: 234.

Illustrations. — MOSER & HORAK (1975: l. c.); HORAK (1979: l. c.).

Habitat and distribution. — On soil under *Nothofagus* spp. — Argentina, Chile.

Material. — Argentina: Neuquen: Pto. Manzano, Lago Nahuel Huapi, 19. III. 1963, leg. MOSER (IB, 63/88, holotype). — Chubut: P. N. Los Alerces, Lago Verde, 29. III. 1980, leg. HORAK (ZT, 339). — For further collections examined cp. HORAK (1979: 345).

11. *Rozites ochraceoazurea* (HORAK) HORAK 1979 — Fig. 1, 2

Fl. Criptogamica Tierra del Fuego 11: 348.

Bas. *Thaxterogaster ochraceoazureus* HORAK ap. HORAK & MOSER 1965: Nova Hedwigia 10: 237.

Syn. *Cortinarius ochroianthinus* HORAK & MOSER ap. MOSER & HORAK 1975: Nova Hedwigia, Beih. 52: 274.

*Cortinarius gliocyclus* HORAK ap. MOSER & HORAK 1975: Nova Hedwigia, Beih. 52: 275.

Illustrations. — HORAK & MOSER (1965: l. c.); MOSER & HORAK (1975: l. c.); HORAK (1979: l. c.).

Habitat and distribution. — On soil under *Nothofagus* spp. — Argentina, Chile.

**Material.** — Argentina: Tierra del Fuego: Ushuaia, Valle Glaciar Martial, 2. III. 1963, leg. HORAK (ZT, 64/22, holotype). — Chubut: P. N. Los Alerces, Lago Verde, 29. III. 1980, leg. HORAK (ZT, 336). — For further collections examined cp. HORAK (1979: 348).

12. *Rozites violacea* HORAK ap. MOSER & HORAK 1975 — Fig. 5

Nova Hedwigia, Beih. 52: 516.

Syn. *Rozites purpurea* MOSER ap. MOSER & HORAK 1975, Nova Hedwigia, Beih. 52: 514.

Illustrations. — MOSER & HORAK (1975: l. c.).




Fig. 5. *Rozites violacea* ap. MOSER & HORAK: 1. carpophore, spores (type). — 2. carpophore (ZT, 412). — 3. spores (type of *Rozites purpurea* MOSER)

**Habitat and distribution.** — On soil under *Nothofagus* spp. — Argentina.

**Material.** — Argentina: Rio Negro, Pto. Blest, El Albuelo, 4. IV. 1962, leg. HORAK (IB and ZT, 70/353, holotype). — Same locality, 8. IV. 1980, leg. HORAK (ZT, 412). — Neuquen, Lago Nahuel Huapi, Brazo Rincon, 18. III. 1973, leg. MOSER (IB, 63/80, holotype of *Rozites purpurea* MOSER). — For further collections studied cp. MOSER & HORAK (1975: l. c.).

13. *Rozites sarmienti* (SPEGAZZINI) HORAK 1979 — Fig. 1, 7  
Fl. Criptogamica Tierra del Fuego 11: 350.  
Bas. *Cortinarius sarmienti* SPEGAZZINI 1887: Bol. Acad. Nac. Cs. Cordoba  
11: 151.  
Syn. *Cortinarius togularis* HORAK ap. MOSER & HORAK 1975: Nova  
Hedwigia, Beih. 52: 277.  
Illustrations. — HORAK (1979: l. c.).  
Habitat and distribution. — On soil under *Nothofagus* spp. —  
Argentina, Chile.  
Material. — Chile: Magallanes: Monte Sarmiento, V. 1882, leg.  
SPEGAZZINI (LPS, 38971, holotype). — Osorno: Refugio Antillanca  
(Volcan Antillanca), 22. IV. 1975, leg. HORAK (ZT, 75/407). — For  
further collections examined cp. HORAK (1979: l. c.).
14. *Rozites gamundiae* HORAK 1979 — Fig. 1, 10  
Fl. Criptogamica Tierra del Fuego 11: 346.  
Illustrations. — HORAK (1979: l. c.).  
Habitat and distribution: On soil under *Nothofagus* spp. —  
Argentina.  
Material. — Argentina: Tierra del Fuego, Ushuaia, Tierra  
Mayor, 5. III. 1974, leg. HORAK (LPS, 37861, holotype). — For  
further collections examined cp. HORAK (1979: l. c.).

#### Excluded Species

1. *Rozites australiensis* CLELAND & CHEEL 1918  
Trans. Proc. Roy. Soc. S. Australia 42: 90.  
= *Cortinarius (Phlegmacium) australiensis* (CLEL. & CHEEL)  
HORAK comb. nov.  
Material. — Australia: South Australia, Mt. Lofty, 7. IV. 1917,  
leg. CLELAND (ADW 13692, holotype). — ACT: Canberra, Brindabella  
Range (Fauna Reserve Area), 24. III. 1972, leg. SHEPHERD, 776  
(CANB 226784).  
Remarks. — The revision of the type material (in addition with  
related taxa from New Zealand and Papua New Guinea) indicates  
that *R. australiensis* belongs to *Cortinarius* subgen. *Phlegmacium*.  
For additional information consult CLELAND (1934: 100), MOSER  
(1953: 170) and WILLIS (1963: 30).
2. *Rozites coerulea* MOSER ap. MOSER & HORAK 1975  
Nova Hedwigia, Beih. 52: 518.  
= *Stephanopus coerulea* (MOSER) HORAK comb. nov.  
Material. — Argentina: Neuquen, Pto. Blest, Los Cantaros,  
15. III. 1959, leg. SINGER, M 1816 (IB, holotype).




Fig. 6. Present distribution of *Nothofagus* spp. (black areas) and *Rozites* (numbers follow enumeration in text): 1. *R. caperata* (approx. southern limit on the northern hemisphere). — 2. *R. emodensis*. — 3. *R. similis*. — 4. *R. neocaledonica*. — 5. *R. meleagris*. — 6. *R. fusipes*. — 7. *R. rugosiceps*. — 8. *R. pallida*. — 9. *R. castanella*. — 10. *R. collarata*. — 11. *R. ochraceoazurea*. — 12. *R. violacea*. — 13. *R. sarmienti*. — 14. *R. gamundiae*

Remarks. — The critical re-examination of the amygdaliform spores demonstrated a distinct plage. Hence this species is transferred from *Rozites* to *Stephanopus* MOSER & HORAK (1975) taxonomically being a close relative of *St. azureus* MOSER & HORAK (1975).

3. *Rozites flavoannulata* VASILIEVA 1950

Bot. Mat. Inst. spor. Rast. 6: 199.

= *Descolea flavoannulata* (VASILIEVA) HORAK 1971. Persoonia 6: 246.

Remarks. — According to YOKOHAMA & al. (1979) this species is widely distributed in Eastern Siberia, Korea and Japan.

### Acknowledgements

I am grateful to Prof. Dr. M. MOSER (Innsbruck, Austria) for the access to yet unpublished data on *Rozites* and to the Curators in K and ADW for the loan of type material and pertinent notes kept in those herbaria.

My thanks are also expressed to the Swiss Society of Natural Sciences for a travelling grant to Indomalaya and Australasia. During this collecting trip material of *Rozites* has been gathered in New Caledonia.

### References

- CLELAND, J. B. (1934). Toadstools and mushrooms and other larger fungi of South Australia, vol. 1, 178 pp. — Gov. Printer, Adelaide.
- & CHEEL, E. (1918). Australian fungi: notes and descriptions. No. 1. — Trans. Proc. Roy. Soc. S. Australia 42: 88—138.
- HORAK, E. (1968). Synopsis generum Agaricalium. — Beitr. Kryptogamen Fl. Schweiz 13: 1—741.
- (1971). Studies on the genus *Descolea* SING. — Persoonia 6: 231—248.
- (1979). Fungi, Basidiomycetes: Agaricales y Gasteromycetes secotioides. — Fl. Criptogam. Tierra del Fuego 11: 1—524.
- & TAYLOR, M. (1981). Fungi Agaricini Novazelandiae. XI. *Rozites* KARSTEN. — N. Z. J. Bot. (in print).
- MOSER, M. (1953). Die Gattung *Rozites* KARSTEN. — Schweiz. Zeitschr. Pilzkunde 31: 164—172.
- & HORAK, E. (1975). *Cortinarius* Fr. und nahe verwandte Gattungen in Südamerika. — Nova Hedwigia Beiheft 52: 1—628.
- SINGER, R. (1975). The Agaricales in modern taxonomy. — Cramer, Vaduz (912 pp., 3rd ed.).
- WILLIS, J. H. (1963). Victorian toadstools and mushrooms. — 88 pp., Field Naturalists Club of Victoria, Melbourne (3rd ed.).
- YOKOHAMA, T., YONG HWAN PARK, YANG SUP KIM, BYONG KAK KIM & HONGO, T. (1979). Distribution of *Descolea flavoannulata* (VASILIEVA) HORAK in Far Eastern Asia. — Trans. Myc. Soc. Japan 20: 63—72.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1981

Band/Volume: [34](#)

Autor(en)/Author(s): Horak Egon

Artikel/Article: [Notes on Taxonomy and Biogeography of Rozites  
KARSTEN. 94-108](#)