

## Aquatic Fungi of Kumaun Himalaya, India: *Pythium* PRINGSHEIM

G. S. MER, B. L. VERMA & R. D. KHULBE

Department of Botany, Kumaun University, Nainital-263 002, India.

**Abstract.** — A total of twenty one isolates belonging to different species of *Pythium* were isolated and studied from water and soil of Kumaun Himalaya, India. Of these, one species viz., *P. pulchrum* MINDEN is a new record for India and ten species viz., *P. aphanidermatum* (EDSON) FITZ., *P. artotrogus* de BARY, *P. debaryanum* HESSE, *P. dissotocum* DRECHSLER, *P. echinulatum* MATTHEWS, *P. gracile* SCHENK, *P. middletonii* SPARROW, *P. paroeacandrum* DRECHSLER, *P. spinosum* SAWADA ap. SAWADA & CHEN and *P. ultimum* TROW are being reported from Kumaun Himalaya for the first time.

### Introduction

In India BUTLER (1907) was the first who reported eight species of *Pythium*. After his work, some of the important contributions on the study of *Pythium* have been made by a number of workers including DASTUR (1935), BHARGAVA (1941), BALKRISHNAN (1948 a, b), RAMKRISHNAN (1949), MAHMUD (1950, 1951, 1952), PRAKASH & SAKSENA (1952), SHARMA & NEEMA (1952), RAMKRISHNAN (1955), SRINIVASAN (1956), DAYAL & TANDON (1962), RAO (1963), DESHPANDE & DESHPANDE (1964), SRIVASTAVA (1964), BHARGAVA & SINGH (1965), RAJGOPALAN & RAMKRISHNAN (1967—68), AULAKH (1971), PRABHUJI & SRIVASTAVA (1976), MANOHARACHARY & RAO (1978), KHULBE (1979), CHOWDHRY & AGARWAL (1981), MANOHARACHARY (1981) and MER & KHULBE (1983). But the Kumaun region in the Himalayas remained almost unexplored. In the present pursuit, different representatives of twenty one species of *Pythium* were isolated and studied from water, soil and infected roots of crops over a period of three years.

### Materials and Methods

Samples of water, soil and infected roots of different crops were collected from a number of selected sites in the Kumaun Himalaya. Isolation of species was exercised by following the method used by ROBERTSON (1980).

Subsequently, the isolates were grown on hempseed halves in sterile distilled water. Preserved specimens of all the species have been deposited in the collections in the museum of Botany Department, Kumaun University, Nainital. Moreover, a few important taxa viz. *Pythium echinulatum* MATTHEWS (IMI 255016), *Pythium gracile* SCHENK (IMI 255013), *P. hypogynum* MIDDLETON (IMI 277403), *P. middletonii* SPARROW (IMI 255019

& 255018), *P. rostratum* BUTLER (IMI 277406), *P. spinosum* SAWADA ap. SAWADA & CHEN (IMI 277405) and *P. vexans* var. *minuta* MER & KHULBE (IMI 255017) have been deposited in the Herbarium, CMI, Kew, England.

**Key to the Himalayan species of *Pythium* PRINGSHEIM**

1. Sporangia filamentous, irregularly lobulate or digitate ..... 3
2. Sporangia spherical, ovoid or ellipsoidal ..... 4
  - 3a. Sex organs present ..... 5
  - 3b. Sex organs absent ..... *P. afertile* (1)
- 4a. Sex organs present ..... 6
- 4b. Sex organs absent ..... 7
  - 5a. Antheridia strictly declinous ..... 8
  - 5b. Antheridia declinous and androgynous ..... 9
  - 5c. Antheridia declinous and monoclinal ..... 10
- 6a. Oogonia smooth walled ..... 11
- 6b. Oogonia spiny ..... 12
  - 7a. Hyphae undulating, sporangia ellipsoidal .. *P. undulatum* (20)
  - 7b. Hyphae not undulating, sporangia not ellipsoidal .....
    - ..... *P. elongatum* (7)
- 8a. Oospore aplerotic ..... *P. gracile* (8)
- 8b. Oospore plerotic ..... *P. inflatum* (10)
  - 9a. Oospore aplerotic ..... *P. dissotocum* (5)
  - 9b. Oospore plerotic ..... *P. monospermum* (12)
- 10a. Oospore aplerotic ..... *P. aphanidermatum* (2)
- 10b. Oospore plerotic ..... *P. torulosum* (18)
  - 11a. Antheridia sessile ..... *P. ultimum* (19)
  - 11b. Antheridia not as above ..... 13
- 12a. Antheridia monoclinal, declinous and hypogynous in addition to androgynous ..... 14
- 12b. Antheridia strictly hypogynous ..... *P. artotrogum* (3)
  - 13a. Antheridia declinous ..... 15
  - 13b. Antheridia monoclinal ..... 16
  - 13c. Antheridia strictly hypogynous ..... 17
  - 13d. Antheridia bell-shaped, hypogynous as well as declinous .....
    - ..... *P. vexans* var. *minutum* (21)
- 14a. Oospore aplerotic ..... *P. echinulatum* (6)
- 14b. Oospore plerotic ..... *P. spinosum* (17)
  - 15a. Oogonia terminal, oospore single ..... *P. debaryanum* (4)
  - 15b. Oospore 1—2 ..... *P. middletonii* (11)
- 16a. Oogonia usually intercalary ..... *P. parocandrum* (13)
- 16b. Oogonia not intercalary ..... 18
  - 17a. Oospore plerotic ..... *P. hypogynum* (9)
  - 17b. Oospore aplerotic ..... *P. pulchrum* (15)
- 18a. Oospore plerotic ..... *P. rostratum* (16)
- 18b. Oospore aplerotic ..... *P. proliferum* (14)

1. *Pythium afertile* KANOUSE & HUMPHREY in Papers Mich. Acad. 8: 129—140, 1927. — Pl.1, fig. 1

Principal hyphae slender, 3—11.2  $\mu\text{m}$  thick at base and 3—3.7  $\mu\text{m}$  at the tip; zoosporangia filamentous, simple or slightly branched, 41—250  $\mu\text{m}$  long and 3—95  $\mu\text{m}$  in diameter; vesicle spherical, sessile, 18—45  $\mu\text{m}$  in diameter, mostly 30  $\mu\text{m}$ ; primary zoospores 8—25 in number, 7.5—11  $\mu\text{m}$  in diameter; gemmae abundant, filamentous with bead-like structures, act as zoosporangia; sex organs not developed.

Isolated from Ram Tal water and soil.

This species was observed in India by KHULBE (1979).

2. *Pythium aphanidermatum* (EDSON) FITZPATRICK in Jour. Agri. Res. 4: 279—292, 1915. — Pl. 1, fig. 19, 20

Colony well developed on hempseed halves; hyphae non-septate, branched, 4—7  $\mu\text{m}$  in diameter; sporangia inflated, filamentous, terminal and sometimes lateral in position, 25—210  $\mu\text{m}$  in length and 8—19  $\mu\text{m}$  in diameter; zoospores reniform, formed within the vesicle, encysted primary zoospores 10  $\mu\text{m}$  in diameter; oogonia spherical, terminal, rarely intercalary, 21 to 28.5  $\mu\text{m}$  in diameter; antheridia monoclinal and declinous, one to two per oogonium; antheridial cells barrel shaped or dome shaped; oospore single, eccentric, aplerotic, 17—20.5  $\mu\text{m}$  in diameter.

Isolated from soil of Khatima.

In India, this species was observed by SUBRAMANIAM (1919), MITRA (1927), RAMKRISHNA (1929), MAHMUD (1950), GATTANI & KAUL (1951), JAIN (1952), SHARMA & NEMA (1952), BHARGAVA & GUPTA (1957), SHARMA & ASTHANA (1958), RAO (1963), RAGHUNATHAN (1968), RAMKRISHNAN & al. (1973), MUTHUSAMY & al. (1974), DIWAKAR & PAYAK (1975). This species is very close to the description given by WATERHOUSE (1967) with the exception in having zoosporangia of smaller size.

3. *Pythium artotrogum* (MONT.) de BARY in Abh. Senckenb. Naturf. Ges. 12: 225—340, 1881. — Pl. 2, fig. 1, 2

= *P. hydnosporum* (MONT.) SCHRÖTER in ENGL. & PRANTL, Nat. Pfl. 1: 104—105, 1897.

= *P. artotrogum* var. *macranthum* SIDERIS, Mycologia 24: 14—61, 1932.

Mycelium showing aerial development; hyphae branched, 4—8  $\mu\text{m}$  in diameter; conidia spherical to sub-spherical, 30—35  $\mu\text{m}$  in diameter, mostly intercalary; oogonia spiny, terminal, 23—28  $\mu\text{m}$  in diameter excluding the spines; antheridia unicellular, cylindrical to clubshaped, tapering, 10—23  $\mu\text{m}$  in length by 5—12  $\mu\text{m}$  in diameter, hypogynous; oospore echinulate or smooth walled, spherical, 18—23  $\mu\text{m}$  in diameter without the spines.

Isolated from soil of Ganai.

In India, this species is observed by BLATTER (1911), BUTLER (1907) and SYDOW & BUTLER (1907).

4. *Pythium debaryanum* HESSE in Inaug. Diss., Halle, 14—34, 1874. — Pl. 2, fig. 5

Principal hyphae branched, non septate, 4—6  $\mu\text{m}$  in diameter; sporangia terminal or intercalary, spherical to subspherical, 15—29  $\mu\text{m}$ , mostly 22  $\mu\text{m}$  in diameter; encysted primary zoospores 8—9  $\mu\text{m}$  in diameter; oogonia terminal or intercalary, spherical, 16—27  $\mu\text{m}$  in diameter, mostly 21  $\mu\text{m}$ ; antheridia monoclinal and declinal, one to four per oogonium; oospore single, eccentric, aplerotic, 12—18  $\mu\text{m}$  in diameter.

Isolated from rice roots and soil of Nanakmatta.

This species has been recorded in India by BUTLER (1913), RAMKRISHNAN (1941), VASUDEVA (1948), SINGH & GUPTA (1951), MAHMUD (1951), BHARGAVA & GUPTA (1957) and RAO (1963). The isolate shows almost complete agreement with the description given by RAO (1963).

5. *Pythium dissotocum* DRECHSLER in Jour. Wash. Acad. 20: 398—418, 1930. — Pl. 1, fig. 2, 3

Hyphae slender, branched moderately, 3—7.5  $\mu\text{m}$  in diameter, predominantly 3.7  $\mu\text{m}$ ; primary zoosporangia abundant, filamentous as well as lobulated, 28—125  $\mu\text{m}$  long and 3.7  $\mu\text{m}$  in diameter; gemmae abundant, lobulated, act as sporangia; oogonia abundant, terminal on short lateral branches, spherical, smooth walled, 13—23  $\mu\text{m}$  in diameter, predominantly 20  $\mu\text{m}$ ; antheridia androgynous and declinal, mostly androgynous, 1—3 in number per oogonium; oospore single, smooth walled, spherical, eccentric, aplerotic or nearly so, 10—19  $\mu\text{m}$  in diameter, mostly 16.5  $\mu\text{m}$ .

Isolated from Ram Tal soil and diseased roots and seedlings of *Trigonella foenum-graecum* L. near Nal-Damayanti Tal.


This species has been reported from India by CHOWDHRY & AGARWAL (1981).

6. *Pythium echinulatum* MATTHEWS in Univ. N. Carol. Press, 136 pp. 1931. — Pl. 1, fig. 12, 13

Hyphae long, slender, frequently branched, 3—11  $\mu\text{m}$  in diameter, predominantly 3—3.7  $\mu\text{m}$ ; zoosporangia terminal or intercalary, mostly oval, spherical, rarely sub-spherical, 20—30  $\mu\text{m}$  in diameter, germinate by 1—3 germ tubes; zoospores not observed; oogonia formed abundantly, terminal or on lateral branches or intercalary, spherical, oval or sub-spherical, predominantly spherical, 16—29  $\mu\text{m}$  in diameter, mostly 18—29  $\mu\text{m}$  exclusive of the spines; spines many, acute, 3.7—13  $\mu\text{m}$  long, predominantly 7.5—11  $\mu\text{m}$ ; antheridia mostly androgynous, rarely declinal and monoclinal, 1—3 per oogonia; oospores single, spherical, eccentric, aplerotic, 14—25  $\mu\text{m}$  in diameter, predominantly 19  $\mu\text{m}$ .

Isolated from Ram-Tal soil. The culture has been deposited in the herbarium CMI (IMI-255016), Kew, England.

This species has been reported from India by PRABHUJI & SRIVASTAVA (1976) and MANOHARACHARI & RAO (1978). This isolate differs from



Pl. 1: Fig. 1 (scale B): *Pythium afertile* KANOUSE & HUMPHREY. — Gemmae.  
 2, 3 (scale A): *P. dissotocum* DRECHSLER. — 2. Mature oogonium with androgynous antheridium. — 3. Mature oogonium with dichlinous antheridium.  
 4, 5 (scale B): *P. undulatum* PETERSON. — 4. Young sporangia. — 5. Formation of secondary sporangium by internal proliferation.  
 6 (scale C): *P. elongatum* MATTHEWS. — Mature zooporangium with zoospores in the vesicle.  
 7 (scale C): *P. monospermum* PRINGSHEIM. — Mature oogonium with dichlinous antheridium.  
 8 (scale D): *P. gracile* SCHENK. — Mature oogonium with dichlinous antheridium.  
 9, 10 (scale B): *P. pulchrum* MINDEN. — 9. Mature oogonium with hypogynous antheridium. — 10. Catenulate oogonia.

the description given by PRABHUJI (1979) in having the preponderance of androgynous antheridial branches.

7. *Pythium elongatum* MATTHEWS in Studies on the genus *Pythium*. Univ. N. Carol. Press 106: 1931. — Pl. 1, fig. 6

Hyphae long, branched, 3–6  $\mu\text{m}$  in diameter; sporangia terminal or intercalary, pyriform, cylindrical or spherical, 10–16  $\mu\text{m}$  in diameter; zoospores produced in a vesicle with a long basal tube; encysted zoospores 5–10  $\mu\text{m}$  in diameter; sex organs not developed.

Isolated from soil of Naina Peak, Nainital.

This species has been observed in India by MANOHARACHARY (1981) and KHULBE (1983).

8. *Pythium gracile* SCHENK in Verh. Phys. Med. Ges. Würzb. 9: 13–20, 1859. — Pl. 1, fig. 8

Hyphae slender, branched moderately, 3.5–7.5  $\mu\text{m}$  in diameter; zoosporangia mostly terminal and rarely lateral, filamentous, mostly branched, 375–800  $\mu\text{m}$  long and 7.5–18  $\mu\text{m}$  in diameter; zoospores formed within a vesicle at the tip of the zoosporangium, 7.5–12  $\mu\text{m}$  in diameter, 10–20 in number; gemmae abundant, branched, often deciduous, act as zoosporangia; oogonia lateral, abundant, formed intramatrically, spherical, 10.7–30  $\mu\text{m}$  in diameter, predominantly 16–26.5  $\mu\text{m}$ ; oogonial wall smooth; antheridia strictly declinuous, one to two per oogonium; oospores always maturing; oospore single, spherical and smooth walled, eccentric, aplerotic, 6.5–20.5  $\mu\text{m}$  in diameter, predominantly 13–16.5  $\mu\text{m}$ .

Isolated from Ram Tal water and gills of *Puntius ticto* L. from Naini Tal.

In India, this species was observed by SYDOW & BUTLER (1907). This species is being reported from the temperate region of India for the first time.

11 (scale D): *P. ultimum* Trow. — Mature oogonium with androgynous antheridium.

12, 13 (scale D): *P. echinulatum* MATTHEWS. — 12. Mature oogonium with androgynous antheridium and aplerotic oospore. — 13. Mature oogonium with hypogynous antheridium.

14, 15 (scale D): *P. spinosum* SAWADA ap. SAWADA & CHEN. — 14. Mature oogonium with plerotic oospore and androgynous antheridium. — 15. Oogonium with monoclinal antheridium.

16 (scale D): *P. rostratum* BUTLER. Mature oogonium with androgynous antheridium and plerotic oospore.

17, 18 (scale D): *P. inflatum* MATTHEWS. — 17. Mature oogonium with plerotic oospore and hypogynous antheridium. — 18. Oogonium with declinuous antheridia.

19, 20 (scale D): *P. aphanidermatum* (Eds.) FITZ. — 19. Mature oogonium with declinuous antheridium. — 20. Oogonium with androgynous antheridium.

21 (scale D): *P. torulosum* COKER & PATTERSON. — Mature oogonium with plerotic oospore and androgynous antheridium.

9. *Pythium hypogynum* MIDDLETON in Mem. Torrey bot. Club 20: 69, 1943. — Pl. 2, fig. 3

Hyp h a e long, non-septate, branched with luxuriant growth on hemp-seed, 2—8.5  $\mu\text{m}$  in diameter, mostly 5  $\mu\text{m}$ ; s p o r a n g i a terminal, spherical to oval, thin-walled, 8.5—37.5  $\mu\text{m}$  in diameter, mostly 22—30  $\mu\text{m}$ ; z o o s p o r e reniform, encysted zoospores 9—12  $\mu\text{m}$  in diameter; o o g o n i a terminal on short lateral branches, spherical to sub-spherical, smooth walled, 12.5—33  $\mu\text{m}$  in diameter, mostly 22.5  $\mu\text{m}$ ; a n t h e r i d i a single, hypogynous, antheridial cells small, 3—8.5  $\mu\text{m}$  wide; o o s p o r e single, spherical, eccentric, plerotic, 11—32  $\mu\text{m}$  in diameter, mostly 21  $\mu\text{m}$ .

Isolated from the diseased roots of *Triticum aestivum* L. and soil of Khatima. The culture has been deposited in the herbarium CMI (IMI-277403), Kew, England. This isolate shows almost complete agreement with the description given by MIDDLETON (1943).

10. *Pythium inflatum* MATTHEWS in Univ. N. Carol. Press. 45: 136, 1931. — Pl. 1, fig. 17, 18

Mycelium well developed on SCHMITTHENNER's agar; s p o r a n g i a inflated or toruloid, complex structure, 8—21  $\mu\text{m}$  in diameter; z o o s p o r e s not observed in culture; sex organs produced in SCHMITTHENNER's agar but not developed in water culture; o o g o n i a terminal or intercalary, spherical, 18—24  $\mu\text{m}$  in diameter; a n t h e r i d i a rarely formed, 1—2 per oogonium when present, declinous; o o s p o r e single, spherical, eccentric, plerotic, 14—22  $\mu\text{m}$  in diameter.

Isolated from diseased roots of tomato and soil of Khatima and Nagartarai.

This species is very close to the description given by ROBERTSON (1980).


11. *Pythium middletonii* SPARROW in Aquatic Phycomycetes (Univ. of Michigan Press), p. 1187, 1960. — Pl. 2, fig. 4

Hyp h a e slender, frequently branched, 2.5—9.5  $\mu\text{m}$  in diameter, predominantly 3.3—5  $\mu\text{m}$ ; z o o s p o r a n g i a abundant, papillated, terminal on lateral branches, spherical subspherical or sometimes ellipsoidal, 19—40  $\mu\text{m}$  in diameter, predominantly 25—36  $\mu\text{m}$ , renewed by internal proliferation; encysted primary z o o s p o r e s 8—11  $\mu\text{m}$  in diameter; c h l a m y d o s p o r e s are of same shape and size like that of sporangia, act as sporangia; o o g o n i a formed abundantly, in some isolates it is delayed, spherical 13.5—32  $\mu\text{m}$  in diameter, predominantly 18.5—25  $\mu\text{m}$ ; a n t h e r i d i a androgynous, declinous and hypogynous, predominantly declinous, 1—3 per oogonium; o o s p o r e s 1—2, mostly one, eccentric, aplerotic, 9.5—22  $\mu\text{m}$  in diameter.

Isolated from Ram Tal water and soil.

The first report of this species from India was by RAJGOPALAN & RAMKRISHNAN 1967—68). Recently, MANOHARACHARY & RAO (1978) reported this species from pond waters and soil of Vikarabad (Hyderabad). But it is being reported from the temparate region of Kumaon for the first time. During the present study of Pythiaceous fungi, this

species was found dominant in all the collections. The present isolate differs from the description given by RAJGOPALAN & RAMKRISHNAN (1967—68) in having mostly androgynous and hypogynous antheridia in addition to diclinous antheridia and also in the size slightly larger as oogonia.



Pl. 2: Fig. 1, 2 (scale D): *Pythium artotrogus* (MONT.) de BARY. — Oogonia with hypogynous antheridia.  
 3 (scale D): *P. hypogynum* MIDDLETON. — Mature oogonium with plerotic oospore and hypogynous antheridium.  
 4 (scale B): *P. middletonii* SPARROW. — Mature oogonium with aplerotic oospore and diclinous antheridia.  
 5 (scale D): *P. debaryanum* HESSE. — Mature oogonium with aplerotic oospore and androgynous and diclinous antheridia.  
 6 (scale A) and 7, 8 (scale B): *P. vexans* var. *minutum* MER & KHULBE. — 6. Mature oogonium with hypogynous antheridium. — 7. Oogonium with hypogynous and diclinous antheridia. — 8. Oogonium with diclinous antheridia.  
 9, 10 (scale C): *P. proliferum* de BARY. — 9. Sporangium showing internal proliferation. — 10. Oogonium with diclinous antheridium.  
 11, 12 (scale D): *P. parocandrum* DRECHSLER. — 11. Intercalary sporangium. — 12. Intercalary oogonium with androgynous antheridia.



12. *Pythium monospermum* PRINGSHEIM in Jahrb. Win. Bot. 1: 284—306, 1868. — Pl. 1, fig. 7

Hyphae long, slender, frequently branched, 2.5—5  $\mu\text{m}$  in diameter; sporangia terminal or lateral, filamentous, branched or unbranched, 25—121  $\mu\text{m}$  long by 2.5—5.5  $\mu\text{m}$  in diameter, wavy in outline; zoospores 4—30, reniform, encysted primary zoospores 9—11.5  $\mu\text{m}$  in diameter, predominantly 11  $\mu\text{m}$ ; oogonia terminal, sometimes intercalary, spherical, 13.2—26.4  $\mu\text{m}$  in diameter, mostly 23  $\mu\text{m}$ ; oogonial wall smooth, thin; antheridial branches declinous, sometimes monoclinal, 1—2 per oogonium, clavate; oospore plerotic, single, occasionally two, eccentric, spherical, 11—16.5  $\mu\text{m}$  in diameter, predominantly 13  $\mu\text{m}$ .

Isolated on apple and tomato fruits and hempseed from Khurpatal water.

13. *Pythium paroecandrum* DRECHSLER in J. Wash. Acad. Sci. 20: 407—407, 1930. — Pl. 2, fig. 11, 12

Colony well developed on hempseed; hyphae measuring 3—8  $\mu\text{m}$  in diameter; sporangia spherical to ellipsoidal, typically intercalary, occasionally terminal on short to long branches, spherical sporangia 15—30  $\mu\text{m}$  in diameter, mostly 23  $\mu\text{m}$ , ellipsoidal sporangia 18—36  $\mu\text{m}$  long and 14—24  $\mu\text{m}$  in diameter; zoospores reniform, encysted, primary zoospores 9—11  $\mu\text{m}$  in diameter; oogonia usually intercalary, smooth, sub-spherical, 11—28  $\mu\text{m}$  in diameter, mostly 22  $\mu\text{m}$ ; antheridia 1—3 per oogonium, androgynous or declinous; oospore single, aplerotic, smooth walled, spherical 10—22  $\mu\text{m}$  in diameter, mostly 18  $\mu\text{m}$ .

Isolated from diseased roots of cabbage and cauliflower and soil of Nagaratarai.

In India this species has been reported by BALKRISHNAN (1948) and RAO (1963).

14. *Pythium proliferum* de BARY in Jahr. Wiss. Bot. 2: 169—192, 1860. — Pl. 2, fig. 9, 10

Hyphae slender, branched, 2.5—4.5  $\mu\text{m}$  in diameter; sporangia terminal, spherical or ovoid, papillate, 25—55  $\mu\text{m}$  in diameter, renewed internally and laterally; vesicle sessile, some with short stalk; zoospores reniform; encysted zoospores 8—11.5  $\mu\text{m}$  in diameter; oogonia terminal or lateral, sometimes intercalary, spherical, 15.5—35  $\mu\text{m}$  in diameter; oogonial wall smooth, thin; antheridia usually monoclinal, rarely declinous, 1—3 per oogonium, clavate, tuberous; oospore single, aplerotic, 13—21  $\mu\text{m}$  in diameter, predominantly 19.6  $\mu\text{m}$ , eccentric.

Isolated on apple fruit and hempseed from Bhimtal and Naukuchital lakes.

15. *Pythium pulchrum* MINDEN in Mykol. Unters. Ber. 1: 227—228, 1923. — Pl. 1, fig. 9, 10

Main hyphae long, branched, 4.5—7  $\mu\text{m}$  in diameter; zoosporangia spherical to sub-spherical, terminal or intercalary, 20—33  $\mu\text{m}$  in diameter, predominantly 24  $\mu\text{m}$ ; zoospores reniform, encysted zoospores 9—11  $\mu\text{m}$  in diameter; oogonia spherical to sub-spherical, terminal or intercalary,

single or catenulate, 1—4 in chains, 18—25  $\mu\text{m}$  in diameter, mostly 22  $\mu\text{m}$ ; antheridia mostly hypogynous, barrel shaped; oospores spherical, 1—2  $\mu\text{m}$  in number, mostly single, eccentric, aplerotic, 12.5—18  $\mu\text{m}$  in diameter, predominantly 15  $\mu\text{m}$ .

Isolated from infected seedlings of *Hibiscus esculentus* L. near Nal Damayanti Tal, Naini Tal.

This species had mostly hypogynous antheridial branches which are similar to the description given by MIDDLETON (1943), who reported that some strains of this species produce only hypogynous antheridia. JOHNSON (1971) also found variation in the origin of antheridial branches in this species. This species is a new record for Indian aquatic fungi.

16. *Pythium rostratum* BUTLER in Mem. Dep. Agric. India Bot. Ser. 1 (5): 84—85, 1907. — Pl. 1, fig. 16

Hyphae non septate, branched, 6—8  $\mu\text{m}$  in diameter; colony well developed on hempseeds; sporangia terminal or intercalary, spherical to oval, 22 to 32.5  $\mu\text{m}$  in diameter, mostly 24  $\mu\text{m}$ ; zoospores produced within the vesicle, reniform; encysted zoospores 7—9  $\mu\text{m}$  in diameter; oogonia typically intercalary or lateral, 15—24  $\mu\text{m}$ , mostly 21  $\mu\text{m}$  in diameter, smooth walled; antheridia generally one per oogonium, monoclinal, often hypogynous; oospore single, eccentric, plerotic, 14—23  $\mu\text{m}$  in diameter.

Isolated from roots of wheat, rice, cabbage, cauliflower, tomato and soil of Khatima. The culture has been deposited in the herbarium CMI (IMI-277406), Kew England.

This isolate shows complete agreement with the description given by MIDDLETON (1943) and ROBERTSON (1980).

17. *Pythium spinosum* SAWADA ap. SAWADA & CHEN in Trans. Nat. Hist. Soc. Formosa 16: 199—200, 1926. — Pl. 1, fig. 14, 15

Hyphae branched, 3—5  $\mu\text{m}$  in diameter; sporangia abundant, spherical to sub-spherical, 12—27  $\mu\text{m}$  in diameter, mostly 22  $\mu\text{m}$  in diameter, germinating by germ tubes; oogonia spherical to sub-spherical, generally terminal, occasionally intercalary, 13—25  $\mu\text{m}$  in diameter, mostly 19  $\mu\text{m}$ ; oogonial wall echinulate, spines straight or curved, 4—8  $\mu\text{m}$  in length; antheridia 1—2 per oogonium, monoclinal, sometimes declinal; oospore single, plerotic, 12—23  $\mu\text{m}$  in diameter, mostly 17.5  $\mu\text{m}$ .

Isolated from rice field soil of Khatima. The culture has been deposited in the herbarium CMI (IMI 277405) Kew, England.

This species was recorded in India by RAMKRISHNAN (1955), RAO (1963), RAGHUNATHAN (1968) and MANOHARACHARY (1973). The characters of this isolate correspond well with the description given by MIDDLETON (1943).

18. *Pythium torulosum* COKER & PATTERSON in J. Elisha Mitchell Sci. Soc. 42: 247—50, 1927. — Pl. 1, fig. 21

Colony well developed on hempseeds, measuring up to 2.5  $\mu\text{m}$  in diameter within 10 days under laboratory conditions; hyphae 2.5—4  $\mu\text{m}$  in

diameter; sporangia abundant, terminal, tubular complex up to 52  $\mu\text{m}$  in length and 12  $\mu\text{m}$  in diameter; zoospores readily produced in water culture; oogonia mostly terminal rarely intercalary, 12–24  $\mu\text{m}$  in diameter, mostly 18  $\mu\text{m}$ ; oogonial wall smooth; antheridia single, androgynous; oospore single, spherical, smooth-walled, eccentric, plerotic, 10–12  $\mu\text{m}$  in diameter.

Isolated from wheat roots and soil of Khatima and Nanakmatta.

The culture is very close to the description given by MIDDLETON (1943).

19. *Pythium ultimum* TROW in Ann. Bot. 15: 300–301, 1901. — Pl. 1, fig. 11

Hypphae branched, 3–6.5  $\mu\text{m}$  in diameter; sporangia chiefly terminal and spherical, 14–28  $\mu\text{m}$  in diameter, mostly 21  $\mu\text{m}$ ; zoospores not observed; oogonia typically terminal, spherical, 19–26  $\mu\text{m}$  in diameter, mostly 21  $\mu\text{m}$  in diameter; oogonial wall smooth; antheridia single, rarely two per oogonium, typically sessile; oospore single, spherical, smooth-walled, eccentric, aplerotic, 15–22  $\mu\text{m}$  in diameter, mostly 18  $\mu\text{m}$ .

Isolated from wheat roots and soil of Nanakmatta.

In India this species has been reported by BHARGAVA & SINGH (1965) and CHOWDHRY & AGARWAL (1981).

20. *Pythium undulatum* PETERSON in Bot. Tidskr. 29: 394, 1909. — Pl. 1, fig. 4, 5

(Reprinted with English text in Ann. Mycol. 8: 531, 1910).

Hypphae undulating, long, measuring 3–6  $\mu\text{m}$  in diameter; sporangia ellipsoidal, acrogenous, often with a small apical papilla, 80–150  $\mu\text{m}$  long and 35–60  $\mu\text{m}$  in diameter; zoospore readily produced in water cultures, encysted zoospores 9–12  $\mu\text{m}$  in diameter; sex organs not developed.

Isolated from rice roots and soil of Khatima and Nanakmatta.

In India, this species has been observed by DAYAL & TANDON (1962), SRIVASTAVA (1964) and KHULBE (1979).

21. *Pythium vexans* de BARY var. *minutum* MER & KHULBE in Current Science 52 (15): 735–736, 1983. — Pl. 2, fig. 6, 8

Hypphae slender, frequently branched, 2–9  $\mu\text{m}$  in diameter; zoosporangia formed abundantly, terminal or intercalary, spherical, pyriform, ovoid or sub-spherical, 12.5–25  $\mu\text{m}$  in diameter, predominantly 16–18  $\mu\text{m}$ , germinated by 1–2 germ tubes; zoospores not observed; chlamydospores are of same size and shape like sporangia when present, germinate by germ tubes; oogonia abundant, terminal on short lateral branches, occasionally intercalary, spherical, smooth walled, 13–31  $\mu\text{m}$  in diameter, predominantly 15–18  $\mu\text{m}$ ; antheridia mostly hypogynous, but androgynous and declinuous antheridial branches are also not uncommon, bell-shaped; oospore single, smooth-walled, spherical, eccentric, mostly aplerotic, rarely plerotic, 10–16  $\mu\text{m}$  in diameter, mostly 11  $\mu\text{m}$ .

Isolated from Ram Tal soil. The culture has been deposited in the herbarium CMI (IMI No. 255017), Kew, England.

The first report of this variety was by MER and KHULBE (1983).

### Acknowledgements

This work is supported financially by CSIR, New Delhi. The authors are much grateful to Prof. K. S. BHARGAVA, Director, College Development Council, University of Gorakhpur for his valuable suggestions and to Professor B. S. MEHROTRA, Head, Department of Botany, Kumaun University, Nainital, for providing necessary laboratory facilities.

### References

- AULAKH, K. S. (1971). Damping off of toria seedlings due to *Pythium butleri*. — Indian Phytopathology 24: 611—612.
- BALKRISHNAN, M. S. (1948 a). South Indian Phycomycetes. I. *Pythium indicum* sp. nov. causing fruit-rot of *Hibiscus esculentus* LINN. — Proc. Ind. Acad. Sci. 27B: 161—173.
- (1948 b). South Indian Phycomycetes. II. Some little known species of *Pythium* occurring in South India. — Proc. Ind. Acad. Sci. 28 B: 27—34.
- BHARGAVA, K. S. (1941). *Pythium aphanidermatum* (EDS.) FITZ. on *Carica papaya*. — Curr. Sci. 10: 212—213.
- & GUPTA, S. C. (1957). Market diseases of fruit and vegetable in Kumaun. I. Cottony Leak of beans. — Ind. Phytopathology 10: 48.
- & SINGH, B. B. (1965). Observations on Indian aquatic fungi. III. Some species of Blastocladales, Monoblepharidales and Peronosporales collected from Gorakhpur. — Proc. Nat. Acad. Sci. India 35 (1) B: 49—58.
- BLATTER, E. (1911). A list of Indian fungi chiefly of the Bombay presidency with the description of two new species. — J. Bombay Nat. Hist. Soc. 21: 146—152.
- BUTLER, E. J. (1907). An account of the genus *Pythium* and some Chytridiaceae. — Mem. Dept. Agr., India, Bot. Ser. 1: 1—160.
- CHOWDHRY, P. N. & AGARWAL, G. P. (1981). Taxonomic studies on aquatic fungi of India. I. Pythiaceae. — Indian Phytopathology 34 (2): 235—236.
- DASTUR, J. F. (1935). Diseases of Pan (*Piper betle* L.). — Proc. Ind. Acad. Sci. India 28: 49—57.
- DAYAL, R. & TANDON, R. N. (1962). Ecological studies of some aquatic Phycomycetes. — Hydrobiologia 20: 121—127.
- DESHPANDE, K. B. & DESHPANDE, K. S. (1964). A new species of *Pythium*. — Sydowia 16: 33—34.
- DIWAKAR, M. C. & PAYAK, M. M. (1975). *Pythium* stalk rot — a deadly disease of maize in India. — Curr. Sci. 44 (6): 209.
- GATTANI, M. L. & KAUL, T. N. (1951). Damping off of tomato seedlings — its cause and control. — Indian Phytopathology 4: 156—161.
- JAIN, A. C. (1952). Control of *Pythium* fruit-rot of cucurbits in Madhya Pradesh. — Nagpur Agric. Coll. Mag. 26: 1—5.
- JOHNSON, T. W. (1971). Aquatic fungi of Iceland. *Pythium*. — Mycologia 63: 517—536.
- KHULBE, R. D. (1979). *Pythium* in some lakes of Naini Tal. — Geobios 6: 335.
- (1983). *Pythium elongatum* MATTHEWS. A new record for temperate Indian aquatic fungi. — Current Science 52 (6): 260.

- MAHMUD, K. A. (1950). Damping off of cabbage, cauliflower and knowl-kohl caused by *Pythium aphanidermatum* (EDSON) FITZ. — *Curr. Sci.* 19: 67—68.
- (1951). Damping off of cotton seedlings by *Pythium debaryanum*. — *Sci. and Cult.* 16: 423—424.
- (1952). *Pythium* — damping off of Brinjal seedlings. — *Sci. and Cult.* 18: 149—150.
- MANOHARACHARY, C. H. (1981). The taxonomy and ecology of fresh water Phycomycetes from India. — *Nova Hedwigia* 63: 187—196.
- & RAO, P. R. (1978). Distribution and seasonal variation of some *Pythium* species in some soils, pond water and mud soils of Hyderabad district. — *Mysore J. Agri. Sci.* 12: 280—284.
- MER, G. S. & KHULBE, R. D. (1983). *Pythium vexans* var. *minuta* var. nov. from Kumaun, the Himalayas, India. — *Current Science* 52 (15): 735—736.
- MIDDLETON, J. T. (1943). The taxonomy, host range and geographic distribution of the genus *Pythium*. — *Mem. Torrey Bot. Club* 20: 1—171.
- MITRA, M. (1927). Root rot disease of cultivated Cucurbitaceae by *Pythium aphanidermatum* (EDS.) FITZ. — *Proc. 14th Indian Sci. Congr. Bot.* 5: 213—214.
- MUTHUSAMY, M., VIDHYASEKARAN, P. & RAJGOPALAN, C. K. S. (1974). Physiology of damped off tomato seedlings (*Pythium aphanidermatum*). — *Madras Agri. Jour.* 61: 27—30.
- PRABHUJI, S. K. & SRIVASTAVA, G. C. (1976). A new record of *Pythium* from India. — *Curr. Sci.* 45: 708.
- PRAKASH, R. & SAKSENA, R. K. (1952). Decomposition of paddy and Bajara (*Pennisetum typhoideum*) straws by fungi commonly found in Allahabad soils. — *Proc. Ind. Acad. Sci.* 36 B 119—128.
- RAGHUNATHAN, V. (1968). Damping off of green gram, cauliflower, Daincha, Raghi and cluster bean. — *Indian Phytopathology* 21: 456—457.
- RAJGOPALAN, C. K. S. & RAMKRISHNAN, K. (1967—68). Phycomycetes in agricultural soils with special reference to Pythiaceae. II. Species of *Pythium* isolated from Madras. — *Madras Univ. J.* 37—38: 100—117.
- RAMKRISHNA Iyer, T. S. (1929). *Pythium aphanidermatum* (EDS.) FITZ. on *Opuntia dilleni* HAW. — *Mem. Dept. Agri. India. Bot.* 16: 191—201.
- RAMKRISHNAN, K. (1955). Some aspect of soil fungal ecology. — *Proc. Ind. Acad. Sci.* 41B: 110—116.
- RAMKRISHNAN, L., THAMBURAJ, S., KAMALNATHAN, S. & KRISHNAMURTHY, C. S. (1973). Studies on the control of damping off (*Pythium aphanidermatum* (EDS.) FITZ.) disease in certain solanaceous vegetable crops. — *Madras Agri. Jour.* 60 (7): 537—538.
- RAMKRISHNAN, T. S. (1943). The occurrence of *Pythium vexans* de BARY in South India. — *Indian Phytopathology* 2: 27—30.
- RAO, V. G. (1963). An account of the genus *Pythium* PRINGSHEIM in India. — *Mycopathologia et Mycologia Applicata* 21 (1): 45—59.
- ROBERTSON, G. I. (1980). The genus *Pythium* in New Zealand. — *New Zealand Journal of Botany* 18: 73—102.
- SHARMA, O. P. & ASTANA, R. P. (1958). Fruit-rot disease of *Momordica dioica* ROXB. due to *Pythium aphanidermatum* (EDS.) FITZ. — *Sci. and Cult.* 23 (8): 435.
- SHARMA, O. P. & NEMA, K. G. (1952). A short note on *Pythium aphanidermatum* (EDS.) FITZ. on *Linum grandiflorum* LINN. — *Mag. Agric. Coll. Nagpur* 26: 27.
- SRINIVASAN, K. U. (1956). *Pythium catenulastrum* MATTHEWS — Sugarcane seedling root-rot. — *Curr. Sci.* 299—300.

- SRIVASTAVA, G. C. (1964). Studies on some lower fungi of Gorakhpur. — Ph. D. Thesis, University of Gorakhpur, Gorakhpur.
- SUBRAMANIAM, L. S. (1919). A *Pythium* disease of ginger, tobacco and papaya. — Mem. Dept. Agric. India. Bot. Ser., 10 (4) : 181—194.
- SYDOW, H. & BUTLER, E. J. (1907). Fungi India Orientalis. Pars II. — Annales Mycologici 5: 485—515.
- WATERHOUSE, G. M. (1967). Key to *Pythium* PRINGSHEIM. — Mycological Papers, Commonwealth Mycological Institute 109: 15 p.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1984

Band/Volume: [37](#)

Autor(en)/Author(s): Mer G. S., Verma B. L., Khulbe R. D.

Artikel/Article: [Aquatic Fungi of Kumaun Himalaya, Indien: \*Phythium PRINGSHEIM.208\* 208-221](#)