

The Polyol Pattern, Chemotaxonomy, and Phylogeny of the Fungi *)

PFYFFER, Gaby E., PFYFFER, B. U. **) & RAST, Dora M.

Department of Plant Biology, University of Zürich,
CH-8008 Zürich, Switzerland, and

**) Department of Forestry, Canton of Lucerne, CH-6002 Lucerne, Switzerland

Abstract. – Acyclic polyols are widely distributed within the fungi and also represent the fungal metabolites studied most intensively. Therefore, the taxonomic evaluation of the alditol pattern of fungi suggests itself as a tool for a model testing of existing classifications at higher levels of hierarchy. To this aim, all data on the systematic distribution of polyols within the Eumycetes (nearly 600 records, representing some 400 species) have been processed with the database facility of the integrated software package Lotus® Symphony 1.1 installed on an IBM® PC XT. When delimiting groups on account of the polyol character (3 states: P₀, polyols absent; P₁, polyols, except mannitol, present; P₂, mannitol [and other polyols] present), the taxa thus formed coincide with the Oomycetes (P₀), the Zygo- and Hemiascomycetes (P₁), and the Chytridio-, Euasco-, Basidio-, and Deuteromycetes group (P₂), as defined by classical criteria. Not unexpectedly, the Blastomycetes are heterogeneous with respect to the polyol pattern. As judged from the very few apparent exceptions (5), the polyol character appears to be extremely conservative and, therefore, lends itself as a marker for the assignment of species of doubtful systematic position to conventional taxa, as well as for the unravelling of evolutionary relationships within the fungi. These patterns, together with other biochemical traits, are integrated into a phylogenetic scheme, using the classification system of AINSWORTH & al. (1971) as the reference.

Introduction

Traditionally, the taxonomy, systematics, and phylogeny of the fungi have been – and mostly still are – based on morphological traits, especially those displayed during reproductive behaviour (cf. [198]). In the continuous quest for an information retrieval system suitable for identification as well as for unravelling the evolution of the fungi, the use of such characters is not only firmly entrenched but also has been vigorously defended (cf. [179]). Nevertheless, this approach has its limitations, especially in cases where distinctive morphological features are scarce or lacking [83].

To compensate for these insufficiencies, various attempts have been made to determine molecular characteristics of fungi and, thence, to introduce chemical differential characters into fungal taxonomy [201, 9, 208, 216, 172, 173, 186, 187, 102, 171, 18, 19, 20,

*) Dedicated to Prof. Dr. Hans WANNER (Zürich, Switzerland) on the occasion of his 70th birthday.

11]. However, many of these efforts have proved unsuccessful or, at the best, unsatisfactory, especially as systematics at higher levels of hierarchy is concerned. Thus, many characteristics studied have not displayed sufficient taxonomic 'resolving power', as e. g. the patterns of fatty acids [172, 173] and carotenoids [118, 11]. For others, the compounds concerned were found to be restricted to groupings of low taxonomic rank only, such as the confinement of psilocybin to *Psilocybe* and *Conocybe* species [207, 225], and of certain complex toxic cyclopeptides to *Amanita* [207, 225] and some *Galerina* [209] and *Lepiota* (cf. [34]) representatives. Occasionally, deductions have been made from studies encompassing merely a few species (e. g. the distribution of some non-protein amino acids in a few agarics [90]), or the chemical characteristics investigated could intrinsically not be expected to yield two-state criteria (e. g. the (G + C) percentage of DNA [187, 188], although a large number of species (ca. 500) was studied in this case).

The notable exceptions to this unsatisfactory situation concern differences in intermediates of the lysine biosynthetic pathway [216], in the sedimentation pattern of enzymes involved in tryptophan biogenesis [102], and in cell wall composition [9, 18, 19]. Thus, on the one hand, the Oomycetes are discrete by any of these three criteria and, on the other, the Zygomycetes by their unique combination of type of organization of the tryptophan pathway and of cell wall construction. The Chytridio-, Euasco-, Homobasidio-, and Deuteromycetes, finally, constitute one single group, inasmuch as their representatives display identical physico-chemical properties of the enzymes involved in tryptophan synthesis as well as a high degree of similarity in cell wall composition, with chitin as the major microfibrillar component. The coincidence of chemotaxonomic and conventional groupings – though data are based on a limited number of species (some 25, 20, and 90, respectively) – is indicative of the highly conservative nature of these traits and, thus, also of their highly predictive systematic value (cf. [93]).

To test further whether the employment of chemical data for the construction of general purpose as well as evolutionary classifications of the fungi, is, indeed, a feasible aim, a consideration of the distribution of the acyclic sugar alcohols (alditols: cf. [35]) suggests itself for a model investigation, because polyols in the free form are the fungal components studied most widely (see Results). The reasons for this are twofold: (1) as polyols crystallize easily from alcoholic solutions, their existence in fungi has been known for nearly two centuries [32, 111], and over this period a considerable amount of information on their distribution has been accumulated (see [231, 123, 156, 105]), particularly as – in contrast to earlier times – polyols are now assessed easily by gas-liquid chromatography [156],

or by thin-layer chromatography using spray reagents which discriminate them from their corresponding monoses [87, 156, 224]; and (2) polyols have physiological functions, of which the storage of carbon and reducing power [123], involvement in the control of growth [61, 62] and of water potential [106], regulation of cytoplasmic pH value [105] and – where applicable – the maintenance of the proper chemical potential gradient for carbon movement in host-parasite associations [176] are presently the most favoured experimental hypotheses.

The aim of the present investigation was to compile all data on the identity and systematic distribution of acyclic sugar alcohols in the fungi and to delimit phenetic groups based on the polyol pattern of their component members, in order to possibly add another “objective criterion” (cf. [50, 16]) for classifying the fungi at higher taxonomic levels and to, thus, finally also gain further insight into their phylogeny.

Compilation, Systematic Evaluation, and Taxonomic Application of Data

This paper presents a comprehensive enumeration of the fungi that have been studied hitherto with respect to the occurrence of acyclic polyols. The species have been arranged either in alphabetical order (Tables 1–5), or according to the systematic scheme of AINSWORTH & al. ([6]; Table 6). *Binomials are as stated by the authors themselves, i. e. any taxonomic or nomenclatural changes which might subsequently have occurred have been disregarded.* Hence, synonymy has not been accounted for. The lists do not include those studies of fungi not identified by their specific epithets, nor do they encompass works in which the presence of a particular sugar alcohol has not conclusively been established but simply been inferred from enzymic tests or from the similarity observed in the mobility of a compound with that of an authentic polyol using merely one single chromatographic method with limited sensitivity. Furthermore, reference is not made to papers stating the absence of a particular polyol in a given species if this has later been found to contain the compound in question. Finally, in view of a taxonomic evaluation of the data (see below), reports based on analyses of mixed fungus/plant or fungal/algal material have also been omitted, since the identity of the polyol-synthesizing partner of the associations remained uncertain. Reference to studies of this type that relate to phytopathogenic symbioses (*sensu* [46]) is made separately in Table 7. The possibility that fungal infection may elicit polyol formation in a green plant is not a far-fetched idea, because: (i) some vascular plants do contain polyols (mainly sorbitol or mannitol: see

[122]), and (ii) the green algal symbionts within lichens apparently synthesize a polyol but cease to do so upon removal of the fungus (see [46, 168]). On the other hand, works that explicitly state the absence of polyols even in phytopathogenic associations of fungi are included in Table 1. The computation was carried out on an IBM® model XT Personal Computer; the database was established with Lotus® Symphony 1.1 (Lotus Development Corp., Cambridge, MA, USA).

The present study comprises nearly 600 records relating to approximately 400 different species representing some 1.3% of the lower and 1.0% of the higher fungi, respectively (based on the estimated total number of species given in [6]). To date, the occurrence of at least one polyol has been reported in some 390 species (Table 2), whereas the absence of sugar alcohols has been established in 7 species (Table 1) – all Oomycetes (cf. Table 6). The other lower fungi are heterogeneous with regard to their polyol pattern: beside of glycerol, ribitol is present in Zygomycetes, and mannitol in Chytridiomycetes. Throughout the higher fungi – except the Hemiascomycetes – mannitol is being accumulated. The most versatile organisms with respect to polyol synthesis appear to be clustered within the Ascomycetes-Heterobasidiomycetes-Deuteromycetes group, inasmuch as glycerol, erythritol, threitol, ribitol, arabitol, xylitol, dulcitol, and sorbitol may be present, e. g. in *Claviceps purpurea*, *Puccinia graminis tritici*, and *Pyrenophaeta terrestris* (Table 6). The polyol pattern of the Homobasidiomycetes, finally, consists almost exclusively of mannitol, as other polyols have only occasionally been found.

Groups of species (A-E: Tables 1–5) have been made according to the following criteria: (i) absence of polyols, (ii) presence of polyols, (iii) presence of polyols, except mannitol, (iv) presence of mannitol (and other polyols), and (v) presence of uncommon polyols. As judged by classical criteria (see Introduction) group A consists entirely of representatives of the Oomycetes (see also above), whereas all species of group B belong to the Chytridio-, Asco-, Basidio-, or the Deuteromycetes. Group B is subdivided into the groups C and D by the criteria (iii) and (iv) as detailed in Tables 3 and 4. Group D encompasses only members of the higher fungi and Chytridiomycetes, group C Zygomycetes and, for the rest, mostly yeasts (Hemiascomycetes as well as Blastomycetes). The appearance, in group C, of a representative of the genus *Torula*, i. e. of a hyphomycete, is almost certainly due to a taxonomic mis-statement as to the fungus actually analyzed, which appears to have been *Candida utilis*, not *Torula utilis*, as judged from the description of the experimental organism by the authors themselves [71] as well as from the notions of BARRETT & al. [19] and AHEARN [5] that *C. utilis* is often wrongly referred to as

Table 1: Chemotaxonomy of fungi: group of species containing no polyols in the water-soluble fraction (group A)

<i>Achlya radiosua</i> 156	<i>Phytophthora cinnamomi</i> 156
<i>Albugo tragopogonis</i> 126	<i>Phytophthora infestans</i> 30
<i>Peronospora parasitica</i> 202	<i>Plasmopara viticola</i> 30, 33
<i>Phytophthora cactorum</i> 27	

Torula yeast. Since all Homobasidiomycetes species analyzed and also the Hyphomycetes do synthesize mannitol (see Table 6), it can, furthermore, safely be assumed that the three *Penicillia* listed in group C have simply not been thoroughly searched for the presence of the hexitol and it, hence, appears justified to allocate them in the neighbourhood of group D. This may also hold for some Blastomycetes listed in group C. In comprising either Hemiascomycetes (most of which contain arabitol) or Zygomycetes (in which the occurrence of ribitol is common: see above), the remainder of group C consists of two well-defined entities. That the group of species synthesizing a rare polyol (group E: Table 5), *viz* threitol, ribitol, xylitol, dulcitol, sorbitol, volemitol, *meso-glycero-ido-heptitol*, or *D-glycero-D-ido-heptitol*, would, as a whole, be heterogeneous with respect to the position in conventional systematics of the organism concerned was to be expected, since these substances are in themselves structurally and, conceivably, also biosynthetically (see [105]) quite different, and since gain of the ability to elaborate a given rare secondary compound by singular members of phylogenetically even very distant lines is a fairly common phenomenon (see [66, 89, 206, 34]). Also, compounds which appear erratically are generally suitable for systematic purposes only at low taxonomic level (species, or below: cf. [91]) and not for the study of taxa of higher ranks as aimed at in the present investigation (see Introduction). Nevertheless, despite of its conceptionally artificial character, group E has been established and its members presented in Table 5 – for the sake of completeness of this compilation.

When now delimiting chemosystematic taxa on account of the characters P_0 (no polyols present), P_1 (a polyol, except mannitol, present), and P_2 (mannitol present), the three groups obtained coincide with (1) the Oomycetes, (2) the Zyg- and Hemiascomycetes, and (3) the Chytridio-, Euasco-, Basidio-, and Deuteromycetes group (except some Imperfect Yeasts), as shown by Table 6. Heterogeneity of the Blastomycetes as to the polyol pattern cannot be a surprise. These and the (apparent) exceptions, i. e. the absence of mannitol in "*T. utilis*" and some *Penicillia* (see also above), as well as the P_2 -pattern of *Coccidioides immitis* and of *Byssochlamys* species (see Tables 4 and 6) are dealt with in the Discussion.

Table 2: Chemotaxonomy of fungi: group of species containing free polyol(s) (group B). * References are not given separately to the original publications antedating this review article.

<i>Acanthocystis petalooides</i>	72	<i>Aspergillus versicolor</i>	85
<i>Acetabula vulgaris</i>	231 *	<i>Aspergillus wentii</i>	24
<i>Actinomucor elegans</i>	156, 157	<i>Blastocladiella emersonii</i>	156
<i>Agaricus arvensis</i>	28	<i>Boletus aereus</i>	28
<i>Agaricus bisporus</i>	166, 60, 81, 59, 61, 182, 84, 155	<i>Boletus appendiculatus</i>	231 *
<i>Agaricus campester</i>	72, 163	<i>Boletus aurantiacus</i>	231 *
<i>Agaricus campestris</i>	134, 63	<i>Boletus badius</i>	231 *
<i>Agaricus eryngii</i>	231 *	<i>Boletus bovinus</i>	231 *, 219, 103, 74
<i>Agaricus silvaticus</i>	28	<i>Boletus calopus</i>	231 *
<i>Agaricus silvicola</i>	72	<i>Boletus chrysenteron</i>	231 *
<i>Agaricus xanthoderma</i>	72	<i>Boletus cyanescens</i>	231 *
<i>Allomyces arbuscula</i>	156	<i>Boletus edulis</i>	231 *, 103, 28, 166
<i>Alternaria alternata</i>	101	<i>Boletus erythropus</i>	231 *, 155
<i>Alternaria tenuissima</i>	156	<i>Boletus luridus</i>	231 *, 28
<i>Amanita aspera</i>	231 *	<i>Boletus luteus</i>	231 *, 103
<i>Amanita bulbosa</i>	231 *	<i>Boletus pachypus</i>	231 *
<i>Amanita caesarea</i>	231 *, 28	<i>Boletus pruinatus</i>	231 *
<i>Amanita citrina</i>	72, 28	<i>Boletus rufus</i>	72, 28
<i>Amanita muscaria</i>	231 *, 72, 28	<i>Boletus satanas</i>	28
<i>Amanita pantherina</i>	231 *, 28	<i>Boletus scaber</i>	231 *, 28
<i>Amanita phalloides</i>	72, 28	<i>Boletus subtomentosus</i>	231 *, 103
<i>Amanita porphyria</i>	28	<i>Boletus variegatus</i>	231 *
<i>Amanita rubescens</i>	231 *, 28	<i>Botrytis cinerea</i>	117, 154, 30
<i>Amanita vaginata</i>	231 *	<i>Bulgaria inquinans</i>	231 *, 72
<i>Amanitopsis vaginata</i>	28	<i>Byssochlamys fulva</i>	165
<i>Armillaria caligata</i>	103	<i>Byssochlamys nivea</i>	156
<i>Armillaria mellea</i>	231 *, 25, 155, 130, 79	<i>Candida albicans</i>	22, 158
<i>Armillariella mellea</i>	72	<i>Candida arborea</i>	148
<i>Ascobolus stercorarius</i>	156	<i>Candida diddensii</i>	78
<i>Aspergillus candidus</i>	175, 190, 143	<i>Candida guilliermondii</i>	150, 196, 170, 68
<i>Aspergillus clavatus</i>	98	<i>Candida lipolytica</i>	197
<i>Aspergillus elegans</i>	23, 80	<i>Candida parapsilosis</i>	22
<i>Aspergillus fischeri</i>	162	<i>Candida pelliculosa</i>	109
<i>Aspergillus glaucus</i>	228	<i>Candida polymorpha</i>	146, 148, 151, 196
<i>Aspergillus nidulans</i>	23	<i>Candida pseudotropicalis</i>	22
<i>Aspergillus niger</i>	230, 15, 117, 2	<i>Candida tropicalis</i>	146, 22, 78
<i>Aspergillus oryzae</i>	192, 99	<i>Candida utilis</i>	100, 154
<i>Aspergillus terreus</i>	184, 116	<i>Candida zeylanoides</i>	86

- Cantharellus cibarius* 231*, 166
Cantharellus tubaeformis 231*
Chaetomium elatum 156
Chaetomium globosum 3, 154
Circinella mucoroides 51
Circinella muscae 51
Circinella rigida 51
Circinella simplex 51
Circinella umbellata 51
Clavaria aurea 72
Clavaria coralloides 231*
Clavaria corniculata 96
Clavaria flava 231*
Clavaria formosa 231*, 103
Clavaria pistillaris 231*, 28
Claviceps curreyana 47
Claviceps nigricans 47
Claviceps purpurea 231*, 215, 47, 48
Clitocybe aurantiaca 28
Clitocybe cyanophaea 103
Clitocybe geotropa 231*
Clitocybe infundibuliformis 72
Clitocybe nebularis 231*, 72
Clitocybe odora 28
Clitocybe socialis 231*
Coccidioides immitis 125
Cochliobolus miyabeanus 156
Colletotrichum gloeosporioides 117
Collybia butyracea 231*
Collybia confluens 231*
Collybia dryophila 231*
Collybia erythropus 72
Collybia fusipes 231*, 72
Collybia maculata 231*
Collybia platyphylla 28
Coprinus atramentarius 231*, 28, 154
Coprinus comatus 72, 28
Coprinus friesii 156
Coprinus micaceus 72, 28
Cordiceps capitata 199
Coriolus versicolor 28
Cortinarius alboviolaceus 28
Cortinarius berkeleyi 72
Cortinarius bivelus 231*
Cortinarius bolaris 28
Cortinarius brunneus 231*
Cortinarius bulliardii 72
Cortinarius collinitus 231*, 28
Cortinarius elatior 231*
Cortinarius fulmineus 28
Cortinarius infractus 231*
Cortinarius largus 28
Cortinarius torvus 28
Cortinarius violaceus 72, 28
Cortinellus shiitake 103
Coryne sarcoides 96
Craterellus cornucopioides 28
Crepidotus nidulans 72
Cryptococcus neoformans 149
Debaryomyces hansenii 150, 1
Debaryomyces sake 146, 195
Dendryphiella salina 8, 97, 128, 155, 107
Diplodia viticola 191
Elaphomyces asperulus 231*
Elaphomyces echinatus 231*
Elaphomyces granulatus 231*
Elaphomyces leveillei 231*
Elaphomyces variegatus 231*
Endothia parasitica 156
Entoloma clypeatum 28
Entoloma lividum 28
Entoloma nidorosum 231*
Entoloma sinuatum 231*
Epichloe typhina 203
Fistulina hepatica 73, 28
Flammula spectabilis 72
Flammulina velutipes 108
Fomes ignarius 58
Fusarium solani 44
Galactinia olivacea 72
Geaster rufescens 231*
Gelasinospora cerealis 154
Geotrichum candidum 42/1, 183, 52

- Gibberella fujikuroi* 156
Gomphidius roseus 28
Gomphidius viscidus 72
- Hansenula anomala* 146
Hansenula suaveolens 196
Helminthosporium geniculatum 24
Helminthosporium oryzae 30
Helvella esculenta 231 *
Hydnnum erinaceus 72
Hydnnum ferrugineum 231 *, 28
Hydnnum hybridum 231 *
Hydnnum imbricatum 154
Hydnnum repandum 231 *, 28, 166, 154
Hydnnum squamosum 231 *
Hygrophoropsis aurantiaca 28
Hygrophorus agathosmus 231 *
Hygrophorus cossus 231 *
Hygrophorus hypotheius 231 *
Hygrophorus nemoreus 72
Hygrophorus olivaceo-albus 231 *
Hygrophorus virgineus 231 *
Hypholoma fasciculare 231 *, 28
Hypholoma hydrophilum 28
Hypholoma sublateritium 231 *, 103
- Inocybe praetervisa* 28
Ixocomus bovinus 28
Ixocomus luteus 28
Ixocomus piperatus 28
Ixocomus variegatus 28
- Kluyveromyces fragilis* 78
- Laccaria laccata* 28
Lactarius blennius 231 *, 72, 28
Lactarius chrysorrheus 72, 28
Lactarius controversus 231 *, 72
Lactarius deliciosus 231 *, 72, 21, 28, 166
Lactarius musteus 28
Lactarius pallidus 231 *, 72
- Lactarius piperatus* 231 *, 103, 72
Lactarius plumbeus 72, 28
Lactarius pubescens 72
Lactarius pyrogalus 231 *, 72
Lactarius quietus 231 *
Lactarius rufus 231 *, 28
Lactarius subdulcis 231 *
Lactarius torminosus 231 *, 72
Lactarius turpis 231 *
Lactarius uvidus 28
Lactarius vellereus 231 *, 72, 28
Lactarius vietus 231 *
Lactarius violascens 72
Lactarius volvemus 31, 231 *
Lactarius zonarius 231 *, 72
Lentinus cochleatus 231 *, 72
Lentinus edodes 137
Lenzites flaccida 28
Lenzites saepiaria 154
Lepidella vittadini 72
Lepiota clypeolaria 72
Lepiota cristata 72
Lepiota excoriata 231 *
Lepiota friesii 231 *
Lepiota procera 231 *
Lepiota rhacodes 231 *
Leucocoprinus rhacodes 72
Lycoperdon pusillum 231 *
Lycoperdon pyriforme 72
- Marasmius hariolorum* 72
Melanopus squamosus 72
Microsporum gypseum 119
Monascus ruber 156
Morchella conica 231 *
Morchella esculenta 231 *
Morchella semilibera 231 *
Mortierella ramanniana 156, 157
Mucor miehei 156, 157
Mucor mucedo 154
Mucor rouxii 156
Mycena galericulata 96
Mycena pelianthina 231 *
Myrothecium verrucaria 132

- Nematoloma fasciculare* 72
Neurospora crassa 193
Neurospora sitophila 220
Omphalia scyphoides 231 *
Oudemansiella mucida 217
Panus conchatus 231 *
Panus stipticus 231 *
Paxillus atrotomentosus 231 *, 72
Paxillus involutus 231 *, 72
Penicillium brefeldianum 156
Penicillium brevi-compactum 152
Penicillium chrysogenum 24, 12, 2
Penicillium cyclopium 152
Penicillium griseofulvum 117
Penicillium herquei 75
Penicillium italicum 156
Penicillium notatum 29
Penicillium oxalicum 156
Peziza badia 231 *
Peziza nigra 231 *
Peziza ochracea 231 *
Peziza onotica 231 *
Peziza venosa 231 *
Phallus impudicus 231 *
Pholiota aurivella 72
Pholiota lubrica 155
Pholiota mutabilis 103, 28
Pholiota praecox 72
Pholiota radicosa 231 *
Pholiota togularis 231 *
Phycomyces blakesleeanus 156
Phymatotrichum omnivorum 67
Pichia farinosa 146
Pichia guilliermondii 136
Pichia miso 146, 147, 195
Pichia quercibus 194, 196
Pleurotus cornucopiae 72
Pleurotus dryinus 231 *
Pleurotus japonicus 103
Pleurotus serinus 103
Pluteus cervinus 72
Pluteus pellitus 72
Pleurotus olearius 72
Pleurotus ostreatus 103, 72, 154, 156
Polyporus frondosus 72
Polyporus intybaceus 103
Polyporus officinalis 231 *
Polyporus pomaceus 28
Polyporus ptychogaster 231 *
Polyporus robustus 28
Polyporus umbellatus 72
Poria vincta 130
Psalliota arvensis 231 *, 103
Psalliota campestris 231 *, 103, 112
Psalliota placomyces 103
Psalliota silvicola 231 *
Puccinia coronata 200
Puccinia graminis tritici 160, 161, 167, 227, 53, 39, 131, 130, 133
Pyrenophaeta terrestris 226
Pyricularia oryzae 154, 30
Rhizoctonia repens 178
Rhizoctonia solani 178, 156, 30
Rhizopogon luteolus 231 *
Rhizophorus oligosporus 154
Rhizophorus stolonifer 156
Rhodosporidium toruloides 78
Rhodotorula graminis 55
Russula adusta 231 *
Russula albonigra 28
Russula coerulea 28
Russula cyanoxantha 231 *, 28
Russula delica 231 *, 72
Russula drymeia 72
Russula fellea 231 *, 28
Russula foetens 231 *, 28
Russula fragilis 103, 28
Russula integra 231 *
Russula lepida 231 *, 28
Russula nigricans 231 *, 72, 28
Russula ochroleuca 231 *, 28, 96
Russula pseudoviolacea 28
Russula queletii 231 *
Russula sanguinea 72
Russula sardonia 28
Russula torulosa 28
Russula virescens 231 *

- Saccharomyces acidifaciens* 196
Saccharomyces carlsbergensis 78
Saccharomyces cerevisiae 78
Saccharomyces diastaticus 78
Saccharomyces fragilis 146
Saccharomyces mellis 223
Saccharomyces monacensis 146
Saccharomyces paradoxus 146
Saccharomyces rouxii 146, 26, 104, 150, 196, 78
Saccharomyces sake 146, 78
Saccharomyces fibuligera 78
Sarcodon repandum 72
Sarcosphaera coronaria 72
Schizophyllum commune 144, 7, 181, 130
Schizosaccharomyces pombe 78
Scleroderma verrucosum 231*, 72
Scleroderma vulgare 231*, 72
Sclerotinia curreyana 47, 4
Sclerotinia sclerotiorum 120, 221
Sclerotinia tuberosa 231*
Serpula lacrimans 36
Spongipellis spumeus 72
Sporobolomyces roseus 130
Stereostratum corticoides 200
Sterigmatocystis nigra 145, 72
Syncephalastrum racemosum 51

Torula utilis 71
Torulopsis anomala 149
Torulopsis candida 78
Torulopsis famata 146, 196
Torulopsis halophila 196
Torulopsis magnoliae 82
Torulopsis mannitofaciens 151, 196
Torulopsis nodaensis 149, 196
Torulopsis versatilis 146, 149, 196
Tricholoma albo-brunneum 28
Tricholoma album 231*
Tricholoma amarum 28
Tricholoma columbetta 231*, 28
Tricholoma equestre 103, 28
Tricholoma flavobrunneum 231*

Tricholoma nudum 231*, 103
Tricholoma pessundatum 231*
Tricholoma resplendens 231*
Tricholoma russula 231*, 103
Tricholoma sulfureum 231*, 72
Tricholoma terreum 231*
Tricholoma ustale 231*, 103
Trichosporon melibiosaceum 78
Trigonopsis variabilis 151
Tuber cibarium 231*
Tylopilus felleus 72

Ungulina betulina 72
Ustilago esculenta 41
Ustilago maydis 232, 155
Ustilago nuda 77

Verticillium albo-atrum 121
Verticillium dahliae 121, 30
Verticillium fungicola 156
Verticillium intertextum 156
Volvaria speciosa 72
Volvaria volvacea 231*

Xerocomus badius 28, 154
Xerocomus chrysenteron 28
Xylaria polymorpha 231*

Zygorhynchus moelleri 156, 157
Zygosaccharomyces acidifaciens 153
Zygosaccharomyces barkeri 153
Zygosaccharomyces fermentati 153
Zygosaccharomyces mandschuricus 153
Zygosaccharomyces mellis 153
Zygosaccharomyces nadsonii 153
Zygosaccharomyces nussbaumeri 153
Zygosaccharomyces priorianus 153
Zygosaccharomyces richteri 153
Zygosaccharomyces rugosus 153
Zygosaccharomyces wyocena 153

Table 3: Chemosystematics of fungi: polyol-synthesizing organisms not reported to contain free mannitol (group C)

<i>Actinomucor elegans</i>	156, 157	<i>Rhodosporidium toruloides</i>	78
<i>Candida albicans</i>	122, 158	<i>Rhodotorula graminis</i>	55
<i>Candida arborea</i>	148	<i>Saccharomyces acidifaciens</i>	196
<i>Candida diddensii</i>	78	<i>Saccharomyces carlsbergensis</i>	78
<i>Candida guilliermondii</i>	150, 196, 170, 68	<i>Saccharomyces cerevisiae</i>	78
<i>Candida parapsilosis</i>	22	<i>Saccharomyces diastaticus</i>	78
<i>Candida pelliculosa</i>	109	<i>Saccharomyces fragilis</i>	146
<i>Candida polymorpha</i>	146, 148, 151, 196	<i>Saccharomyces mellis</i>	223
<i>Candida pseudotropicalis</i>	22	<i>Saccharomyces monacensis</i>	146
<i>Candida tropicalis</i>	146, 22, 78	<i>Saccharomyces paradoxus</i>	146
<i>Circinella mucoroides</i>	51	<i>Saccharomyces rouxii</i>	146, 26, 104, 150, 196, 78
<i>Circinella muscae</i>	51	<i>Saccharomyces sake</i>	146, 78
<i>Circinella rigida</i>	51	<i>Saccharomycopsis fibuligera</i>	78
<i>Circinella simplex</i>	51	<i>Schizosaccharomyces pombe</i>	78
<i>Circinella umbellata</i>	51	<i>Syncephalastrum racemosum</i>	51
<i>Debaryomyces hansenii</i>	150, 1	<i>Torula utilis</i>	71
<i>Debaryomyces sake</i>	146, 195	<i>Torulopsis candida</i>	78
 		<i>Torulopsis famata</i>	146, 196
<i>Hansenula anomala</i>	146	<i>Torulopsis halophila</i>	196
<i>Hansenula suaveolens</i>	196	<i>Torulopsis magnoliae</i>	82
 		<i>Trichosporon melibiosaceum</i>	78
<i>Kluyveromyces fragilis</i>	78	<i>Trigonopsis variabilis</i>	151
<i>Mortierella ramanniana</i>	156, 157	<i>Zygorhynchus moelleri</i>	156, 157
<i>Mucor miehei</i>	156, 157	<i>Zygosaccharomyces acidifaciens</i>	153
<i>Mucor mucedo</i>	154	<i>Zygosaccharomyces barkeri</i>	153
<i>Mucor rouxii</i>	156	<i>Zygosaccharomyces fermentati</i>	153
 		<i>Zygosaccharomyces mandschuricus</i>	153
<i>Penicillium brevi-compactum</i>	152	<i>Zygosaccharomyces mellis</i>	153
<i>Penicillium cyclopium</i>	152	<i>Zygosaccharomyces nadsonii</i>	153
<i>Penicillium herquei</i>	75	<i>Zygosaccharomyces nussbaumeri</i>	153
<i>Phycomyces blakesleeanus</i>	156	<i>Zygosaccharomyces priorianus</i>	153
<i>Pichia farinosa</i>	146	<i>Zygosaccharomyces richteri</i>	153
<i>Pichia guilliermondii</i>	136	<i>Zygosaccharomyces rugosus</i>	153
<i>Pichia miso</i>	146, 147, 195	<i>Zygosaccharomyces wyocena</i>	153
<i>Pichia quercibus</i>	194, 196		
<i>Rhizopus oligosporus</i>	154		
<i>Rhizopus stolonifer</i>	156		

Table 4: Chemotaxonomy of fungi: group of species containing free mannitol (group D). * References are not given separately to the original publications antedating this review article.

<i>Acanthocystis petalooides</i>	72	<i>Blastocladiella emersonii</i>	156
<i>Acetabula vulgaris</i>	231 *	<i>Boletus aereus</i>	28
<i>Agaricus arvensis</i>	28	<i>Boletus appendiculatus</i>	231 *
<i>Agaricus bisporus</i>	166, 60, 81, 59, 61, 182, 84, 155	<i>Boletus aurantiacus</i>	231 *
<i>Agaricus campester</i>	72, 163	<i>Boletus badius</i>	231 *
<i>Agaricus campestris</i>	134, 63	<i>Boletus bovinus</i>	231 *, 103
<i>Agaricus eryngii</i>	231 *	<i>Boletus calopus</i>	231 *
<i>Agaricus silvaticus</i>	28	<i>Boletus chrysenteron</i>	231 *
<i>Agaricus silvicola</i>	72	<i>Boletus cyanescens</i>	231 *
<i>Agaricus xanthoderma</i>	72	<i>Boletus edulis</i>	231 *, 103, 28, 166
<i>Allomyces arbuscula</i>	156	<i>Boletus erythropus</i>	231 *, 155
<i>Alternaria alternata</i>	101	<i>Boletus luridus</i>	231 *, 28
<i>Alternaria tenuissima</i>	156	<i>Boletus luteus</i>	231 *, 103
<i>Amanita aspera</i>	231 *	<i>Boletus pachypus</i>	231 *
<i>Amanita bulbosa</i>	231 *	<i>Boletus pruinatus</i>	231 *
<i>Amanita caesarea</i>	231 *, 28	<i>Boletus rufus</i>	72, 28
<i>Amanita citrina</i>	72, 28	<i>Boletus satanas</i>	28
<i>Amanita muscaria</i>	231 *, 72, 28	<i>Boletus scaber</i>	231 *, 28
<i>Amanita pantherina</i>	231 *, 28	<i>Boletus subtomentosus</i>	231 *, 103
<i>Amanita porphyria</i>	28	<i>Boletus variegatus</i>	231 *
<i>Amanita rubescens</i>	231 *, 28	<i>Botrytis cinerea</i>	117, 154, 30
<i>Amanita phalloides</i>	72, 28	<i>Bulgaria inquinans</i>	231 *, 72
<i>Amanita vaginata</i>	231 *	<i>Byssochlamys fulva</i>	165
<i>Amanitopsis vaginata</i>	28	<i>Byssochlamys nivea</i>	156
<i>Armillaria caligata</i>	103	<i>Candida lipolytica</i>	197
<i>Armillaria mellea</i>	231 *, 155, 130, 79	<i>Candida utilis</i>	154
<i>Armillariella mellea</i>	72	<i>Candida zeylanoides</i>	86
<i>Ascobolus stercorarius</i>	156	<i>Cantharellus cibarius</i>	231 *, 166
<i>Aspergillus candidus</i>	175, 190, 143	<i>Cantharellus tubaeformis</i>	231 *
<i>Aspergillus clavatus</i>	98	<i>Chaetomium elatum</i>	156
<i>Aspergillus elegans</i>	23, 80	<i>Chaetomium globosum</i>	3, 154
<i>Aspergillus fischeri</i>	162	<i>Clavaria aurea</i>	72
<i>Aspergillus glaucus</i>	228	<i>Clavaria coralloides</i>	231 *
<i>Aspergillus nidulans</i>	23	<i>Clavaria corniculata</i>	96
<i>Aspergillus niger</i>	15, 117, 2	<i>Clavaria flava</i>	231 *
<i>Aspergillus oryzae</i>	192, 99	<i>Clavaria formosa</i>	231 *, 103
<i>Aspergillus terreus</i>	116	<i>Clavaria pistillaris</i>	231 *, 28
<i>Aspergillus versicolor</i>	85	<i>Claviceps curreyana</i>	47
<i>Aspergillus wentii</i>	24	<i>Claviceps nigricans</i>	47
		<i>Claviceps purpurea</i>	231 *, 215, 47, 48
		<i>Clitocybe aurantiaca</i>	28

- Clitocybe cyanophaea* 103
Clitocybe geotropa 231 *
Clitocybe infundibuliformis 72
Clitocybe nebularis 231 *, 72
Clitocybe odora 28
Clitocybe socialis 231 *
Coccidioides immitis 125
Cochliobolus miyabeanus 156
Colletotrichum gloeosporioides 117
Collybia butyracea 231 *
Collybia confluens 231 *
Collybia dryophila 231 *
Collybia erythropus 72
Collybia fusipes 231 *, 72
Collybia maculata 231 *
Collybia platyphylla 28
Coprinus atramentarius 231 *, 28, 154
Coprinus comatus 72, 28
Coprinus friesii 156
Coprinus micaceus 72, 28
Cordiceps capitata 199
Coriolus versicolor 28
Cortinarius alboviolaceus 28
Cortinarius berkeleyi 72
Cortinarius bivelus 231 *
Cortinarius bolaris 28
Cortinarius brunneus 231 *
Cortinarius bulliardii 72
Cortinarius collinitus 231 *, 28
Cortinarius elatior 231 *
Cortinarius fulmineus 28
Cortinarius infractus 231 *
Cortinarius largus 28
Cortinarius torvus 28
Cortinarius violaceus 72, 28
Cortinellus shiitake 103
Coryne sarcoïdes 96
Craterellus cornucopioides 28
Crepidotus nidulans 72
Cryptococcus neoformans 149
Dendryphiella salina 8, 97, 128, 155, 107
Diplodia viticola 191
Elaphomyces asperulus 231 *
Elaphomyces echinatus 231 *
Elaphomyces granulatus 231 *
Elaphomyces leveillei 231 *
Elaphomyces variegatus 231 *
Endothia parasitica 156
Entoloma clypeatum 28
Entoloma lividum 28
Entoloma nidorosum 231 *
Entoloma sinuatum 231 *
Epichloe typhina 203
Fistulina hepatica 28
Flammula spectabilis 72
Flammulina velutipes 108
Fomes ignarius 58
Fusarium solani 44
Galactinia olivacea 72
Geaster rufescens 231 *
Gelasinospora cerealis 154
Geotrichum candidum 42/1, 183, 52
Gibberella fujikuroi 156
Gomphidius roseus 28
Gomphidius viscidus 72
Helminthosporium geniculatum 24
Helminthosporium oryzae 30
Helvella esculenta 231 *
Hydnnum erinaceus 72
Hydnnum ferrugineum 231 *, 28
Hydnnum hybridum 231 *
Hydnnum imbricatum 154
Hydnnum repandum 231 *, 28, 166, 154
Hydnnum squamosum 231 *
Hygrophoropsis aurantiaca 28
Hygrophorus agathosmus 231 *
Hygrophorus cossus 231 *
Hygrophorus hypotheius 231 *
Hygrophorus nemoreus 72
Hygrophorus olivaceo-albus 231 *
Hygrophorus virgineus 231 *
Hypholoma fasciculare 231 *, 28
Hypholoma hydrophilum 28

- Hypholoma sublateritium* 231*, 103, 28
Inocybe praetervisa 28
Ixocomus bovinus 28
Ixocomus luteus 28
Ixocomus piperatus 28
Ixocomus variegatus 28
Laccaria laccata 28
Lactarius blennius 231*, 72, 28
Lactarius chrysorrheus 72, 28
Lactarius controversus 231*, 72
Lactarius deliciosus 231*, 72, 21, 28, 166
Lactarius musteus 28
Lactarius pallidus 231*, 72
Lactarius piperatus 231*, 103, 72
Lactarius plumbeus 72, 28
Lactarius pubescens 72
Lactarius pyrogalus 231*, 72
Lactarius quietus 231*
Lactarius rufus 231*, 28
Lactarius subdulcis 231*
Lactarius torminosus 231*, 72, 28
Lactarius turpis 231*
Lactarius uvidus 28
Lactarius vellereus 231*, 72, 28
Lactarius vietus 231*
Lactarius violascens 72
Lactarius volvulus 231*
Lactarius zonarius 231*, 72
Lentinus cochleatus 231*, 72
Lentinus edodes 137
Lenzites flaccida 28
Lenzites seapiaria 154
Lepidella vittadinii 72
Lepiota clypeolaria 72
Lepiota cristata 72
Lepiota excoriata 231*
Lepiota friesii 231*
Lepiota proceria 231*
Lepiota rhacodes 231*
Leucocoprinus rhacodes 72
Lycoperdon pusillum 231*
Lycoperdon pyriforme 72
Marasmius hariolorum 72
Melanopus squamosus 72
Microsporum gypseum 119
Monascus ruber 156
Morchella conica 231*
Morchella esculenta 231*
Morchella semilibera 231*
Mycena galericulata 96
Mycena pelianthina 231*
Myrothecium verrucaria 132
Nematoloma fasciculare 72
Neurospora crassa 193
Neurospora sitophila 220
Omphalia scyphoides 231*
Oudemansiella mucida 217
Panus conchatus 231*
Panus stipticus 231*
Paxillus atrotomentosus 231*, 72
Paxillus involutus 231*, 72
Penicillium brefeldianum 156
Penicillium chrysogenum 24, 12, 2
Penicillium griseofulvum 117
Penicillium italicum 156
Penicillium notatum 29
Penicillium oxalicum 156
Peziza badia 231*
Peziza nigra 231*
Peziza ochracea 231*
Peziza onotica 231*
Peziza venosa 231*
Phallus impudicus 231*
Pholiota aurivella 72
Pholiota lubrica 155
Pholiota mutabilis 103, 28
Pholiota praecox 72
Pholiota radicosa 231*
Pholiota togularis 231*
Phymatotrichum omnivorum 67
Pleurotus cornucopioides 72
Pleurotus dryinus 231*
Pleurotus japonicus 103
Pleurotus olearius 72
Pleurotus ostreatus 103, 72, 154, 156
Pleurotus serinus 103

- Pluteus cervinus* 72
Pluteus pellitus 72
Polyporus frondosus 72
Polyporus intybaceus 103
Polyporus officinalis 231 *
Polyporus pomaceus 28
Polyporus ptychogaster 231 *
Polyporus robustus 28
Polyporus umbellatus 72
Poria vincta 130
Psalliota arvensis 231 *, 103
Psalliota campestris 231 *, 103, 112
Psalliota placomyces 103
Psalliota silvicola 231 *
Puccinia coronata 200
Puccinia graminis tritici 161, 167, 227, 53, 39, 131, 130, 133
Pyrenophaeta terrestris 226
Pyricularia oryzae 155, 30
Rhizoctonia repens 178
Rhizoctonia solani 178, 156, 30
Rhizopogon luteolus 231 *
Russula adusta 231 *
Russula albonigra 28
Russula coerulea 28
Russula cyanoxantha 231 *, 28
Russula delica 231 *, 72
Russula drymeia 72
Russula fellea 231 *, 28
Russula foetens 231 *, 28
Russula fragilis 103, 28
Russula integra 231 *
Russula lepida 231 *, 28
Russula nigricans 231 *, 72, 28
Russula ochroleuca 231 *, 28, 96
Russula pseudoviolacea 28
Russula queletii 231 *
Russula sanguinea 72
Russula sardonia 28
Russula torulosa 28
Russula virescens 231 *
Sarcodon repandum 72
Sarcosphaera coronaria 72
Schizophyllum commune 144, 7, 181, 130
Scleroderma verrucosum 231 *, 72
Scleroderma vulgare 231 *, 72
Sclerotinia curreyana 47, 4
Sclerotinia sclerotiorum 120, 221
Sclerotinia tuberosa 231 *
Serpula lacrimans 36
Spongipellis spumeus 72
Sporobolomyces roseus 130
Stereostratum corticoides 200
Sterigmatocystis nigra 145, 72
Torulopsis anomala 149
Torulopsis mannitofaciens 151
Torulopsis nodaensis 149
Torulopsis versatilis 149
Tricholoma albo-brunneum 28
Tricholoma album 231 *
Tricholoma amarum 28
Tricholoma columbetta 231 *, 28
Tricholoma equestre 103, 28
Tricholoma flavobrunneum 231 *
Tricholoma nudum 231 *, 103
Tricholoma pessundatum 231 *
Tricholoma resplendens 231 *
Tricholoma russula 231 *, 103
Tricholoma sulfureum 231 *, 72
Tricholoma terreum 231 *
Tricholoma ustale 231 *, 103
Tuber cibarium 231 *
Tylopilus felleus 72
Ungulina betulina 72
Ustilago esculenta 41
Ustilago maydis 232, 155
Ustilago nuda 77
Verticillium albo-atrum 121
Verticillium dahliae 121, 30
Verticillium fungicola 156
Verticillium intertextum 156
Volvaria speciosa 72
Volvaria volvacea 231 *
Xerocomus badius 28, 154
Xerocomus chrysenteron 28
Xylaria polymorpha 231 *

Table 5: Chemosystematics of fungi: group of species containing free threitol, ribitol, xylitol, dulcitol, sorbitol, volemitol, *meso-glycero-ido-heptitol*, or *D-glycero-D-ido-heptitol* (group E)

<i>Agaricus bisporus</i>	59, 61	<i>Pichia farinosa</i>	146
<i>Armillaria mellea</i>	25, 155, 130	<i>Pichia guilliermondii</i>	136
<i>Boletus bovinus</i>	219	<i>Pichia miso</i>	146, 147
<i>Boletus erythropus</i>	155	<i>Pichia quercibus</i>	194
<i>Candida albicans</i>	158	<i>Pleurotus ostreatus</i>	156
<i>Candida diddensii</i>	78	<i>Poria vincta</i>	130
<i>Candida guilliermondii</i>	150, 170	<i>Psalliota campestris</i>	112
<i>Candida pelliculosa</i>	109	<i>Puccinia graminis tritici</i>	53, 131, 130, 133
<i>Candida polymorpha</i>	146, 148, 196	<i>Pyrenophaeta terrestris</i>	226
<i>Candida tropicalis</i>	146, 78	<i>Rhodosporidium toruloides</i>	78
<i>Candida utilis</i>	100	<i>Rhodotorula graminis</i>	55
<i>Chaetomium elatum</i>	156	<i>Saccharomyces acidifaciens</i>	196
<i>Circinella mucoroides</i>	51	<i>Saccharomyces carlsbergensis</i>	78
<i>Circinella muscae</i>	51	<i>Saccharomyces cerevisiae</i>	78
<i>Circinella rigida</i>	51	<i>Saccharomyces diastaticus</i>	78
<i>Circinella simplex</i>	51	<i>Saccharomyces rouxii</i>	146, 196, 78
<i>Circinella umbellata</i>	51	<i>Saccharomyces sake</i>	78
<i>Claviceps purpurea</i>	215, 48	<i>Saccharomycopsis fibuligera</i>	78
<i>Endothia parasitica</i>	156	<i>Schizophyllum commune</i>	130
<i>Geotrichum candidum</i>	183	<i>Schizosaccharomyces pombe</i>	78
<i>Hansenula anomala</i>	146	<i>Sclerotinia sclerotiorum</i>	221
<i>Kluyveromyces fragilis</i>	78	<i>Sporobolomyces roseus</i>	130
<i>Lactarius volemus</i>	31	<i>Syncephalastrum racemosum</i>	51
<i>Mucor miehei</i>	156, 157	<i>Torula utilis</i>	71
<i>Mucor rouxii</i>	156	<i>Torulopsis candida</i>	78
		<i>Torulopsis famata</i>	146
		<i>Torulopsis versatilis</i>	146
		<i>Trichosporon melibiosaceum</i>	78
		<i>Zygorhynchus moelleri</i>	156, 157

Discussion

Alditols in the free form have been found in autotrophs as well as heterotrophs, but especially occur in lower forms of life (see [88, 205, 159, 123, 204, 105]). In particular, most of the Eumycetes synthesize at least one polyol (mainly mannitol). Exceptions appear

to be taxonomically restricted, or can otherwise be rationalized (see below). Thus, the Oomycetes lack polyols as a whole (Tables 1 and 6), and the Zygomyco- and Hemiascomycetes do seem not to elaborate mannitol (Tables 3 and 6). Therefore, the metabolism of soluble carbohydrates of the large majority of the fungi (i) displays likeness to that of the invertebrates, especially the arthropods (cf. [43]) – where mannitol is, however, replaced by sorbitol that, conversely, is rare among the fungi (Table 5) – (ii) resembles that of certain heterofermentative Lactobacillariaceae (cf. [57]), and (iii) sets off nicely against the corresponding features of the higher plants, although a few of them do synthesize mannitol, dulcitol or sorbitol (see below); but, then, there is co-occurrence with sucrose, and not with trehalose as in the Eumycetes, bacteria and invertebrates (cf. [65]).

Table 6: The distribution of polyols (G, E, T, R, A, X, M, D, S, V, H, I, 0) in the fungi. The major taxa are according to AINSWORTH & BISBY's Dictionary of the Fungi (1971); within these, species are listed in alphabetical order.

G glycerol	X xylitol	H meso-glycero-ido-heptitol
E erythritol	M mannositol	I D-glycero-D-ido-heptitol
T threitol	D dulcitol	O no polyols
R ribitol	S sorbitol	
A arabitol	V volemitol	

MASTIGOMYCOTINA

Oomycetes

<i>Achlya radiosa</i>	0
<i>Albugo tragopogonis</i>	0
<i>Peronospora parasitica</i>	0
<i>Phytophthora cactorum</i>	0
<i>Phytophthora cinnamomi</i>	0
<i>Phytophthora infestans</i>	0
<i>Plasmopara viticola</i>	0

Chytridiomycetes

Allomyces arbuscula	G	M
Blastocladiella emersonii	G	M

ZYgomycotina

Zygomycetes

<i>Actinomucor elegans</i>	G	
<i>Circinella mucoroides</i>		R
<i>Circinella muscae</i>		R
<i>Circinella rigida</i>		R
<i>Circinella simplex</i>		R
<i>Circinella umbellata</i>		R
<i>Coccidioides immitis</i>		M

<i>Mortierella rammaniana</i>	G	
<i>Mucor miehei</i>	G	R
<i>Mucor mucedo</i>	G	
<i>Mucor rouxi</i>	G	R A
<i>Phycomyces blakesleeanus</i>	G	
<i>Rhizopus oligosporus</i>	G	
<i>Rhizopus stolonifer</i>	G	
<i>Syncephalastrum racemosum</i>		R
<i>Zygorhynchus moelleri</i>	G	R

ASCOMYCOTINA

Hemiascomycetes

<i>Byssochlamys fulva</i>			M
<i>Byssochlamys nivea</i>	G E		M
<i>Debaryomyces hansenii</i>	G	A	
<i>Debaryomyces sake</i>		A	
<i>Hansenula anomala</i>	G	A	D
<i>Hansenula suaveolens</i>		A	
<i>Kluyveromyces fragilis</i>		X	
<i>Pichia farinosa</i>		A	D
<i>Pichia guilliermondii</i>	R A		
<i>Pichia miso</i>	G E	A X	D
<i>Pichia quercibus</i>		A X	
<i>Saccharomyces acidifaciens</i>	G	A X	
<i>Saccharomyces carlsbergensis</i>		X	
<i>Saccharomyces cerevisiae</i>		X	
<i>Saccharomyces diastaticus</i>		X	
<i>Saccharomyces fragilis</i>	G	A	
<i>Saccharomyces mellis</i>		A	
<i>Saccharomyces monacensis</i>	G		
<i>Saccharomyces paradoxus</i>	G		
<i>Saccharomyces rouxii</i>	G	A X	D
<i>Saccharomyces sake</i>	G	X	
<i>Saccharomyces fibuligera</i>		X	
<i>Schizosaccharomyces pombe</i>		X	
<i>Zygosaccharomyces acidifaciens</i>	G	A	
<i>Zygosaccharomyces barkeri</i>	G	A	
<i>Zygosaccharomyces fermentati</i>		A	
<i>Zygosaccharomyces mandschuricus</i>		A	
<i>Zygosaccharomyces mellis</i>		A	
<i>Zygosaccharomyces nadsonii</i>	G	A	
<i>Zygosaccharomyces nussbaumeri</i>	G	A	
<i>Zygosaccharomyces priorianus</i>	G	A	
<i>Zygosaccharomyces richteri</i>	G	A	
<i>Zygosaccharomyces rugosus</i>	G	A	
<i>Zygosaccharomyces wyocena</i>	G	A	

Plectomycetes

<i>Monascus ruber</i>	G E	M
-----------------------	-----	---

Pyrenomycetes

<i>Chaetomium elatum</i>	T	A	M
<i>Chaetomium globosum</i>	G		M
<i>Claviceps curreyana</i>		A	M
<i>Claviceps nigricans</i>		A	M
<i>Claviceps purpurea</i>	G E T	A X	M D
<i>Cochliobolus miyabeanus</i>	G		M
<i>Endothia parasitica</i>	G	A	M S
<i>Epichloe typhina</i>			M
<i>Gelasinospora cerealis</i>	G		M
<i>Gibberella fujikuroi</i>	G	A	M
<i>Neurospora crassa</i>			M
<i>Neurospora sitophila</i>			M
<i>Xylaria polymorpha</i>			M

Discomycetes

<i>Acetabula vulgaris</i>			M
<i>Ascobolus stercorarius</i>			M
<i>Bulgaria inquinans</i>			M
<i>Coryne sarcoides</i>	G	A	M
<i>Elaphomyces asperulus</i>			M
<i>Elaphomyces echinatus</i>			M
<i>Elaphomyces granulatus</i>			M
<i>Elaphomyces leveillei</i>			M
<i>Elaphomyces variegatus</i>			M
<i>Galactinia olivacea</i>			M
<i>Helvella esculenta</i>			M
<i>Morchella conica</i>			M
<i>Morchella esculenta</i>			M
<i>Morchella semilibera</i>			M
<i>Peziza badia</i>			M
<i>Peziza nigra</i>			M
<i>Peziza ochracea</i>			M
<i>Peziza onotica</i>			M
<i>Peziza venosa</i>			M
<i>Sarcosphaera coronaria</i>			M
<i>Sclerotinia curreyana</i>		A	M
<i>Sclerotinia sclerotiorum</i>	G	R A X	M D S
<i>Sclerotinia tuberosa</i>			M
<i>Tuber cibarium</i>			M

DEUTEROMYCOTINA

Blastomycetes

<i>Candida albicans</i>	G	R A	
<i>Candida arborea</i>		A	
<i>Candida diiddensii</i>			X
<i>Candida guilliermondii</i>	G	R A X	
<i>Candida lipolytica</i>	E	A	M
<i>Candida parapsilosis</i>		A	
<i>Candida pelliculosa</i>			X
<i>Candida polymorpha</i>	E	R A X	D

<i>Candida pseudotropicalis</i>		A		
<i>Candida tropicalis</i>	G	A	X	D
<i>Candida utilis</i>	G	A	X	M D
<i>Candida zeylanoides</i>	E		M	
<i>Cryptococcus neoformans</i>	G		M	
<i>Rhodotorula graminis</i>	G	A		S
<i>Sporobolomyces roseus</i>	E	A	X	M S
<i>Torulopsis anomala</i>	G		M	
<i>Torulopsis candida</i>			X	
<i>Torulopsis famata</i>	E	A		D
<i>Torulopsis halophila</i>			A	
<i>Torulopsis magnoliae</i>	G			
<i>Torulopsis mannitofaciens</i>			A	M
<i>Torulopsis nodaensis</i>	G		A	M
<i>Torulopsis versatilis</i>	G	A	M D	
<i>Trichosporon melibiosaceum</i>			X	
<i>Trigonopsis variabilis</i>	E			

Hypocreomycetes

<i>Alternaria alternata</i>			M	
<i>Alternaria tenuissima</i>			A	M
<i>Aspergillus candidus</i>				M
<i>Aspergillus clavatus</i>	G E	R A	M	
<i>Aspergillus elegans</i>			M	
<i>Aspergillus fischeri</i>			M	
<i>Aspergillus glaucus</i>			M	
<i>Aspergillus nidulans</i>			M	
<i>Aspergillus niger</i>	G E	A	M	
<i>Aspergillus oryzae</i>			M	
<i>Aspergillus terreus</i>	E		M	
<i>Aspergillus versicolor</i>			M	
<i>Aspergillus wentii</i>	G		M	
<i>Botrytis cinerea</i>	G	A	M	
<i>Dendryphiella salina</i>	G	A	M	
<i>Fusarium solani</i>			M	
<i>Geotrichum candidum</i>	G E	R A	M	
<i>Helminthosporium geniculatum</i>	G		M	
<i>Helminthosporium oryzae</i>			M	
<i>Microsporum gypseum</i>			A	M
<i>Myrothecium verrucaria</i>				M
<i>Penicillium brefeldianum</i>	E		M	
<i>Penicillium brevi-compactum</i>	E			
<i>Penicillium chrysogenum</i>	G E	A	M	
<i>Penicillium cyclopium</i>	E			
<i>Penicillium griseofulvum</i>			A	M
<i>Penicillium herquei</i>	E			
<i>Penicillium italicum</i>	G E	A	M	
<i>Penicillium notatum</i>				M
<i>Penicillium oxalicum</i>	G E	A	M	
<i>Phymatotrichum omnivorum</i>				M
<i>Pyricularia oryzae</i>	G E	A	M	
<i>Rhizoctonia repens</i>				M
<i>Rhizoctonia solani</i>	G	A	M	

<i>Sterigmatocystis nigra</i>			M	
<i>Torula utilis</i>			D	
<i>Verticillium albo-atrum</i>	G E	A	M	
<i>Verticillium dahliae</i>	G E	A	M	
<i>Verticillium fungicola</i>		E	M	
<i>Verticillium intertextum</i>	G	A	M	
<i>Coelomycetes</i>				
<i>Colletotrichum gloeosporioides</i>		A	M	
<i>Diplodia viticola</i>		A	M	
<i>Pyrenopeziza terrestris</i>	G E	A	M	S

BASIDIOMYCOTINA

Teliomycetes

<i>Puccinia coronata</i>			M		
<i>Puccinia graminis tritici</i>	G E	R A X	M		S
<i>Rhodosporidium toruloides</i>			X		
<i>Stereostratum corticoides</i>				M	
<i>Ustilago esculenta</i>		E		M	
<i>Ustilago maydis</i>	G E		A	M	
<i>Ustilago nuda</i>	G E			M	

Hymenomycetes

<i>Acanthocystis petalooides</i>			M		
<i>Agaricus arvensis</i>			M		
<i>Agaricus bisporus</i>	G	A X	M		
<i>Agaricus campester</i>			M		
<i>Agaricus campestris</i>			M		
<i>Agaricus eryngii</i>			M		
<i>Agaricus silvaticus</i>			M		
<i>Agaricus silvicola</i>			M		
<i>Agaricus xanthoderma</i>			M		
<i>Amanita aspera</i>			M		
<i>Amanita bulbosa</i>			M		
<i>Amanita caesarea</i>			M		
<i>Amanita citrina</i>			M		
<i>Amanita muscaria</i>			M		
<i>Amanita pantherina</i>			M		
<i>Amanita phalloides</i>			M		
<i>Amanita porphyria</i>			M		
<i>Amanita rubescens</i>			M		
<i>Amanita vaginata</i>			M		
<i>Amanitopsis vaginata</i>			M		
<i>Armillaria caligata</i>			M		
<i>Armillaria mellea</i>	G E T	A X	M		S
<i>Armillariella mellea</i>			M		
<i>Boletus aereus</i>			M		
<i>Boletus appendiculatus</i>			M		
<i>Boletus aurantiacus</i>			M		

<i>Boletus badius</i>		M	
<i>Boletus bovinus</i>	A	M	S
<i>Boletus calopus</i>		M	
<i>Boletus chrysenteron</i>		M	
<i>Boletus cyanescens</i>		M	
<i>Boletus edulis</i>		M	
<i>Boletus erythropus</i>	G	A X	M
<i>Boletus luridus</i>		M	
<i>Boletus luteus</i>		M	
<i>Boletus pachypus</i>		M	
<i>Boletus pruinatus</i>		M	
<i>Boletus rufus</i>		M	
<i>Boletus satanas</i>		M	
<i>Boletus scaber</i>		M	
<i>Boletus subtomentosus</i>		M	
<i>Boletus variegatus</i>		M	
<i>Cantharellus cibarius</i>		M	
<i>Cantharellus tubaeformis</i>		M	
<i>Clavaria aurea</i>		M	
<i>Clavaria coraloides</i>		M	
<i>Clavaria corniculata</i>		M	
<i>Clavaria flava</i>		M	
<i>Clavaria formosa</i>		M	
<i>Clavaria pistillaris</i>		M	
<i>Clitocybe aurantiaca</i>		M	
<i>Clitocybe cyanophaea</i>		M	
<i>Clitocybe geotropa</i>		M	
<i>Clitocybe infundibuliformis</i>		M	
<i>Clitocybe nebularis</i>		M	
<i>Clitocybe odora</i>		M	
<i>Clitocybe socialis</i>		M	
<i>Collybia butyracea</i>		M	
<i>Collybia confluens</i>		M	
<i>Collybia dryophila</i>		M	
<i>Collybia erythropus</i>		M	
<i>Collybia fusipes</i>		M	
<i>Collybia maculata</i>		M	
<i>Collybia platyphylla</i>		M	
<i>Coprinus atramentarius</i>	A	M	
<i>Coprinus comatus</i>		M	
<i>Coprinus friesii</i>	G	M	
<i>Coprinus micaceus</i>		M	
<i>Cordiceps capitata</i>		M	
<i>Coriolus versicolor</i>		M	
<i>Cortinarius alboviolaceus</i>		M	
<i>Cortinarius berkeleyi</i>		M	
<i>Cortinarius bivelus</i>		M	
<i>Cortinarius bolaris</i>		M	
<i>Cortinarius brunneus</i>		M	
<i>Cortinarius bulliardii</i>		M	
<i>Cortinarius collinitus</i>		M	
<i>Cortinarius elatior</i>		M	
<i>Cortinarius fulmineus</i>		M	
<i>Cortinarius infractus</i>		M	
<i>Cortinarius largus</i>		M	

<i>Cortinarius torvus</i>	M
<i>Cortinarius violaceus</i>	M
<i>Cortinellus shiitake</i>	M
<i>Craterellus cornucopioides</i>	M
<i>Crepidotus nidulans</i>	M
<i>Entoloma clypeatum</i>	M
<i>Entoloma lividum</i>	M
<i>Entoloma nidorosum</i>	M
<i>Entoloma sinuatum</i>	M
<i>Fistulina hepatica</i>	A M
<i>Flammula spectabilis</i>	M
<i>Flammulina velutipes</i>	A M
<i>Fomes ignarius</i>	M
<i>Gomphidius roseus</i>	M
<i>Gomphidius viscidus</i>	M
<i>Hydnnum erinaceus</i>	M
<i>Hydnnum ferrugineum</i>	M
<i>Hydnnum hybridum</i>	M
<i>Hydnnum imbricatum</i>	M
<i>Hydnnum repandum</i>	M
<i>Hydnnum squamosum</i>	M
<i>Hygrophoropsis aurantiaca</i>	M
<i>Hygrophorus agathosmus</i>	M
<i>Hygrophorus cossus</i>	M
<i>Hygrophorus hypotheius</i>	M
<i>Hygrophorus nemoreus</i>	M
<i>Hygrophorus olivaceo-albus</i>	M
<i>Hygrophorus virgineus</i>	M
<i>Hypholoma fasciculare</i>	M
<i>Hypholoma hydrophilum</i>	M
<i>Hypholoma sublateritium</i>	M
<i>Inocybe praetervisa</i>	M
<i>Ixocomus bovinus</i>	M
<i>Ixocomus luteus</i>	M
<i>Ixocomus piperatus</i>	M
<i>Ixocomus variegatus</i>	M
<i>Laccaria laccata</i>	M
<i>Lactarius blennius</i>	M
<i>Lactarius chrysorrheus</i>	M
<i>Lactarius controversus</i>	M
<i>Lactarius deliciosus</i>	M
<i>Lactarius musteus</i>	M
<i>Lactarius pallidus</i>	M
<i>Lactarius piperatus</i>	M
<i>Lactarius plumbeus</i>	M
<i>Lactarius pubescens</i>	M
<i>Lactarius pyrogalus</i>	M
<i>Lactarius quietus</i>	M
<i>Lactarius rufus</i>	M
<i>Lactarius subdulcis</i>	M
<i>Lactarius torminosus</i>	M
<i>Lactarius turpis</i>	M
<i>Lactarius uvidus</i>	M
<i>Lactarius vellereus</i>	M
<i>Lactarius vietus</i>	M

<i>Lactarius violascens</i>			M	
<i>Lactarius volemus</i>			M	V
<i>Lactarius zonarius</i>			M	
<i>Lentinus cochleatus</i>			M	
<i>Lentinus edodes</i>		A	M	
<i>Lenzites flaccida</i>			M	
<i>Lenzites saeparia</i>	G	A	M	
<i>Lepidella vittadinii</i>			M	
<i>Lepiota clypeolaria</i>			M	
<i>Lepiota cristata</i>			M	
<i>Lepiota excoriata</i>			M	
<i>Lepiota friesii</i>			M	
<i>Lepiota procera</i>			M	
<i>Lepiota rhacodes</i>			M	
<i>Leucocoprinus rhacodes</i>			M	
<i>Marasmius hariovorum</i>			M	
<i>Melanopus squamosus</i>			M	
<i>Mycena galericulata</i>	G	A	M	
<i>Mycena pelianthina</i>			M	
<i>Nematoloma fasciculare</i>			M	
<i>Omphalia scyphoides</i>			M	
<i>Oudemansiella mucida</i>	G	A	M	
<i>Panus conchatus</i>			M	
<i>Panus stipticus</i>			M	
<i>Paxillus atrotomentosus</i>			M	
<i>Paxillus involutus</i>			M	
<i>Pholiota aurivella</i>			M	
<i>Pholiota lubrica</i>	E	A	M	
<i>Pholiota mutabilis</i>			M	
<i>Pholiota praecox</i>			M	
<i>Pholiota radicosa</i>			M	
<i>Pholiota togularis</i>			M	
<i>Pleurotus cornucopioides</i>			M	
<i>Pleurotus dryinus</i>			M	
<i>Pleurotus japonicus</i>			M	
<i>Pleurotus olearius</i>			M	
<i>Pleurotus ostreatus</i>	G E		M	S
<i>Pleurotus serinus</i>			M	
<i>Pluteus cervinus</i>			M	
<i>Pluteus pellitus</i>			M	
<i>Polyporus frondosus</i>			M	
<i>Polyporus intybaceus</i>			M	
<i>Polyporus officinalis</i>			M	
<i>Polyporus pomaceus</i>			M	
<i>Polyporus ptychogaster</i>			M	
<i>Polyporus robustus</i>			M	
<i>Polyporus umbellatus</i>			M	
<i>Poria vincta</i>	E	A X	M	
<i>Psalliota arvensis</i>			M	
<i>Psalliota campestris</i>		X	M	
<i>Psalliota placomyces</i>			M	
<i>Psalliota silvicola</i>			M	
<i>Russula adusta</i>			M	
<i>Russula albonigra</i>			M	
<i>Russula coerulea</i>			M	

<i>Russula cyanoxantha</i>				M
<i>Russula delica</i>				M
<i>Russula drymeia</i>				M
<i>Russula fellea</i>				M
<i>Russula foetens</i>				M
<i>Russula fragilis</i>				M
<i>Russula integra</i>				M
<i>Russula lepida</i>				M
<i>Russula nigricans</i>				M
<i>Russula ochroleuca</i>	E			M
<i>Russula pseudoviolacea</i>				M
<i>Russula queletii</i>				M
<i>Russula sanguinea</i>				M
<i>Russula sardonia</i>				M
<i>Russula torulosa</i>				M
<i>Russula virescens</i>				M
<i>Sarcodon repandum</i>				M
<i>Schizophyllum commune</i>	G E	A	X	M
<i>Serpula lacrimans</i>	G E	A		M
<i>Spongipellis spumeus</i>				M
<i>Tricholoma albo-brunneum</i>				M
<i>Tricholoma album</i>				M
<i>Tricholoma amarum</i>				M
<i>Tricholoma columbetta</i>				M
<i>Tricholoma equestre</i>				M
<i>Tricholoma flavobrunneum</i>				M
<i>Tricholoma nudum</i>				M
<i>Tricholoma pessundatum</i>				M
<i>Tricholoma resplendens</i>				M
<i>Tricholoma russula</i>				M
<i>Tricholoma sulfureum</i>				M
<i>Tricholoma terreum</i>				M
<i>Tricholoma ustale</i>				M
<i>Tylopilus felleus</i>				M
<i>Ungulina betulina</i>				M
<i>Volvaria speciosa</i>				M
<i>Volvaria volvacea</i>				M
<i>Xerocomus badius</i>	G			M
<i>Xerocomus chrysenteron</i>				M

Gasteromycetes

<i>Geaster rufescens</i>		M
<i>Lycoperdon pusillum</i>		M
<i>Lycoperdon pyriforme</i>		M
<i>Phallus impudicus</i>		M
<i>Rhizopogon luteolus</i>		M
<i>Scleroderma verrucosum</i>		M
<i>Scleroderma vulgare</i>		M

Whereas in vascular plants some cases are quoted where the distribution of alditol seems to follow a systematic pattern (the presence of mannitol is characteristic of the Oleaceae and Apiaceae, dulcitol has been found in certain Celastrales and Scrophulariales,

and sorbitol is synthesized by Rosaceae: for refs., see [177] and [122]), sugar alcohols have attracted hardly any attention in the taxonomy of fungi despite the fact that they represent the class of fungal compounds studied most widely (see Introduction).

Any consideration of polyols as taxonomic markers at higher levels of hierarchy requires their being not only widespread (what, indeed, holds: Tables 2 and 6) but also displaying constancy [54]. This appears to be the case: considering the factor most liable to influence the polyol composition of a fungus, i. e. the carbohydrate nutrient source, the polyol fraction expectedly varied quantitatively in response to changes in the medium composition inasmuch as mostly the synthesis of that alditol was favoured whose immediate sugar precursor served as the C-source (e. g. in *C. purpurea* [215], *P. terrestris* [226], and *S. sclerotiorum* [221]), whereas the general pattern of sugar alcohols remained constant under any of the conditions tested (studied extensively in *G. candidum* [42/1], *P. chrysogenum* [12], *S. sclerotiorum* [45], and *D. salina* [97], as well as in *A. bisporus*, *P. italicum*, and *Z. moelleri* [155]).

Using the criteria (i) absence of polyols (Table 1), (ii) presence of polyols, except mannitol (Table 3), and (iii) presence of mannitol (Table 4), three phenetic groups, displaying characters P_0 , P_1 , and P_2 , were established. These perfectly coincide with taxa, or groups of taxa, as delimited in some widely accepted conventional classifications, i. e., the Oomycetes, the Zyg- and Hemiascomyces, and the Chytridio-, Euasco-, Basidio-, and Deuteromycetes complex (for a discussion of the few apparent exceptions, see below). Thus, the P_0 group (Table 6) comprises only Oomycetes that also single out using three other biochemical markers, namely, the pathway of lysine biogenesis, the sedimentation pattern of enzymes of tryptophan biosynthesis, and the type of cell wall construction, with cellulose as the major microfibrillar component (characters L_1 , T_1 , and W_1 , respectively: Fig. 1). The very special position of the Oomycetes with respect to all other classes of Eumycetes is, furthermore, underlined by additional chemical traits, i. e. the molecular weight of the 25 S ribosomal RNA components (Oomycetes, $1.4\text{--}1.43 \times 10^6$; all other fungi, $1.3\text{--}1.36 \times 10^6$: [127]), significant amounts of hydroxyprolin in cell wall protein [10], the presence of histones [164], and the occurrence of desmosterol, 24-methylene-cholesterol, and fucosterol – in the absence of ergosterol [135, 38, 164, 222]. Also, they do not accumulate polyphosphates, in contrast to representatives of all other groups of fungi [42/2]. When considering all these features, together with characters P_0 , L_1 , T_1 and W_1 , it would be justified to ally the Oomycetes, or their respective ancestors, to green algae (see also [102, 164]), an opinion already brought forward on account of comparative morphology [76, 218, 40].

The P₁ group (Table 3) is homogeneous also with respect to the L character. It is subdivided into two taxa when considering the different states of characters T and W (T₂/T₃; W₂/W₃: chitosan/chitin vs mannan-glucan type as delimited by BARTNICKI-GARCIA [18, 19]; see Fig. 1) as well as the ability to synthesize trisporic acids [37, 210]. The phenetic groups thus obtained embody the Zygomycetes on the one hand, and the Hemiascomycetes on the other. In the former, accumulation of ribitol appears to prevail whereas arabitol is rare; for representatives of the Hemiascomycetes the reverse holds true (Table 6; see also [156]).

The component members of the Zyg- and Hemiascomycetes having polyol-synthesizing ability and state of character L in common with the Chytridio-, Euasco-, and Basidiomycetes indicates a monophyletic origin of all of these groups, with rhodophytean and/or phaeophytean types of organisms being discussed as precursors (cf. [164] and [142]). Because of their distinctive polyol pattern and other special chemical characteristics (see above and Fig. 1), the Zygomycetes did, however, probably separate early from the common ancestor pool originally shared with present-day higher fungi, as suggested already before [102, 19]. By the same token, the Hemiascomycetes would also represent a side-line from the main evolutionary trunk [102, 19]. When, additionally, considering characteristics of sexual reproductive behaviour, the Zygomycetes do, however, show a lower degree of phenetic and, hence, very likely also of cladistic relationship with the Euasco- and Basidiomycetes, although – on account of the P, T and W characters only – the diversity of the Zygomycetes from the latter two groups would be the same as that of the Hemiascomycetes. Supposing a closer kinship of the Ustilaginales with the Hemiascomycetes than with the other Basidiomycetes (cf. [76, 218]) and accounting for the fact that the *Ustilago* species studied are of the P₂-type (Table 6), one would have to conclude that the P₁-pattern of the Hemiascomycetes has arisen by a loss of the ability, or tendency, to synthesize mannitol. – For an interpretation of the fact that the P₁-group also contains (some) Blastomycetes, see below.

In contrast to all other groups of the lower fungi, the Chytridiomycetes posses the P₂ character ([156]; Table 6). An analogous situation exists with respect to the L, T and W characters (Fig. 1): in exhibiting the L₂/T₄/W₅ combination they are, thus, unique within the lower fungi and belong to the same phenetic group as the Euasco- and Basidiomycetes (as well as anamorphs assigned to them). Hence, chytridiomycetous fungi single out as the likely ancestors of these (Fig. 1), as proposed already on account of classical criteria (cf. [141]). For reasons detailed above, the Zygomycetes can, however, not represent a link between the Chytridiomycetes and the




Fig. 1. Chemotaxonomy and macroevolutionary scheme of the Eumycetes, as based on classical criteria and chemical traits (polyol character: states P_0 , P_1 , and P_2 [this work]; cell wall composition: types W_1 – W_5 , partly according to BARTNICKI-GARCIA [18, 19]; organization of the tryptophan pathway: categories T_1 – T_4 [102]; and lysine biosynthetic pathway: alternatives L_1 and L_2 [216]).

Euasco-, Basidio-, and Deuteromycetes group, an idea that has been strongly favoured by some classical mycologists [76, 110, 17], and rejected by others ([102, 19]; see also [141]).

Since the classification of fungi on account of their alditol composition affords groups that coincide almost completely with major taxa of a conventional, i. e. mainly morphologically-based classification (see caption to Fig. 1), and since the polyol pattern is retained even in dual organismic interactions (for refs. concerning phytopathogenic associations, see Table 7), the polyol character must be regarded as highly conservative and, therefore, entitles itself as a taxonomic marker (see also below). The correctness of this conclusion is underlined by the fact that the five situations not conforming to the general rule that Oomycetes are of the P₀-, Zygo- and Hemiascomycetes of the P₁-, and the other Eumycetes of the P₂-type, probably represent only apparent exceptions. The following reasons may account for these: (i) mis-quotation of the experimental organism (case *T. utilis* instead of *C. utilis* ([71]; see also above), or incomplete description of the material analyzed (case *Rhodosporidium toruloides*, a member of the Ustilaginales [14], where actually only the anamorph *Rhodotorula gracilis* has been studied [78], and which would, therefore, formally have to be included in the paragraph Blastomycetes of Table 6); (ii) incomplete analysis (*P. brevi-compactum* and *P. cyclopium* [152] as well as *P. herquei* [75] and possibly also *Rh. toruloides* anamorph [see also below] which should all be of the P₂-type); and (iii) doubtful position of species within the classification used here as the reference (cases *C. immitis* in the Zygomycetes, and *Byssochlamys* in the Hemiascomycetes, but both displaying the P₂- and not the P₁-character: see Table 6). The former has in the meantime been transferred to the Hyphomycetes [174]. Similarly, *Byssochlamys* is no longer held as belonging to the Hemiascomycetes, but to the Plectomycetes [185], where it is considered to be transitional between the Thermoascaceae and the Eurotiaceae [70].

The Blastomycetes is well known as the 'junk yard' of fungi growing in the yeast form that have a generally unknown teleomorph, or may not have any at all, and within which the "delimitation of a species is rather subjective", i. e. "largely depending on the insight and the ideas of the taxonomist studying the group" (cited from [124]). Heterogeneity even within a genus is, therefore, considered a common feature of the taxonomy of Imperfect Yeasts [124]. The *Candida-Torulopsis* group represents an extreme in this respect [213, 214]; in addition, the two are distinguished only by a "totally inadequate intergeneric criterion" [16]. – Hence, Blastomycetes is *eo ipso* an artificial taxon [124], its component members waiting for transfer into a systematically more established, natural (cf. [93])

Table 7: Polyol Synthesis in Phytopathogenic Associations

Fungus/Plant Association		Polyol pattern ^a	Reference
Claviceps purpurea/ <i>Secale</i> sp.		A M	56
Claviceps purpurea/ <i>Triticum</i> sp.		A M	56
Epichloe typhina/Agrocystis stolonifera		M	203
Erysiphe cichoracearum/			
<i>Cucumis sativus</i>	G E	A M	30
Erysiphe graminis hordei/Hordeum sp.		A M	64
Melampsora aecidiooides/			
<i>Populus canescens</i>		A M	169
Melampsora lini/ <i>Linum</i> sp.		A M	140
Microsphaera alphitoides/ <i>Quercus robur</i>		A M	92
Puccinia coronata/avenae/ <i>Avena sativa</i>		A M	138
Puccinia graminis tritici/			
<i>Triticum sativum</i>		M	129
Puccinia graminis tritici/ <i>Triticum</i> sp.	R A	M S	131
Puccinia graminis tritici/			
<i>Triticum aestivum</i>		A M	138
Puccinia malvacearum/ <i>Althaea rosea</i>		A M	139
Puccinia pelargonii-zonalis/			
<i>Pelargonium inquinans</i>		A M	139
Puccinia poarum/Tussilago farfara		A M	94, 95
Sclerotinia sclerotiorum/Daucus carota		M	45
Sclerotinia trifoliorum/Daucus carota		M	45
Uncinula necator/Vitis vinifera		A M	30, 33
Uromyces appendiculatus/			
<i>Vigna sesquipedalis</i>		A M	180
Ustilago esculenta/Zizania caduciflora	E	M	41

^a For abbreviations, see caption to Table 6

group. Considering now the highly predictive value of the polyol pattern with respect to the position of a species within the major (other) higher taxa in a conventional classification system of the fungi (see Table 6 and above) and, therefore, also its high *a priori* weight (cf. [93]) for the assignment of a species of unknown systematic position to one of these, the application of the polyol character suggests itself as an "objective criterion" (see Introduction) for the taxonomy of Imperfect Yeasts. By this token, *C. lipolytica*, *C. utilis*, and *C. zeylanoides* (Tables 4 and 6) would have to be associated with Basidiomycetes and not with Hemiascomycetes. For some candidas the genus *Leucosporidium* has been established [69]. Also *C. albicans* has been regarded as a yeast of heterobasidiomycetous affinity [211]. However, this has never been confirmed by others, and based on our analyses, *C. albicans* appears to be of the P₁-type (and this irrespective of growth form [158]) – in contrast to *T. versatilis* (syn. *T. anomala* [214], after the transfer of *Torulopsis* to the genus *Candida* [229], now *C. versatilis*, *T. nodaensis* (nom. invalid. [214] =

C. nodaensis [229]), and *T. mannitofaciens* (nom. nud., see [16]). Similarly, *S. roseus* has since quite some time been considered to be of basidiomycetous nature ([124]; see also [13, 189]). For *C. neoformans*, finally, teleomorphs have been found that form basidiospores – though in a somewhat extravagant manner – and for which the genus *Filobasidiella* has been created [113–115]. The likelihood of a (hetero)basidiomycetous kinship of *C. neoformans* is further supported by its possessing the T₂-type (as do some *Tremella*, *Ustilago*, *Sporobolomyces*, and *Rhodotorula* species [102]), in addition to the P₂-character (Fig. 1), as well as its developing an intensely dark-red colour upon exposition to aqueous solutions of stabilized aromatic diazonium salts [212]). The other Blastomycetes species listed in Table 6 would have to be linked to Hemiascomycetes – if the analyses of their polyol composition are complete, a condition which, at least in some cases, does not seem to have yet been met (e. g. for some of the species studied by SALEWSKI & al. [170], MIER SCH & al. [136], BERNARD & al. [22], and GONG & al. [78]; for further refs., see Table 3).

Acknowledgements

Financial support of this work by the Swiss National Science Foundation (grant no. 3.347–0.82), and the permission to use the computer equipment of the Department of Forestry, Canton of Lucerne, are gratefully acknowledged. It is also a pleasure to thank Professor D. H. LEWIS, University of Sheffield, for many useful suggestions, and Prof. Dr. E. MÜLLER, (Swiss Federal Institute of Technology Zürich) for advice in some taxonomic problems. Professor E. ROULEAU, Université de Montréal, kindly read the manuscript.

References

- ADLER, L. & GUSTAFSSON, L. (1980). Polyhydric alcohol production and intracellular amino acid pool in relation to halotolerance of the yeast *Debaryomyces hansenii*. – Arch. Microbiol. 124: 123–130.
- ADLER, L., PEDERSEN, A. & TUNBLAD-JOHANSSON, I. (1982). Polyol accumulation by two filamentous fungi grown at different concentrations of NaCl. – Physiol. Plant. 56: 139–142.
- ADOMAKO, D., KAYE, M. A. G. & LEWIS, D. H. (1971). Carbohydrate metabolism in *Chaetomium globosum*. I. Cellular carbohydrates of resting mycelium and their changes following utilization of hexoses. – New Phytol. 70: 51–59.
- AGGAB, A. M. & COOKE, R. C. (1981). Carbohydrate changes in germinating sclerotia of *Sclerotinia curreyana*. – Trans. Br. Mycol. Soc. 76: 147–149.
- AHEARN, D. G. (1983). The yeasts. – In: HOWARD, D. H. & HOWARD, L. F. (eds.), *Fungi Pathogenic for Humans and Animals, Part A Biology* (Mycology Series, vol. III): 73–80. Marcel Dekker Inc., New York.
- AINSWORTH, G. C., JAMES, P. W. & HAWKSWORTH, D. L. (1971). Ainsworth & Bisby's Dictionary of the Fungi, 6th edn. Commonwealth Mycological Institute, Kew, Surrey.
- AITKEN, W. B. & NIEDERPRUEM, D. J. (1970). Ultrastructural changes and biochemical events in basidiospore germination of *Schizophyllum commune*. – J. Bacteriol. 104: 981–988.

8. ALLAWAY, A. E. & JENNINGS, D. H. (1970). The influence of cations on glucose transport and metabolism by, and loss of sugar alcohols from, the fungus *Dendryphiella salina*. – New Phytol. 69: 581–593.
9. ARONSON, J. M. (1965). The cell wall. – In: AINSWORTH, G. C. & SUSSMAN, A. S. (eds.). The Fungi, vol. I: 49–76. Academic Press, New York.
10. ARONSON, J. M. & FULLER, M. S. (1969). Cell wall structure of the marine fungus, *Atkinsiella dubia*. – Arch. Mikrobiol. 68: 295–305.
11. ARPIN, N. & FIASSON, J. L. (1971). The pigments of the Basidiomycetes: their chemotaxonomic interest. – In: PETERSEN, R. H. (ed.). Evolution in the Higher Basidiomycetes: 63–98. University of Tennessee Press, Knoxville.
12. BALLIO, A., DI VITTORIO, V. & RUSSI, S. (1964). The isolation of trehalose and polyols from the conidia of *Penicillium chrysogenum* Thom. – Arch. Biochem. Biophys. 107: 177–183.
13. BANDONI, R. J., LOBO, K. J. & BREZDEN, S. A. (1971). Conjugation and chlamydospores in *Sporobolomyces odoratus*. – Can. J. Bot. 49: 683–686.
14. BANNO, I. (1967). Studies on the sexuality of *Rhodotorula*. – J. Gen. Appl. Microbiol. 13: 167–196.
15. BARKER, S. A., GOMEZ-SANCHEZ, A. & STACEY, M. (1958). Studies of *Aspergillus niger*. X. Polyol and disaccharide production from acetate. – J. Chem. Soc. 52: 2583–2586.
16. BARNETT, J. A., PAYNE, R. W. & YARROW, D. (1979). A Guide to Identifying and Classifying Yeasts. Cambridge University Press, Cambridge.
17. BARR, M. E. (1975). Perspectives in the Ascomycotina. – Mem. New York Bot. Gard. 28: 1–8.
18. BARTNICKI-GARCIA, S. (1968). Cell wall chemistry, morphogenesis, and taxonomy of fungi. – Annu. Rev. Microbiol. 22: 87–108.
19. — (1970). Cell wall composition and other biochemical markers in fungal phylogeny. – In: HARBORNE, J. B. (ed.). Phytochemical Phylogeny: 81–103. Academic Press, London.
20. BENEDICT, R. G. (1970). Chemotaxonomic relationships among the Basidiomycetes. – Adv. Appl. Microbiol. 13: 1–23.
21. BENEŠOVÁ, V., HEROUT, V. & ŠORM, F. (1954). O Rostlinných látkách III. O látkách z ryzce pravého (*Lactarius deliciosus* L.) – Chem. Listy 48: 882–885.
22. BERNARD, E. M., CHRISTIANSEN, K. J., TSANG, S.-F., KIEHN, T. E. & ARMSTRONG, D. (1981). Rate of arabinitol production by pathogenic yeast species. – J. Clin. Microbiol. 14: 189–194.
23. BIRKINSHAW, J. H., CHARLES, J. H. V., HETHERINGTON, A. C. & RAISTRICK, H. (1931). Studies in the biochemistry of micro-organisms. IX. On the production of mannitol from glucose by species of *Aspergillus*. – Trans. Roy. Soc. Lond. 220 B: 153–171.
24. BIRKINSHAW, J. H. & RAISTRICK, H. (1931). Studies in the biochemistry of micro-organisms. XVII. The products of glucose metabolism formed by various species of fungi. – Trans. Roy. Soc. Lond. 220 B: 331–354.
25. BIRKINSHAW, J. H., STICKINGS, C. E. & TESSIER, P. (1948). Biochemistry of the wood-rotting fungi. V. The production of D-threitol (L-erythritol) by *Armillaria mellea* (Vahl) Quél. – Biochem. J. 42: 329–332.
26. BLAKELEY, E. R. & SPENCER, J. F. T. (1962). Studies on the formation of D-arabitol by osmophilic yeasts. – Can. J. Biochem. Physiol. 40: 1737–1748.
27. BOKHARY, H. A. & COOKE, R. C. (1974). Translocation of [¹⁴C] glucose by *Phytophthora cactorum*. – Trans. Br. Mycol. Soc. 63: 535–540.
28. BONNET, J.-L. (1959). Application de la chromatographie sur papier à l'étude de divers champignons (Basidiomycètes-Hyménomycètes). – Thèse (Pharm.), Université de Paris.
29. BOONSAENG, V., SULLIVAN, P. A. & SHEPERD, M. G. (1976). Mannitol production in fungi during glucose catabolism. – Can. J. Microbiol. 22: 808–816.

30. BORASCHI-GAIA, C. (1984). Vorkommen von Polyolen und Trehalose bei phytopathogenen Pilzen – in Reinkultur und in der parasitären Situation. – M. Sc. thesis, University of Zürich.
31. BOURQUELOT, E. (1889). Les hydrates de carbone chez les champignons. – Bull. Soc. Mycol. Fr. 5: 132–163.
32. BRACONNOT, H. (1811). Recherches analytiques sur la nature des champignons. – Ann. Chim. (Paris) 79: 265–304.
33. BREM, S., RAST, D. M. & RUFFNER, H. P. (1986). Partitioning of photosynthate in leaves of *Vitis vinifera* infected with *Uncinula necator* or *Plasmopara viticola*. – Physiol. Molec. Plant Pathol. 29: 285–291.
34. BRESINSKY, A. (1977). Chemotaxonomie der Pilze. – In: FREY, W., HURKA, H. & OBERWINKLER, F. (eds.). Beiträge zur Biologie der niederen Pflanzen: 25–42. Gustav Fischer, Stuttgart.
35. BRIMACOMBE, J. S. & WEBBER, J. M. (1972). Alditols and derivatives. – In: PIGMAN, W. & HORTON, D. (eds.). The Carbohydrates, vol. IA: 479–518. Academic Press, New York.
36. BROWNLEE, C. & JENNINGS, D. H. (1981). The content of soluble carbohydrates and their translocation in mycelium of *Serpula lacrimans*. – Trans. Br. Mycol. Soc. 77: 615–619.
37. BU'LOCK, J. P. (1976). Hormones in fungi. – In: SMITH, J. E. & BERRY, D. R. (eds.). The Filamentous Fungi, vol. II: 345–368. Edward Arnold, London.
38. BU'LOCK, J. P. & OSAGIE, A. U. (1976). Sterol biosynthesis via cycloartenol in *Saprolegnia*. – Phytochemistry 15: 1249–1251.
39. BURGER, A., PRINZING, A. & REISENER, H. J. (1972). Untersuchungen über den Glucosestoffwechsel keimender Uredosporen von *Puccinia graminis* var. *tritici*. – Arch. Mikrobiol. 83: 1–16.
40. BURNETT, J. H. (1976). Fundamentals of Mycology, 2nd edn. Edward Arnold, London.
41. CHAN, Y.-S. & THROWER, L. B. (1980). The host-parasite relationship between *Zizania caduciflora* Turcz. and *Ustilago esculenta* P. Henn. III. Carbohydrate metabolism of *U. esculenta* and the host-parasite combination. – New Phytol. 85: 217–224.
- 42/1. CHANG, S.-C. & LI, K.-H. (1964). Identification of D-mannitol in *Geotrichum candidum* Link and the mechanism of its formation. – Sci. Sin. 13: 621–630.
- 42/2. CHILVERS, G. A., LAPEYRIE, F. F. & DOUGLASS, P. A. (1985). A contrast between Oomycetes and other taxa of mycelial fungi in regard to metachromatic granule formation. – New Phytol. 99: 203–210.
43. CHIPPENDALE, G. M. (1978). The functions of carbohydrates in insect life processes. – In: ROCKSTEIN, M. (ed.). Biochemistry of Insects: 1–55. Academic Press, New York.
44. COCHRANE, V. W., COCHRANE, J. C., COLLINS, C. B. & SERAFIN, F. G. (1963). Spore germination and carbon metabolism in *Fusarium solani*. II. Endogenous respiration in relation to germination. – Am. J. Bot. 50: 806–814.
45. COOKE, R. C. (1969). Changes in soluble carbohydrates during sclerotia formation by *Sclerotinia sclerotiorum* and *S. trifoliorum*. – Trans. Br. Mycol. Soc. 53: 77–86.
46. — (1977). The Biology of Symbiotic Fungi. John Wiley, New York.
47. COOKE, R. C. & MITCHELL, D. T. (1969). Sugars and polyols in sclerotia of *Claviceps purpurea*, *C. nigricans*, and *Sclerotinia curreyana* during germination. – Trans. Br. Mycol. Soc. 52: 356–372.
48. CORBETT, K., DICKERSON, A. G. & MANTLE, P. G. (1975). Metabolism of the germinating sclerotium of *Claviceps purpurea*. – J. Gen. Microbiol. 90: 55–68.
49. CORINA, D. L. & MUNDAY, K. A. (1971). Studies on polyol function in *Aspergillus clavatus*: a role for mannitol and ribitol. – J. Gen. Microbiol. 69: 221–227.

50. CULLEN, J. (1968). Botanical problems of numerical taxonomy. – In: HEYWOOD, V. H. (ed.). Modern Methods in Plant Taxonomy: 175–183. Academic Press, London.
51. CURTIS, F. C., LEWIS, D. H. & COOKE, R. C. (1980). Distribution of a pentitol in the Mucorales. – Trans. Br. Mycol. Soc. 74: 421–423.
52. DA COSTA, M. S. & NIEDERPRUEM, D. J. (1982). Arabitol accumulation in *Geotrichum candidum*. – Arch. Mikrobiol. 131: 283–286.
53. DALY, J. M., KNOCHE, H. W. & WIESE, M. V. (1967). Carbohydrate and lipid metabolism during germination of uredospores of *Puccinia graminis tritici*. – Plant Physiol. 42: 1633–1642.
54. DAVIS, P. H. & HEYWOOD, V. H. (1963). Principles of Angiosperm Taxonomy. Oliver & Boyd, Edinburgh.
55. DEINEMA, M. H. & LANDHEER, C. A. (1960). Extracellular lipid production by a strain of *Rhodotorula graminis*. – Biochim. Biophys. Acta 37: 178–179.
56. DICKERSON, A. G., MANTLE, P. G. & NISBET, L. J. (1976). Carbon assimilation by *Claviceps purpurea* growing as a parasite. – J. Gen. Microbiol. 97: 267–276.
57. DOELLE, H. W. (1975). Bacterial Metabolism: pp. 631. Academic Press, New York.
58. DOMINGUEZ, X. A., BUTRUILLE, D., ZAMUDIO, A., REYES, C. & CASTILLO, J. T. (1972). Ergosterol and L-mannitol dans des champignons parasites. – Phytochemistry 11: 2616.
59. DÜTSCH, G. A. (1971). Biogenese von Mannit und ihre physiologische Bedeutung in *Agaricus bisporus*. – Ph. D. thesis, University of Zürich.
60. DÜTSCH, G. A. & RAST, D. (1969). Biosynthese von Mannitol in *Agaricus bisporus*. – Arch. Mikrobiol. 65: 195–207.
61. — (1972 a). Über die Rolle der Mannitbildung in *Agaricus bisporus*. – Mushroom Sci. 8: 387–397.
62. — (1972 b). Biochemische Beziehung zwischen Mannitbildung und Hexose-monophosphatzzyklus in *Agaricus bisporus*. – Phytochemistry 11: 2677–2681.
63. EDMUNDOWICZ, J. M. & WRISTON, J. C. (1963). Mannitol dehydrogenase from *Agaricus campestris*. – J. Biol. Chem. 238: 3539–3541.
64. EDWARDS, H. H. & ALLEN, P. J. (1966). Distribution of the products of photosynthesis between powdery mildew and barley. – Plant Physiol. 41: 683–688.
65. ELBEIN, A. D. (1974). The metabolism of α,α -trehalose. – In: TRIPSON, R. S. & HORTON, D. (eds.). Advances in Carbohydrate Chemistry and Biochemistry, vol. 30: 227–256. Academic Press, New York.
66. ERDTMAN, H. (1963). Some aspects of chemotaxonomy. – In: SWAIN, T. (ed.). Chemical Plant Taxonomy: 89–125. Academic Press, London.
67. ERGLE, D. R. (1948). The carbohydrate metabolism of germinating *Phycomyces trichocomoides* sclerotia with special reference to glycogen. – Phytopathology 38: 142–151.
68. ERMAKOVA, I. T., LENSIKIH, G. V., GAIDEREKO, T. A., MEDVEDEVA, T. N., LITVINOVA, E. G. & KORPACHEV, A. V. (1984). Biosynthesis of polyols by the yeast *Candida guilliermondii* in media with different carbon sources. – Mikrobiologiya (Engl. Transl.) 53: 663–668.
69. FELL, J. W., STATZELL, A. C., HUNTER, I. L. & PHAFF, H. J. (1969). *Leucosporidium* gen. n., the heterobasidiomycetous stage of several yeasts of the genus *Candida*. – Antonie van Leeuwenhoek 35: 433–462.
70. FENNELL, D. J. (1973). Plectomycetes; Eurotiales. – In: AINSWORTH, G. C., SPARROW, F. K. & SUSSMAN, A. S. (eds.). The Fungi, vol. IVA: 45–68. Academic Press, New York.
71. FINK, H. & JUST, F. (1938). Zur Biochemie der *Torula utilis*. 1. Mitteilung: die Auffindung von Dulcitol in Holzzuckerhefe. – Biochem. Z. 296: 306–314.
72. FRÉREjacque, M. (1939 a). Le mannitol chez les champignons. – Rev. Mycol. 4: 88–97.

73. — (1939 b). Présence du d-arabitol dans *Fistulina hepatica*. — C. R. Acad. Sci. (Paris) 208: 1123–1124.
74. — (1943). Sur la présence de l'arabitol dans *Boletus bovinus* L. — C. R. Acad. Sci. (Paris) 217: 251–252.
75. GALARRAGA, J. A., NEILL, K. G. & RAISTRICK, H. (1955). Studies in the biochemistry of micro-organisms. XCVI. The colonizing matters of *Penicillium herquei* Bainier & Sartory. — Biochem. J. 61: 456–464.
76. GAUMANN, E. A. (1964). Die Pilze, 2nd edn. Birkhäuser, Basel.
77. GAUNT, R. E. & MANNERS, J. G. (1973). The production of trehalose and polyols by *Ustilago nuda* in culture and their utilization in healthy and infected wheat plants. — New Phytol. 72: 321–327.
78. GONG, C.-S., CHEN, L. F. & TSAO, G. T. (1981). Quantitative production of xylitol producing yeast mutant *Candida tropicalis* HXP 2. — Biotechnol. Lett. 3: 130–135.
79. GRANLUND, H. I., JENNINGS, D. H. & THOMPSON, W. (1985). Translocation of solutes along rhizomorphs of *Armillaria mellea*. — Trans. Br. Mycol. Soc. 84: 111–119.
80. GRÉGOIRE, P. E. (1949). Présence de mannitol dans le mycelium d'*Aspergillus elegans gasperrini*. — Bull. Soc. Chim. Biol. 31: 801–803.
81. GRIFFIN, P. F. S., BRENNAN, P. J. & LÖSEL, D. M. (1970). Free lipids and carbohydrates of *Agaricus bisporus* mycelium. — Biochem. J. 119: 11 P–12 P.
82. HAJNY, G. J., HENDERSHOT, W. F. & PETERSON, W. H. (1960). Factors affecting glycerol production by a newly isolated osmophilic yeast. — Appl. Microbiol. 8: 5–11.
83. HALL, R. (1969). Molecular approaches to taxonomy of fungi. — Bot. Rev. 35: 285–304.
84. HAMMOND, J. B. W. & NICHOLS, R. (1976). Carbohydrate metabolism in *Agaricus bisporus* (Lange) Sing.: changes in soluble carbohydrates during growth of the mycelium and sporophore. — J. Gen. Microbiol. 93: 309–320.
85. HATSUDA, Y. & KUYAMA, S. (1956). Metabolic products of *Aspergillus versicolor*. — Chem. Abstr. 50: 15522.
86. HATTORI, K. & SUZUKI, T. (1975). Enzymatic studies on the conversion of erythritol fermentation to D-mannitol fermentation in n-alkane-grown yeast *Candida zeylanoides*. — Agric. Biol. Chem. (Jap.) 39: 57–61.
87. HAY, G. W., LEWIS, B. A. & SMITH, F. (1963). Thin-film chromatography in the study of carbohydrates. — J. Chromatogr. 11: 479–486.
88. HEGNAUER, R. (1962). Chemotaxonomie der Pflanzen, vol. I. Birkhäuser, Basel.
89. — (1967). Chemical characters in plant taxonomy: some possibilities and limitations. — Pure Appl. Chem. 14: 173–187.
90. HEINEMANN, P. & CASIMIR, J. (1961). Distribution des acides aminés libres dans le genre *Agaricus* Fr. sensu stricto (= *Psalliota*). — Rev. Mycol. 26: 24–33.
91. HEROUT, V. (1973). A chemical compound as taxonomical character. — In: BENDZ, G. & SANTESSON, J. (eds.). Chemistry in Botanical Classification: 55–62. Nobel Symposium 25, Almqvist-Wiksells, Uppsala.
92. HEWITT, H. G. & AYRES, P. G. (1976). Effect of infection by *Microsphaera alphitoides* (powdery mildew) on carbohydrate levels and translocation in seedlings of *Quercus robur*. — New Phytol. 77: 379–390.
93. HEYWOOD, V. H. (1966). Phytochemistry and taxonomy. — In: SWAIN, T. (ed.). Comparative Phytochemistry: 1–20. Academic Press, London.
94. HOLLIGAN, P. M., CHEN, C. & LEWIS, D. H. (1973). Changes in the carbohydrate composition of leaves of *Tussilago farfara* during infection by *Puccinia poarum*. — New Phytol. 72: 947–955.
95. HOLLIGAN, P. M., CHEN, C., McGEE, E. E. M. & LEWIS, D. H. (1974). Carbohydrate metabolism in healthy and rusted leaves of coltsfoot. — New Phytol. 73: 881–888.

96. HOLLIGAN, P. M. & DREW, E. A. (1971). Routine analysis by gas-liquid chromatography of soluble carbohydrates in extracts of plant tissues. II: Quantitative analysis of standard carbohydrates, and the separation and estimation of soluble sugars and polyols from a variety of plant tissues. – New Phytol. 71: 271–297.
97. HOLLIGAN, P. M. & JENNINGS, D. H. (1972). Carbohydrate metabolism in the fungus *Dendryphiella salina*. II. The influence of different carbon and nitrogen sources on the accumulation of mannitol and arabitol. – New Phytol. 71: 583–594.
98. HOLLIGAN, P. M. & LEWIS, D. H. (1973). The soluble carbohydrates of *Aspergillus clavatus*. – J. Gen. Microbiol. 75: 155–159.
99. HORIKOSHI, K., IIDA, S. & IKEDA, Y. (1965). Mannitol and mannitol dehydrogenases in conidia of *Aspergillus oryzae*. – J. Bacteriol. 89: 326–330.
100. HORITSU, H., TOMOEDA, M. & KUMAGAI, K. (1968). Pentose metabolism in *Candida utilis*. IV. NADP specific polyol dehydrogenase. – Agric. Biol. Chem. (Jap.) 32: 514–517.
101. HULT, K. & GATENBECK, S. (1978). Production of NADPH in the mannitol cycle and its relation to polyketide formation in *Alternaria alternata*. – Eur. J. Biochem. 88: 607–612.
102. HUTTER, R. & DEMOSS, J. A. (1967). Organization of the tryptophan pathway: a phylogenetic study of the fungi. – J. Bacteriol. 94: 1896–1907.
103. INAGAKI, S. (1934). Untersuchung über Bestandteile der in Japan einheimischen Pilze (1. Mitteilung). Mannitgehalt der Pilze. – J. Pharm. Soc. (Jap.) 54: 125–127.
104. INGRAM, J. M. & WOOD, W. A. (1965). Enzymatic basis for D-arabitol production by *Saccharomyces rouxii*. – J. Bacteriol. 89: 1186–1194.
105. JENNINGS, D. H. (1984). Polyol metabolism in fungi. – Adv. Microbial Physiol. 25: 149–193.
106. JENNINGS, D. H. & AUSTIN, S. (1973). The stimulatory effect of the non-metabolized sugar 3-O-methyl glucose on the conversion of mannitol and arabitol to polysaccharide and other insoluble compounds in the fungus *Dendryphiella salina*. – J. Gen. Microbiol. 75: 287–294.
107. JENNINGS, D. H. & THORNTON, J. D. (1984). Carbohydrate metabolism in the fungus *Dendryphiella salina*. VII. The effect of L-sorbose on ethanol-soluble carbohydrate. – New Phytol. 98: 399–403.
108. KITAMOTO, Y. & GRUEN, H. E. (1976). Distribution of cellular carbohydrates during development of the mycelium and fruitbodies of *Flammulina velutipes*. – Plant Physiol. 58: 485–491.
109. KITPREECHAVANICH, V., HAYASHI, M., NISHIO, N. & NAGAI, S. (1984). Conversion of D-xylose into xylitol by xylose reductase from *Candida pelliculosa* coupled with the oxidoreductase system of methanogen strain HU. – Biotechnol. Lett. 6: 651–655.
110. KLEIN, R. M. & CRONQUIST, A. (1976). A consideration of the evolutionary and taxonomic significance of some biochemical, micromorphological, and physiological characters in the Thallophytes. – Quart. Rev. Biol. 42: 105–296.
111. KNOP, H. & SCHNEDERMANN, A. (1844). Über den Mannit. – J. Prakt. Chem. 32: 411–416.
112. KRATZL, K. & SILBERNAGEL, H. (1963). Über das Vorkommen von Xylit im Speisepilz Champignon (*Psalliota campestris*). – Naturwissenschaften 50: 154.
113. KWON-CHUNG, K. J. (1975). A new genus, *Filobasidiella*, the perfect state of *Cryptococcus neoformans*. – Mycologia 67: 1197–1200.
114. — (1976 a). Morphogenesis of *Filobasidiella neoformans*, the sexual state of *Cryptococcus neoformans*. – Mycologia 68: 821–833.
115. — (1976 b). A new species of *Filobasidiella*, the sexual state of *Cryptococcus neoformans* B and C serotypes. – Mycologia 68: 942–946.

116. LAHOZ, R. (1967). Quantitative changes in the content of non-nitrogenous compounds during autolysis of *Aspergillus terreus*. – J. Gen. Microbiol. 46: 451–456.
117. LAHOZ, R., BALLESTEROS, A. M. & IBEAS, J. G. (1971). Chemical and physiological changes in filamentous fungi during autolysis. X. Changes in the concentration of polyols in autolyzing cultures of several fungi. – Mycopathol. Mycol. Appl. 43: 223–228.
118. LEDERER, E. (1938). Sur les caroténoïdes des cryptogames. – Bull. Soc. Chim. Biol. 20: 611–634.
119. LEIGHTON, T. J., STOCK, J. J. & KELLN, R. A. (1970). Macroconidial germination in *Microsporum gypseum*. – J. Bacteriol. 103: 439–446.
120. LE TOURNEAU, D. (1966). Trehalose and acyclic polyols in sclerotia of *Sclerotinia sclerotiorum*. – Mycologia 58: 934–942.
121. — (1974). Alcohol soluble sugars and sugar alcohols produced from glucose by *Verticillium* spp. – Trans. Br. Mycol. Soc. 62: 619–622.
122. LEWIS, D. H. (1984). Physiology and metabolism of alditols. – In: LEWIS, D. H. (ed.). Storage Carbohydrates in Vascular Plants: 157–179. Society for Exp. Biol. Seminar Series 19. Cambridge University Press, Cambridge.
123. LEWIS, D. H. & SMITH, D. C. (1967). Sugar alcohols (polyols) in fungi and green plants. I. Distribution, physiology and metabolism. – New Phytol. 66: 143–184.
124. LODDER, J. (1970). General classification of the yeasts. – In: LODDER, J. (ed.). The Yeasts: 1–33. North-Holland Publishing Company, Amsterdam.
125. LONES, G. W. & PEACOCK, C. (1964). Metabolism of mannitol by *Coccidioides immitis*. – J. Bacteriol. 87: 1114–1117.
126. LONG, D. E. & COOKE, R. C. (1974). Carbohydrate composition and metabolism of *Senecio squalidus* L. leaves infected with *Albugo tragopogonis* (Pers.) S. F. Gray. – New Phytol. 73: 889–899.
127. LOVETT, J. S. & HASELBY, J. A. (1971). Molecular weights of the ribosomal ribonucleic acid of fungi. – Arch. Mikrobiol. 80: 191–204.
128. LOWE, D. A. & JENNINGS, D. H. (1975). Carbohydrate metabolism in the fungus *Dendryphiella salina*. V. The pattern of label in arabitol and polysaccharide after growth in the presence of specifically labelled carbon sources. – New Phytol. 74: 67–79.
129. LUNDERSTAEDT, L. (1966). Effect of rust infection on hexokinase activity and carbohydrate dissimilation in primary leaves of wheat. – Can. J. Bot. 44: 1345–1364.
130. MACLEAN, D. J. (1982). Axenic culture and metabolism of rust fungi. – In: SCOTT, K. J. & CHAKRAVORTY, A. K. (eds.). The Rust Fungi: 37–120. Academic Press, London.
131. MACLEAN, D. J. & SCOTT, K. J. (1976). Identification of glucitol (sorbitol) and ribitol in a rust fungus, *Puccinia graminis* f. sp. *tritici*. – J. Gen. Microbiol. 97: 83–89.
132. MANDELS, G. R., VITOLS, R. & PARRISH, F. W. (1965). Trehalose as an endogenous reserve in spores of *Myrothecium verrucaria*. – J. Bacteriol. 90: 1589–1598.
133. MANNERS, J. M., MACLEAN, D. J. & SCOTT, K. J. (1984). Hexitols as major intermediates of glucose assimilation by mycelium of *Puccinia graminis*. – Arch. Microbiol. 139: 158–161.
134. MCCONNELL, J. E. W. & ESSELEN, W. B. (1947). Carbohydrates in cultivated mushrooms (*Agaricus campestris*). – Food Res. 12: 118–121.
135. MCCORKINDALE, N. J. (1976). The biosynthesis of terpenes and steroids. – In: SMITH, J. E. & BERRY, D. R. (eds.). The Filamentous Fungi, vol. II: 369–422. Edward Arnold, London.
136. MIERSCH, J., LAPF, H. & REINBOTHE, H. (1980). Ribitstoffwechsel in der flavinogenen Hefe *Pichia guilliermondii* Wickerham. – Biochem. Physiol. Pflanz. 175: 732–744.

137. MIKAMO, M. (1974). A sugar alcohol in *Lentinus edodes* (Berk.). – J. Agric. Chem. Soc. (Jap.) 48: 69–71.
138. MITCHELL, D. T., FUNG, A. K. & LEWIS, D. H. (1978). Changes in the ethanol-soluble carbohydrate composition and acid invertase in infected first leaf tissues susceptible to crown rust of oat and wheat stem rust. – New Phytol. 80: 381–392.
139. MITCHELL, D. T. & ROBERTS, S. M. (1973). Carbohydrate composition of tissues infected by the autoecious rusts, *Puccinia pelargonii-zonalis* and *P. malvacearum*. – Physiol. Plant Pathol. 3: 481–488.
140. MITCHELL, D. T. & SHAW, M. (1968). Metabolism of glucose-¹⁴C, pyruvate-¹⁴C, and mannitol-¹⁴C by *Melampsora lini*. II. Conversion to soluble products. – Can. J. Bot. 46: 453–460.
141. MÜLLER, E. (1977). Systemfragen bei Ascomyceten. – In: FREY, W., HURKA, H. & OBERWINKLER, F. (eds.). Beiträge zur Biologie der niederen Pflanzen: 43–57. Gustav Fischer, Stuttgart.
142. MÜLLER, E. & LOEFFLER, W. (1982). Mykologie, 4th edn., Georg Thieme, Stuttgart.
143. NELSON, G. E. N., JOHNSON, D. E. & CIEGLER, A. (1971). Production of D-mannitol by conidia of *Aspergillus candidus*. – Appl. Microbiol. 22: 484–485.
144. NIEDERPRÜM, D. J. & HUNT, S. (1967). Polyols in *Schizophyllum commune*. – Am. J. Bot. 54: 241–245.
145. OBATON, M. (1932). Le mannitol, aliment de choix pour le *Sterigmatocystis nigra*. – C. R. Acad. Sci. (Paris) 194: 302–304.
146. ONISHI, H. (1960). Studies on osmophilic yeasts. VII. Polyalcohol production by various genera and species of yeasts. – Bull. Agric. Soc. (Jap.) 24: 131–140.
147. ONISHI, H. & PERRY, M. B. (1965). The production of *meso-glycero-ido-heptitol* and *D-glycero-D-ido-heptitol* by *Pichia miso*. – Can. J. Microbiol. 11: 929–934.
148. ONISHI, H. & SUZUKI, T. (1966). The production of xylitol, L-arabinitol and ribitol by yeasts. – Agric. Biol. Chem. (Jap.) 30: 1139–1144.
149. — (1968). Production of D-mannitol and glycerol by yeasts. – Appl. Microbiol. 16: 1847–1852.
150. — (1969). Microbial production of xylitol from glucose. – Appl. Microbiol. 18: 1031–1035.
151. — (1970). Production of erythritol, D-arabitol, D-mannitol and a heptitol-like compound from glycerol by yeasts. – J. Ferment. Technol. 48: 563–566.
152. OXFORD, A. E. & RAISTRICK, H. (1935). Studies in the biochemistry of micro-organisms. XLVI. *i*-Erythritol, a metabolic product of *Penicillium brevicompactum* Diercks and *P. cyclopium* Westling. – Biochem. J. 29: 1599–1601.
153. PETERSON, W. H., HENDERSHOT, W. F. & HAJNY, G. J. (1958). Factors affecting production of glycerol and D-arabitol by representative yeasts of the genus *Zygosaccharomyces*. – Appl. Microbiol. 6: 349–357.
154. PFYFFER VON ALTISHOFEN-GAUTSCHI, G. (1976). Polyole in Pilzen: Analytik und systematische Verbreitung. – M. Sc. thesis, University of Zürich.
155. — (1980). Identität und systematische Verbreitung acyclischer Polyole von Eumyceten. – Ph. D. thesis, University of Zürich, Switzerland.
156. PFYFFER, G. E. & RAST, D. M. (1980 a). The polyol pattern of some fungi not hitherto investigated for sugar alcohols. – Exp. Mycol. 4: 160–170.
157. — (1980 b). Systematic distribution of ribitol in the fungi. – New Phytol. 85: 163–168.
158. PFYFFER, G. E. & RAST, D. M. – Unpublished data.
159. PLOUVIER, V. (1963). Distribution of aliphatic polyols and cyclitols. – In: SWAIN, T. (ed.). Chemical Plant Taxonomy: 313–336. Academic Press, London.
160. PRENTICE, N. & CUENDET, L. S. (1954). Composition of uredospores in wheat stem rust. – Nature (London) 174: 1151.

161. PRENTICE, N., CUENDET, L. S., GEDDES, W. F. & SMITH, F. (1959). The constitution of a glucomannan from wheat stem rust (*Puccinia graminis tritici*) uredospores. – *J. Am. Chem. Soc.* 81: 684–688.
162. PRUESS, L. M., PETERSON, W. H. & FRED, E. B. (1933). Isolation and identification of ergosterol and mannitol from *Aspergillus fischeri*. – *J. Biol. Chem.* 97: 483–489.
163. QUILLET, M. & LEGRAND, G. (1952). Sur le métabolisme glucidique des champignons supérieurs. I. Relation du mannitol avec l'activité du système oxidoréducteur. – *C. R. Acad. Sci. (Paris)* 234: 876–878.
164. RAGAN, M. A. & CHAPMAN, D. J. (1978). A biochemical phylogeny. – In: *A Biochemical Phylogeny of the Protists*: 211–232. Academic Press, New York.
165. RAISTRICK, H. & SMITH, G. (1933). Studies in the biochemistry of micro-organisms. XXXV. The metabolic products of *Byssochlamys fulva* Olliver & Smith. – *Biochem. J.* 27: 1814–1819.
166. RAST, D. (1965). Zur stoffwechselphysiologischen Bedeutung von Mannit und Trehalose in *Agaricus bisporus*. – *Planta* 64: 81–93.
167. REISENER, H. J., GOLDSCHMID, H. R., LEDINGHAM, G. A. & PERLIN, A. S. (1962). Formation of trehalose and polyols by wheat stem rust (*Puccinia graminis tritici*) uredospores. – *Can. J. Biochem. Physiol.* 40: 1248–1251.
168. RICHARDSON, D. H. S. (1985). The surface physiology of lichens with particular reference to carbohydrate transfer between the symbionts. – In: VICENTE, C., BROWN, D. H. & LEGAZ, M. E. (eds.). *Surface Physiology of Lichens*: 25–55. Editorial de la Universidad Complutense de Madrid.
169. ROBERTS, S. M. & MITCHELL, D. T. (1979). Carbohydrate composition and invertase activity in poplar leaf tissues infected by *Melampsora aecidioides* (D. C.) Schroeter. – *New Phytol.* 83: 499–508.
170. SALEWSKI, L., MIERSCH, J. & REINBOTHE, H. (1976). Polyol formation from glucose in flavingenic yeast *Candida guilliermondii* (Cast) Lang et G. – *Biochem. Physiol. Pflanz.* 170: 501–508.
171. SEELIGER, H. P. R. (1968). Serology as an aid to taxonomy. – In: AINSWORTH, G. C. & SUSSMAN, A. S. (eds.). *The Fungi*, vol. III: 597–624. Academic Press, New York.
172. SHAW, R. (1965). The occurrence of linolenic acid in fungi. – *Biochim. Biophys. Acta* 98: 230–237.
173. — (1966). The polyunsaturated fatty acids of microorganisms. – *Adv. Lipid Res.* 4: 107–174.
174. SIGLER, L. & CARMICHAEL, J. W. (1976). Taxonomy of *Malbranchea* and some other Hyphomycetes with arthroconidia. – *Mycotaxon* 4: 349–488.
175. SMILEY, K. L., CADMUS, M. C. & LIEPINS, P. (1967). Biosynthesis of D-mannitol from D-glucose by *Aspergillus candidus*. – *Biotechnol. Bioeng.* 9: 365–374.
176. SMITH, D., MUSCATINE, L. & LEWIS, D. (1969). Carbohydrate movement from autotrophs to heterotrophs in parasitic and mutualistic symbiosis. – *Biol. Rev.* 44: 17–90.
177. SMITH, P. M. (1976). *The Chemotaxonomy of Plants*. Edward Arnold, London.
178. SMITH, S. E. (1967). Carbohydrate translocation in orchid mycorrhizas. – *New Phytol.* 66: 371–378.
179. SNYDER, W. C. & TOUSSON, T. A. (1965). Current status of taxonomy in *Fusarium* species and their perfect stages. – *Phytopathology* 55: 833–837.
180. SO, M. L. & THROWER, L. B. (1976). The host-parasite relationship between *Vigna sesquipedalis* and *Uromyces appendiculatus*. III. Uptake of sugars by the parasite and respiration of the host-parasite combination. – *Phytopathol. Z.* 86: 302–309.
181. SPETH, J. L. & NIEDERPRUEM, D. J. (1976). Enzyme activities associated with arabinol and mannitol biosynthesis and catabolism in *Schizophyllum commune*. – *Arch. Microbiol.* 107: 81–86.

182. STAÜBLE, E. (1972). Vorkommen und Stoffwechsel leichtflüchtiger Substanzen von bekannter oder vermuteter physiologischer Wirksamkeit in *Agaricus bisporus*. – Ph. D. thesis, University of Zürich, Switzerland.
183. STEELE, S. D. (1972). Sugars and sugar alcohols in relation to life cycle phases of *Geotrichum candidum*. – Trans. Br. Mycol. Soc. 59: 502–506.
184. STODOLA, F. H. (1946). A note on *meso*-erythritol, a metabolic product of *Aspergillus terreus*. – J. Biol. Chem. 166: 79.
185. STOLK, A. C. & SAMSON, R. A. (1971). Studies on *Talaromyces* and related genera. I. *Hamigera* gen nov. and *Byssochlamys*. – Persoonia (Leiden) 6: 341–357.
186. STORCK, R. (1965). Nucleotide composition of nucleic acids of fungi. I. Ribonucleic acids. – J. Bacteriol. 90: 1260–1264.
187. — (1966). Nucleotide composition of nucleic acids of fungi. II. Deoxyribonucleic acids. – J. Bacteriol. 91: 227–230.
188. STORCK, R. & ALEXOPOULOS, C. J. (1970). Deoxyribonucleic acid of fungi. – Bacteriol. Rev. 34: 126–154.
189. STORCK, R., ALEXOPOULOS, C. J. & PHAFF, H. J. (1969). Nucleotide composition of deoxyribonucleic acid of some species of *Cryptococcus*, *Rhodotorula* and *Sporobolomyces*. – J. Bacteriol. 98: 1069–1072.
190. STRANDBERG, G. W. (1969). D-Mannitol metabolism by *Aspergillus candidus*. – J. Bacteriol. 97: 1305–1309.
191. STROBEL, G. A. & KOSUGE, T. (1965). Polyol metabolism in *Diplodia viticola* Desm. – Arch. Biochem. Biophys. 109: 622–626.
192. SUMI, M. (1928). Über die chemischen Bestandteile der Sporen in *Aspergillus oryzae*. – Biochem. Z. 195: 161–174.
193. SUSSMAN, A. S. & LINGAPPA, B. T. (1959). Role of trehalose in ascospores of *Neurospora tetrasperma*. – Science 130: 1343.
194. SUZUKI, T. & ONISHI, H. (1967). The production of xylitol and D-xylonic acid by yeasts. – Agric. Biol. Chem. (Jap.) 31: 1233–1236.
195. — (1968). Carbon balance of polyalcohol fermentation by *Pichia miso* and *Debaryomyces sake*. – J. Ferment. Technol. 46: 719–724.
196. — (1970). Aerobic dissimilation of D-ribulose by yeasts. – Antonie van Leeuwenhoek 36: 227–232.
197. TABUCHI, T. & HARA, S. (1973). Conversion of citrate fermentation to polyol fermentation in *Candida lipolytica*. – J. Agric. Chem. Soc. (Jap.) 47: 485–490.
198. TALBOT, P. H. B. (1971). Principles of Fungal Taxonomy. – The Macmillan Press, London.
199. TAMM, C. (1962). Über die Inhaltsstoffe von *Cordiceps capitata* (Holmsk.) Lk. – Planta Med. 10: 134–137.
200. TANI, T. & NAITO, N. (1960). D-Mannitol in the spores of *Puccinia coronata* and *Stereostromatum corticoides*. – Techn. Bull. Fac. Agric. Kagawa Univ. 12: 93–96.
201. TAYLOR, M. M. & STORCK, R. (1964). Uniqueness of bacterial ribosomes. – Proc. Natl. Acad. Sci. USA 52: 958–965.
202. THORNTON, J. D. & COOKE, R. C. (1974). Changes in respiration, chlorophyll content and soluble carbohydrates of detached cabbage cotyledons following infection with *Peronospora parasitica* (Pers. ex Fr.) Fr. – Physiol. Plant Pathol. 4: 117–125.
203. THROWER, L. B. & LEWIS, D. H. (1973). Uptake of sugars by *Epichloe typhina* (Pers. ex Fr.) Tul. in culture and from its host, *Agrostis stolonifera* L. – New Phytol. 72: 501–508.
204. TOUSTER, O. (1974). The metabolism of polyols. – In: SIPPLE, H. L. & McNUTT, K. W. (eds.). Sugars in Nutrition: 227–239. Academic Press, New York.
205. TOUSTER, O. & SHAW, D. R. D. (1962). Biochemistry of acyclic polyols. – Physiol. Rev. 42: 181–225.

206. TYLER, V. E. Jr. (1971). Chemotaxonomy in the Basidiomycetes. – In: PETERSEN, R. H. (ed.). Evolution in the Higher Basidiomycetes: 29–62. University of Tennessee Press, Knoxville.
207. TYLER, V. E. Jr., BENEDICT, R. G., BRADY, L. R. & ROBBERS, J. E. (1966). Occurrence of amanita toxins in American collections of deadly amanitas. – J. Pharm. Sci. 55: 590–593.
208. TYLER, V. E. Jr., BENEDICT, R. G. & STUNTZ, D. E. (1965). Chemotaxonomic significance of urea in the higher fungi. – Lloydia 28: 342–353.
209. TYLER, V. E. Jr. & SMITH, A. H. (1963). Chromatographic detection of Amanita toxins in *Galerina venenata*. – Mycologia 55: 358–359.
210. VAN DEN ENDE, H. (1978). Morphogenesis in the Phycomycetes. – In: SMITH, J. E. & BERRY, D. R. (eds.). The Filamentous Fungi, vol. III: 257–274. Edward Arnold, London.
211. VAN DER WALT, J. P. (1969). The genus *Syringospora* Quinquad emend. – Antonie van Leeuwenhoek 35 Supplement Yeast Symposium: A1.
212. VAN DER WALT, J. P. & HOPSU-HAVU, V. K. (1976). A colour reaction for the differentiation of ascomycetous and hemibasidiomycetous yeasts. – Antonie van Leeuwenhoek 42: 157–163.
213. VAN UDEN, N. & BUCKLEY, H. (1970). *Candida* Berkhoult. – In: LODDER, J. (ed.). The Yeasts: 893–1087. North-Holland Publishing Company, Amsterdam.
214. VAN UDEN, N. & VIDAL-LEIRIA, M. (1970). *Torulopsis* Berlese. – In: LODDER, J. (ed.). The Yeasts: 1235–1308. North-Holland Publishing Company, Amsterdam.
215. VINING, L. C. & TABER, W. A. (1964). Analysis of the endogenous sugars and polyols of *Claviceps purpurea* (Fr.) Tul. by chromatography on ion exchange resins. – Can. J. Microbiol. 10: 647–657.
216. VOGEL, H. J. (1965). Lysine biosynthesis and evolution. – In: BRYSON, V. & VOGEL, H. J. (eds.). Evolving Genes and Proteins: 25–40. Academic Press, New York.
217. VOLC, J., WURST, M. & MUSÍLEK, V. (1978). Soluble sugars in the mycelium of the basidiomycete *Oudemansiella mucida*. – Folia Microbiol. (Prague) 23: 448–452.
218. VON ARX, J. A. (1968) Pilzkunde. J. Cramer, Lehre.
219. VON LIPPmann, E. O. (1912). Über das Vorkommen von Trehalose, Vanillin und d-Sorbit. – Ber. Dtsch. Chem. Ges. 45: 3431–3434.
220. WAIN, R. L. & WILKINSON, E. H. (1946). Studies upon copper fungicides. IX. Investigations with the exudate from fungus spores. – Ann. Appl. Biol. 33: 401–405.
221. WANG, S.-Y. C. & LE TOURNEAU, D. (1971). Carbon sources, growth, sclerotium formation and carbohydrate composition of *Sclerotinia sclerotiorum*. – Arch. Mikrobiol. 80: 219–233.
222. WEETE, J. D. (1980). Lipid Biochemistry of Fungi and other Organisms. Plenum Press, New York.
223. WEIMBERG, R. (1962). Mode of formation of D-arabitol by *Saccharomyces mellis*. – Biochem. Biophys. Res. Commun. 8: 442–445.
224. WEINSTEIN, A. N. & SEGAL, S. (1967). Thin-layer chromatographic separation of hexitols and corresponding hexoses. – Anal. Biochem. 20: 558–561.
225. WIELAND, T. (1968). Poisonous principles of mushrooms in the genus *Amanita*. – Science 159: 946–952.
226. WRIGHT, J. R. & LE TOURNEAU, D. (1965). Utilization and production of carbohydrates by *Pyrenophaeta terrestris*. – Physiologia 18: 1044–1053.
227. WYNN, W. K., STAPLES, R. C., STROUSE, B. & GADUSEK, C. (1966). Physiology of uredospores during storage. – Contrib. Boyce Thompson Inst. 23: 229–242.
228. YAMASAKI, I. & SINOMURA, M. (1937). Über die Mannitbildung aus Glycerin durch Schimmelpilze der *Aspergillus glaucus*-Gruppe. – Biochem. Z. 291: 340–348.

229. YARROW, D. & MEYER, S. A. (1978). Proposal for amendment of the diagnosis of the genus *Candida* Berkhoult nom. cons. – Int. J. Syst. Bacteriol. 28: 611–615.
230. YUILL, J. L. (1948). Production of *i*-erythritol by *Aspergillus niger*. – Nature (London) 162: 652.
231. ZELLNER, J. (1907). Chemie der höheren Pilze. Wilhelm Engelmann, Leipzig.
232. — (1910). Zur Chemie der höheren Pilze. V. Mitteilung: über den Maisbrand (*Ustilago maydis* Tulasne). – Monatsh. Chem. 31: 617–634.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1986/1987

Band/Volume: [39](#)

Autor(en)/Author(s): Pfyffer Gaby E., Pfyffer B. U., Rast Dora M.

Artikel/Article: [The Polyol Pattern, Chemotaxonomy, and Phylogeny of the Fungi. 160-201](#)