

***Lauderlindsaya*, a new genus in the Verrucariales for  
*Sphaerulina chlorococca* (LEIGHTON) R. SANT.**

J. C. DAVID & D. L. HAWKSWORTH

CAB International Mycological Institute, Ferry Lane, Kew, Richmond, Surrey,  
TW9 3AF, UK

DAVID, J. C. & D. L. HAWKSWORTH (1989). *Lauderlindsaya*, a new genus in the Verrucariales for *Sphaerulina chlorococca* (LEIGHTON) R. SANT. – SYDOWIA 41: 108–121.

The new generic name *Lauderlindsaya* is introduced for the lichenicolous parasite of *Normandina pulchella* (BORRER) NYL., previously known as *Sphaerulina chlorococca* (LEIGHTON) R. SANT. The biology and systematics of the new genus are discussed, and it is referred to the Verrucariales. It is compared with other multisepate-spored members of that order.

The presence of perithecia associated with the squamules of *Normandina pulchella* (BORRER) NYL. has been known since BORRER (1829: tab. 2602, fig. 1) described it as *Verrucaria pulchella* BORRER but the precise nature of that association has been a source of considerable problems, both nomenclaturally and systematically. The nomenclatural problems were resolved (HENSSSEN, 1976: 129), when *N. pulchella* was lectotypified on a specimen of BORRER's from Sussex, preserved in his herbarium at K, which lacked the pyrenomycete.

The systematic confusion derives from the fact that *N. pulchella* is now known to be a sterile lichen of uncertain affinity. The presence of the parasite gives the impression that *N. pulchella* is fertile, as for the most part the parasite does no apparent damage. The illustration in BORRER (1829) of *N. pulchella* includes one specimen from Sussex which lacks the parasite (the lectotype; see above), and one from Bantry in Eire which clearly is infected. His text also includes a reference to the perithecia of the parasite as an integral part of his description of *Verrucaria pulchella*. This was followed by subsequent lichenologists, so that HOOKER (1833) treated it as a species of *Endocarpon* HEDW. DELISE, in DESMAZIÈRES' exsiccata, Plantes Cryptogames du Nord de la France, fasc. 23, no. 1144 (1841), described a new genus, *Lenormandia* based on material of *N. pulchella*, but his name must be rejected as a later homonym of *Lenormandia* DELISE, *L. jungermanniae* DELISE, was considered identical to *N. pulchella* by KÖRBER (1855: 101). NYLANDER (1855: 191) described

the genus *Normandina*, again including descriptions of the parasite's ascomata as if they were the perithecia of the lichen. However TULASNE (1852: 128), commenting on DELISE's description of the genus *Lenormandia*, said that the "fruits" which had been illustrated as part of the lichen in BORRER (1829) and by LEIGHTON (1851) were those of a parasite to which he gave the name *Sphaeria borrieri* TULASNE. KÖRBER (1855) noted the similarity between *Lenormandia* and TULASNE's parasitic fungus. LINDSAY (1869: 54) states, in his notes on *Normandina jungermanniae* (DELISE) NYL. [i. e. *N. pulchella*], that the spores found in the thallus belong to the parasitic fungus *Sphaeria borrieri*, thus demonstrating that he also was aware of the allochthonous nature of the perithecia described for *N. pulchella*.

Unfortunately the contributions of TULASNE and LINDSAY went unnoticed by later lichenologists (LEIGHTON, 1879; VAINIO, 1890; SMITH, 1926; FINK, 1935; SWINSCOW, 1963) who included *Normandina* as a member of the Verrucariaceae, on the assumption that the presence of the parasite represented its ascomata. VAINIO (1921: 231), however, treated *Normandina* pro parte as a synonym of *Coriscium* VAINIO, as a sterile lichen but did include a reference to the description of "apothecia" by LEIGHTON (1851, 1879). He left the actual position of the lichen as uncertain.

LEIGHTON (1879: 484) described *Verrucaria chlorococca* with exactly the same features as the "fertile *Normandina*" except in the thallus which he gave as, "green, thickish, granulate, somewhat areolato-diffract". This has subsequently been referred to informally as the "free-living form". SMITH (1926) transferred *V. chlorococca* to *Arthopyrenia* MASSAL., emending LEIGHTON's description to include, "paraphyses slender, mucilaginous, disappearing", as opposed to LEIGHTON's, "paraphyses none".

SWINSCOW (1963) realised that LEIGHTON's *V. chlorococca* was the same as the fungus on *N. pulchella* but continued to regard the fungus as being the fertile state of *Normandina*. The problem was not considered again until HENSSEN (1976: 128) confirmed the view that the squamulose part of the association is a sterile lichen. Although she drew comparisons with *Coriscium*, she concluded, due to the absence of dolipore septa in the hyphae, that *Normandina* could not be a basidiomycete. The identity of the parasite was dealt with by SANTESSON (in HENSSEN 1976: 129), who considered it *faute de mieux*, "to be a species of *Sphaerulina* (a somewhat heterogenous assembly . . .)", and transferred it to that genus as *Sphaerulina chlorococca* (LEIGHTON) R. SANT., LEIGHTON's epithet being lectotypified by LARBALESTIER's specimen in BM, which is *S. chlorococca* not associated with *N. pulchella*.

## Materials and Methods

Specimens examined were from BM, E, UPS and IMI. The specimens were sectioned using a freezing microtome and mounted in lactophenol or Lugol's iodine. Squash preparations of the perithecia were made, mounted in 10% KOH or Lugol's iodine. Slides were examined using a Zeiss Universal microscope with Nomarski and phase optics.

## Biology

The rediscovery of "free-living" material prompted an investigation to determine, (a) whether *Sphaerulina chlorococca* really was free-living in association with algae, and, (b) if this morph represented a separate new species. Examination of sections of material from bark and mosses showed the ascomata surrounded by algae, with brownish hyphae spreading out from the ascomata into the algal mass. Sections of the "*Sphaerulina*" immersed in the thallus of *Normandina* show that the ascomata protrude on the underside of the squamules, but are not attached at the base to the substrate on which the *Normandina* is growing. The algae of *Normandina* and those found associated with the "free-living" ascomata were chlorococcalean. Initially, the presence of the ascomata have no apparent effect on the algae or the thallus of *Normandina*, but later on the thallus can become bleached or even degenerate to a powdery mass. The ascomata and ascospore sizes were recorded and the peridial structure was examined in both the "free-living" and the "parasitic" collections in order to determine whether there were any differences between them (Tab. 1, Fig. 1).

Tab. 1. – Range and average (with standard deviations) in length and width of spores of *Lauderlindsaya borrieri* on different substrata

substrate	length $\mu\text{m}$		width $\mu\text{m}$	
	range	average	range	average
<i>Sambucus</i> (COPPINS 4009)	31.1 – 37.8	33.6 $\pm$ 2.2	5.9 – 8.4	7.0 $\pm$ 0.9
<i>Quercus</i> (COPPINS 11223)	26.9 – 38.6	31.9 $\pm$ 3.2	6.7 – 8.4	7.4 $\pm$ 0.7
with <i>Parmeliella</i> (COPPINS 4165)	26.9 – 36.1	32.1 $\pm$ 2.7	6.7 – 8.4	7.6 $\pm$ 0.7
<i>Salix</i> (COPPINS 2574)	31.5 – 38.4	35.3 $\pm$ 2.2	6.9 – 8.8	8.0 $\pm$ 0.5
Moss (indet.) (COPPINS 9905)	26.0 – 31.9	29.3 $\pm$ 1.8	6.7 – 8.4	7.6 $\pm$ 0.4
<i>Normandina</i> (a) (IMI 196191)	35.3 – 41.6	38.5 $\pm$ 2.3	6.3 – 8.2	7.5 $\pm$ 0.6
<i>Normandina</i> (b) (IMI 296945)	27.7 – 42.0	34.7 $\pm$ 4.0	6.7 – 9.2	7.8 $\pm$ 0.9




Fig. 1: Ascospores. – A. *Lauderlindsaya borrieri*, from perithecia on *Normandina pulchella*. – B. From perithecia on *Salix*.



The "free-living" material of *Sphaerulina chlorococca* proved to be almost identical to that found on *Normandina* thalli (Tab. 1). The only noticeable difference was in the ascomatal wall; when "free-living" the perithecia are more heavily thickened than when "parasitic" (see Plate 2, b & c), to the extent that the ascomatal wall of "free-living" material shows a distinct tendency to break up into plates (in squash preparations) (Plate 3, a). This is probably a phenotypic response to their different ecologies. It seems unlikely that the "free-living" type represents an independently lichenized morph of *S. chlorococca*, and it is probably best to consider it as weakly pathogenic or commensalistic (HAWKSWORTH, 1988). The similarity of the host algae suggests that the relationship is a specific one; indeed the difference between the attachment of the ascomata of *S. chlorococca* suggests that the "free-living" material is not a result of the degeneration of *Normandina* thalli.

### Taxonomy

Examination of both "free-living" and "parasitic" material confirmed the suspicion of HAWKSWORTH (1983: 15) that the fungus did not belong to *Sphaerulina* SACC. It was therefore necessary to determine where its affinities lay. The asci have such delicate walls that there were difficulties in examining them in detail. The walls of the asci were seen to have more than one layer (Plate 1, fig. d) and are similar to *Dermatocarpon* ESCHW., as SANTESSON (in HENSSEN 1976: 129) noted: "The ascus type is difficult to recognize on account of the very easily bursting walls . . . have a similar structure to the bitunicate ascus of *Dermatocarpon*". JANEX-FAVRE (1970) and ERIKSSON (1981: 55) however, state the *Dermatocarpon* asci have a distinct ocular chamber, which *S. chlorococca* does not. *S. chlorococca* was compared with several similar genera, using data from direct observations and available literature, and was found to have the most characters in common with the Verrucariales, a placement supported by its ascus features. That it is not a *Sphaerulina*, a genus in the Dothideales, is clear as it has prominent periphyses (Plate 1, figs b & c) and a different peridial structure (Plate 2, figs b & c); the ascomata are also much larger than in any accepted *Sphaerulina* species, and the asci lack the apical nasse seen in *Sphaerulina*.

The Verrucariales contains almost entirely lichenized taxa, but also a few lichenicolous fungi such as *Muellerella* species and lichenicolous lichens, particularly species of *Verrucaria* (ZEHETLEITNER, 1978; MCCARTHY, 1988). Whilst the Verrucariales have mainly non-septate spores, all the basic spore septation types can be found; trans-septate genera are *Thelidium* MASSAL., *Macentina* VÉZDA, *Placidiopsis* BELTR. and *Heterocarpon* MÜLL. ARG. and muriform ones also occur, in *Agonimia* ZAHLBR., *Polyblastia* MASSAL.,




Plate 1: *Lauderlindsaya borneri*. – A. Ascoma ( $\times 250$ ). – B. Ostiole of ascoma showing arrangement of periphyses ( $\times 650$ ). – C. Detail of periphyses ( $\times 650$ ). – D. Asci and ascospores ( $\times 650$ ).

and *Leucocarpia* VÉZDA. Compared with the septate genera, *Macentina* differs from *S. chlorococca* in having small pellucid perithecia. The squamulose/foliose genera *Placidiopsis* and *Heterocarpon* usually have 1-septate ascospores but may have up to 3-septate spores. Thallus morphology, insofar as the perithecia do not project below the lower surface of the thalli, precludes further consideration here. The differentiation between *Thelidium* and *S. chlorococca* is more problematical. ZSCHACKÉ's (1934) account of *Thelidium* includes a number of taxa which occur on substrata considered unusual for *Thelidium* and also with a degree of spore septation which similarly is not typical of that genus. A number of these taxa have been examined in order to be able to more clearly define the separation between *Thelidium* and *S. chlorococca*, and some of them have indeed proved to be identical to the latter as anticipated. Further work is needed to determine the nature of these taxa. A. ORANGE (pers. comm.) has noted that the ascus dehiscence in *Thelidium* s. str. is quite different, in that there is an extrusion of the endotunica after the release of spores. ERIKSSON (1981: 158) observed that the type species of *Thelidium*, *T. amylaceum* MASSAL. has a "distinct, low, broad ocular chamber" which *S. chlorococca* does not. *Pocsia* VÉZDA, a genus of foliicolous lichens currently treated as a member of the Trichotheliaceae (ERIKSSON & HAWKSWORTH, 1987), was suggested by VÉZDA (1975) to be possibly related to the Verrucariales, particularly through the absence of paraphyses, the presence of periphyses and the perithecium structure; the asci also appear to be Verrucarialean. *Pocsia* differs from *S. chlorococca* in having perithecial walls which are not black in colour and are covered by a thalline layer.

Thus it is clear that a new genus is necessary to accommodate *S. chlorococca*.

***Lauderlindsaya* J. C. DAVID & D. HAWKSW., gen. nov.**

Genus lichenicola aut algicola, ad Verrucariales pertinens. Ascomata perithecia subglobosa aut conoideo-globosa, nigra, pseudoparenchymatica, quaeque ostiolo punctiformi terminali induta. Paraphyses non manifestae, et periphyses faciles visae, simplices aut ramosae. Centrum iodo fusco-rubescens. Asci clavati ad lati-clavati, muris tenuibus sed ad apicem incrassatis, 8-spori. Ascosporae ellipsoideo-fusiformes, plerumque 7-septatae, ad septa strictae, primum decolores, dein pallide fuscae, laeves.

Species holotypica, adhuc unica, est *Lauderlindsaya borrieri* (TULASNE) J. C. DAVID & D. HAWKSW. (syn. *Sphaeria borrieri* TULASNE).

Lichenicolous or algicolous ascomycete, belonging to the Verrucariales. — Ascomata perithecioid subglobose or conical-globose, black, pseudoparenchymatous with a terminal punctiform ostiole. — Paraphyses not seen, periphyses well developed, branched or unbranched. Centrum turning red-brown with iodine. — Asci clavate




Plate 2: *Lauderlindsaya borrieri*. – A. Ascospores ( $\times 650$ ). – B. Detail of ascomatal wall of *L. borrieri* on *Normandina pulchella* thallus. – C. Detail of ascomatal wall in “free-living form” ( $\times 650$ ).



or broadly clavate, with thin walls but thickened at the apices, 8-spored. – Ascospores ellipsoid-fusiform, usually 7-septate (immature spores 3-5-septate), constricted at the septa, at first colourless then becoming pale brown (with age), smooth walled.

This genus is named in honour of WILLIAM LAUDER LINDSAY (1829–1880) one of the foremost pioneers of studies of lichenicolous fungi.

The generic name *Normandinomyces* CIFERRI & TOMASELLI (1953: 30), was also considered as a possible earlier name for *Lauderlindsaya*. The former name can be excluded as nomenclaturally illegitimate; being based on the same type as *Normandina* it is automatically typified by the lectotype of BORRER's name, thus rendering it superfluous under Article 63 of the Code.

***Lauderlindsaya borrieri* (L. R. TULASNE) J. C. DAVID & D. HAWKSW., comb nov.**

Bas.: *Sphaeria borrieri* L. R. TULASNE, Ann. Sci. nat., Bot., sér. 3, 17: 128, 1852.

Syn.: *Verrucaria chlorococca* LEIGHTON, Lich. Fl. Br., edn 3: 484, 1879.

*Arthopyrenia chlorococca* (LEIGHTON) A. L. SMITH, Br. Lich., edn II, 2: 361, 1926.

*Thelidium chlorococcum* (LEIGHTON) KESSLER, Rabenh. Krypt. Fl. edn 2: 9, 1(2): 191, 1938.

*Sphaerulina chlorococca* (LEIGHTON) R. SANT., in HENSSEN, apud BROWN et al., Lichenology: Progr. Probl.: 129, 1976.

= *Thelidium erichsenii* KESSLER, Revue mycol., 1: 179, 1936.

= *Polyblastia armericola* W. WATSON, J. Bot., Lond., 77: 42, 1939.

Typification. – TULASNE (l. c.) cites BORRER's illustration (2602 fig. 1) as showing *Sphaeria borrieri*, the parasite of *Normandina*; the specimens used for this illustration are kept in BORRER's herbarium at Kew. One of the specimens was used to lectotypify *Normandina pulchella* (HENSSEN, 1976: 129). The other two are from Ireland, of which one collected by Miss HUTCHINS from Bantry was the other specimen used for the drawings, and is the top right specimen figured. The other Irish specimen was collected by Dr. T. TAYLOR and was used for LEIGHTON's (1851) plate 3 fig. 1. Since TULASNE in his protologue cites BORRER's illustration, the top right-hand specimen on the sheet must be considered as the lectotype material of the name *Sphaeria borrieri*. VOUAUX (1913) considered TULASNE's name to be "trop hypothétique", but whilst TULASNE did not give a description as such, it was clear from his protologue what he intended when he said, "simul et Sphaeriam (*S. Borrieri* Nob.) hospitem sparsim suis in penetralibus immersam exhiberet", and in his citation of relevant literature.

Illustrations. – *Lauderlindsaya* has been illustrated by BORRER (1829: tab. 2602 fig. 1), LEIGHTON (1851: Plate 3 fig. 1), SMITH




Plate 3: *Lauderlindsaya borrieri*. – A. Squash preparation of ascematal wall of "free-living form", under phase contrast ( $\times 250$ ). – B. Lectotype of *Sphaeria borrieri* TULASNE, in Herb. K.



(1926: Plate 40), SWINSCOW (1963: 168, fig. 1) and HENSSEN (1976: Plates XVII, XVIII & XIX).

**Description.** – Fungus occurring on *Normandina pulchella* or on free-living chlorococcalean algae, not forming an independent thallus. – Ascomata perithecioid, solitary, immersed in the host thallus, protruding through the underside of host squamules, or not on a distinct thallus and surrounded by an algal mass; black, matt, with a distinct apical ostiole, entire, globose at the base, becoming conical at the apex; peridium *textura angularis* with fairly elongated cells composed of 10–15 cell layers, 30–45(–55)  $\mu\text{m}$  thick, the outer cells having densely pigmented walls, the inner layers  $\pm$  colourless, (160–)200–300(–350)  $\times$  (150–)170–250(–330)  $\mu\text{m}$  diam. – Centrum I+ red–brown. – Hamathecium lacking interascal filaments, periphyses well developed at the ostiole, septate, branched or unbranched. – Asci clavate to broadly clavate, short stalked, with delicate walls, ascus apex lacking any apparatus, but clearly showing the 2-layered nature of the walls, I–; (15–)17–23  $\times$  85–95(–115)  $\mu\text{m}$ , 8-spored. – Ascospores elongate-ellipsoid, ends pointed, (5–)7-trans-septate, colourless, becoming brown with age (usually after discharge), initially not constricted at the septa but becoming so with age, smooth-walled, 25–35(–45)  $\times$  6–9  $\mu\text{m}$ , after discharge the spores tend to remain clustered together. – Anamorph not observed.

**Distribution.** – In the British Isles the fungus occurs along the western coasts, and is also present in continental Europe (DIEDERICH, 1986), in Chile, W. Australia, the Philippines and Madeira. This distribution tends to indicate that *Lauderlindsaya* will be found where its host occurs.

The distribution of *Normandina pulchella* is temperate – subtropical to tropical, mainly occurring in damp woodlands/rainforests. It is recorded from all five continents. North America (CULBERSON & HALE, 1966); South America (eg. DEGELIUS, 1935; FIGUEIRAS, 1986; XAVIER FILHO & TOLEDO RIZZINI, 1976); Central America (PLUNTSKE, 1984); Africa (eg. ALMBORN, 1966; SWINSCOW & KROG, 1988); Australia (FILSON & ROGERS, 1979); New Zealand (GALLOWAY, 1985); Papua New Guinea (SHAW, 1984); Japan (YOSHIMURA, 1974); USSR (OXNER, 1956); Europe (DEGELIUS, 1935).

Many records of the host in floras tend to include a description of the parasite (usually as the fertile perithecia), but this may not always mean that they had recorded it with the parasite themselves.

**Specimens examined.** – EIRE: Co. Cork, on a mountain near Bantry, on stems of a heath, Miss HUTCHINS (K-BORRER – Lectotype of *Sphaeria borrieri* TUL.). [Plate 3, b]. – UK: Buckinghamshire: Stokenchurch, on mossy bark of Ash (*Fraxinus excelsior*), ii. 1876, C. LARBALESTIER, (BM-Lectotype of *Verrucaria chlorococca* LEIGHTON). – V. C. 1, Cornwall: Lizard, on *Armeria maritima*, 1928, H. H. KNIGHT & D. A.

JONES (BM-Holotype of *Polyblastia armericola* W. WATSON). – V. C. 3, Devon: Newton Abbott, Bradley Manor, on *Normandina pulchella*, 15. viii. 1975, J. FILDES (IMI 196191). – Holne Chase below the Hotel, on *N. pulchella* on *Salix* branch, 3. viii. 1985, D. L. HAWKSWORTH 5519 (IMI 296945). – V. C. 68, Northumberland: Easington, Spindlestone Haugh, on *Sambucus* bark, 9. iv. 1979, B. J. COPPINS 4009 (E). – V. C. 96, Easternness: Glen Cannich, by R. Cannich south west of Muchrachd, on *Populus*, 22. vi. 1976, B. J. COPPINS 3226 (E). – V. C. 97, Westernness: north side of Loch Sunart, c. 4 km, east of Salen, Resipole ravine, on *N. pulchella*, with *Capronia* sp., 10. iii. 1983, B. J. COPPINS & P. M. JØRGENSEN 9405 (E). – V. C. 98, Argyll: Seil, Ballachuan, wood south of Port Mor, on *Salix*, 30. vi. 1976, B. J. COPPINS 3226 (E). – V. C. 103, Mid-Ebudes: Coll, west side of Loch Fada, on a horizontal branch of *Salix*, iv. 1983, B. J. COPPINS 9637 (E); Coll, Arinagour, The Lodge, on *Acer* with, but not on *N. pulchella*, 16. iv. 1983, B. J. COPPINS 9666 (E). – V. C. 104, Skye: Dunvegan Castle woods, over (not in) thallus of *Parmeliella atlantica* on *Fraxinus*, 26. vi. 1979, B. J. COPPINS 4165 (E). – V. C. 105, West Ross: Loch Maree, Talladale oakwood, on *Quercus*, 19. v. 1984, B. J. COPPINS et al. 11223 (E). – V. C. 108, West Sutherland: Bettyhill, Invernaver NNR, over mosses amongst *Dryas*, 22. viii. 1983, B. J. COPPINS et al. 9823 (E) & B. J. COPPINS et al. 9905 (E). – EIRE: V. C. H 1 or 2, Co. Kerry: sine loc., on *N. pulchella*, Dr. MOORE (E). – GERMANY: Schleswig-Holstein, near Ackerum on Föhr island (Nordfriesische Inseln), on bark of *Populus tremula*, 22. v. 1929, C. F. E. ERICHSEN (HBG-Lectotype of *Thelidium erichsenii* KESSLER). – CHILE: Prov. Chiloé: Isla Chiloé, Peninsular Lacui, Punta Corona, on *N. pulchella*, on *Berberis buxifolia* in a dense forest near the seashore, 20. x. 1940, R. SANTESSON No. 4078 (UPS). – AUSTRALIA: W. Australia: 15 km W. of Pemberton, Beedelup National Park, at Beedelup Falls, 34° 26' S, 115° 53' E, on *N. pulchella* on bark of *Eucalyptus* sp., 10. x. 1983, L. TIBELL No. 14001 (UPS). – PHILIPPINES: Luzon Island, Benguet subprovince, Baguio "and vicinity", on *N. pulchella* growing with isotype specimen of *Anaptychia propagulifera* (PERS.) VAIN. var. *propagulifera* VAIN., in BM (UPS). – MADEIRA: Concelho da Santana: Ribiero Frio, by the "levada" E. of Ribiero Frio, alt. 850–870 m, on *N. pulchella*, 6. v. 1978, R. SANTESSON No. 28096 (UPS).

### Acknowledgments

We are grateful to the curators of the herbaria E, K, BM and UPS for the loan of specimens; to Professor R. SANTESSON, Dr B. J. COPPINS and Mr A. ORANGE for helpful comments; to Dr D. W. MINTER for help with the Latin diagnosis; to Mrs A. ANSELL for technical assistance in the preparation of specimens; and to Miss G. GODWIN for help with photography.

### References

- ALMBORN, O. (1966). Revision of some lichen genera in Southern Africa I. – *Botaniska Notiser* 119: 70–112.
- BORRER, W. (1829). *English Botany. Supplement. Vol. 1.* – London: Sowerby.
- CIFERRI, R. & R. TOMASELLI. (1953). Saggio di una sistematica micolichenologica. – *Atti dell'Istituto Botanico e Laboratorio Crittogamico di Pavia ser. 5*, 10: 25–84.
- CULBERSON, W. L. & M. E. HALE. (1966). The range of *Normandina pulchella* in North America. – *Bryologist* 69: 365–367.
- DEGELIUS, G. (1935). Das ozeanische Element der Strauch- und Laubflechten-Flora von Skandinavien. – *Acta Phytogeographica Suecica* 7: 1–411.
- DIEDERICH, P. (1986). Lichenicolous fungi from the Grand Duchy of Luxembourg. – *Lejeunia*, n. s. 119: 1–26.

- ERIKSSON, O. (1981). The families of bitunicate ascomycetes. – Opera botanica 60: 1–209.
- & D. L. HAWKSWORTH (1987). Outline of the ascomycetes – 1987. – Systema Ascomycetum 6: 259–337.
- FIGUEIRAS, M. L. (1986). Censo do macroliquenes Venezolanos dos los estados Falcon, Lara, Merida, Tachira y Trujillo. – Universidad de los Andes, Facultad de Farmacia.
- FILSON, R. B. & R. W. ROGERS. (1979). Lichens of South Australia. – South Australia: Government Printer.
- FINK, B. (1935). The Lichen Flora of the United States. – Ann Arbor: Michigan University Press.
- GALLOWAY, D. J. (1985). Flora of New Zealand Lichens. – Wellington: Government Printer.
- HAWKSWORTH, D. L. (1983). A key to the lichen-forming, parasitic, parasymbiotic and saprophytic fungi occurring on lichens in the British Isles. – Lichenologist 15: 1–44.
- (1988). The variety of fungal-algal symbioses, their evolutionary significance, and the nature of lichens. – Botanical Journal of the Linnean Society 96: 3–20.
- HENSSEN, A. (1976). Studies in the developmental morphology of lichenized ascomycetes. In Lichenology: Progress and Problems (D. H. BROWN, D. L. HAWKSWORTH & R. H. BAILEY eds): 107–138. – London: Academic Press.
- HOOKE, W. J. (1933). The English Flora of J. E. SMITH. Vol. 5. – London: Longman, Brown, Green & Longmans.
- JANEX-FAVRE, M. (1970). Recherches sur l'ontogénie, l'organisation et les asques de quelque Pyrénéen lichens. – Revue Bryologique et Lichénologique 37: 421–649.
- KÖRBER, W. (1855). Systema lichenum Germaniae. – Breslau: Trewendt & Granier.
- LEIGHTON, W. A. (1851). The British Species of Angiocarpous Lichens. – London: Ray Society.
- (1879). The Lichen-Flora of Great Britain, Ireland and the Channel Islands. 3rd ed. – Shrewsbury.
- LINDSAY, W. L. (1869). Enumeration of micro-lichens parasitic on other lichens. – Quarterly Journal of Microscopical Science, n. s., 9: 49–57.
- MCCARTHY, P. M. (1988). New and interesting species of *Verrucaria* II. – Lichenologist 20: 245–252.
- NYLANDER, W. (1955). Essai d'une nouvelle classification des lichens; Second mémoire. – Mémoires de la Société Nationale des Sciences Naturelles de Cherbourg 3: [161]–202.
- OXNER, A. M. (1956). Flora Lishainikov Ukraini. – Kiev: Vidavmitsvo akademii nauk Ukrain'skoi RSR.
- PLUNTKKE, M. (1984). Die Flechtenflora Kubas – Bibliographie. – Halle (Saale): Universitäts- und Landesbibliothek: Sachsen-Anhalt. (Terrestrische Oekologie: Sonderheft 4).
- SHAW, D. E. (1984). Microorganisms in Papua New Guinea. – Research Bulletin No. 33 of the Department of Primary Industry, Port Moresby, Papua New Guinea.
- SMITH, A. L. (1926). A Monograph of the British Lichens, 2<sup>nd</sup> edn. Vol. 2. – London: British Museum (Natural History).
- SWINSCOW, T. D. V. (1963). Pyrenocarpous lichens: 5. Fruiting *Normandina pulchella* (BORR.) NYL. – a cause for confusion. – Lichenologist 2: 167–169.
- & H. KROG. (1988). Macrolichens of East Africa. – London: British Museum (Natural History).
- TULASNE, L.-R. (1852). Mémoire pour servir à l'histoire organographique et physiologique des lichens . . . – Ann. Sci. nat., Bot., sér. 3., 17: [5]–128, 153–249.
- VAINIO, E. A. (1890). Étude sur la classification naturelle et la morphologie des lichens du Brésil. – Acta Societatis pro Fauna et Flora fennica 7: 1–256.

- (1921). Lichenographica Fennica I. – Acta Societatis pro Fauna et Flora Fennica 49(2): 1–274.
- VÉZDA, A. (1975). Foliicole Flechten aus Tanzania (Ost-Afrika). – Folia Geobotanica et Phytotaxonomica 10: 383–432.
- VOUAUX, L. (1913). Synopsis des Champignons parasites de Lichens. – Bulletin de la société Mycologique de France 29: 33–128.
- XAVIER FILHO, L. & C. TOLEDO RIZZINI. (1976). Manual de Liquenologia Brasileiro. – Recife: Universidade de Pernambuco.
- YOSHIMURA, I. (1974). Lichen Flora of Japan. – Osaka: Hoikusha Publ. Co.
- ZEHETLEITNER, G. (1978). Über einige parasitische Arten der Flechtengattung *Verrucaria*. – Nova Hedwigia 29: 683–734.
- ZSCHACKE, H. (1934). Verrucariaceae and Dermatocarpaceae. In Rabenhorst's Kryptogamen-Flora 9 1/1. – Leipzig: Akademische Verlagsgesellschaft.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 1989

Band/Volume: [41](#)

Autor(en)/Author(s): David J. C., Hawksworth David Leslie

Artikel/Article: [Lauderlindsaya, a new genus in the Verrucariales for Sphaerulina chlorococca \(LEIGHTON\) R. SANT. 108-121](#)