

***Phyllosticta* species on orchids (Orchidaceae), introducing  
*Phyllosticta speewahensis* sp. nov. (Phyllostictaceae,  
Ascomycota) from northern Australia**

Roger Graham Shivas<sup>1</sup>,\*, Yu Pei Tan<sup>1</sup> & Kathy R. E. Grice<sup>2</sup>

<sup>1</sup> Plant Pathology Herbarium, Ecosciences Precinct, GPO Box 267, Brisbane, Qld 4001, Australia.

<sup>2</sup> Agri-Science Queensland, PO Box 1054, Mareeba 4880, Queensland, Australia.

Shivas R. G., Tan Y. P. & K. R. E. Grice (2013) *Phyllosticta* species on orchids (Orchidaceae), introducing *Phyllosticta speewahensis* sp. nov. (Phyllostictaceae, Ascomycota) from northern Australia. – *Sydowia* 65 (1): 139–146.

Several species of *Phyllosticta* (syn. *Guignardia*) have been described from orchids worldwide. A new species, *Phyllosticta speewahensis*, is proposed for a specimen isolated from leaf spots on a hybrid *Vanda* orchid in northern Queensland, Australia. Phylogenetic analysis of the nrDNA internal transcribed spacer region (ITS) and partial translation elongation factor 1-alpha (TEF1) gene sequences showed that *P. speewahensis* is most closely related to *P. hostae*. The likelihood that orchids harbour further cryptic species of endophytic and pathogenic *Phyllosticta* species is discussed.

Keywords: cryptic species, *Phyllosticta*, molecular phylogeny, taxonomy, Orchidaceae.

*Phyllosticta* (syn. *Guignardia*) species are widely distributed as plant pathogens, endophytes and saprobes on a range of hosts (van der Aa 1973; van der Aa & Vanev 2002; Wikee *et al.* 2011, 2013). Several species of *Phyllosticta* were originally described from orchids (Orchidaceae) based on their distinct morphology and host association (Tab. 1). Van der Aa & Vanev (2002) revised *Phyllosticta* based on the literature and examination of type specimens, accepting only *P. aplectri* and *P. capitalensis* amongst the taxa described from orchids.

Recently, several new species of *Phyllosticta* have been discovered from a wide range of plants, using phylogenetic analysis of DNA gene sequences, supported by morphology and host association (Wulanderi *et al.* 2009, Wang *et al.* 2012, Wong *et al.* 2012, Su & Cai 2012, Shivas *et al.* 2012, Crous 2012), including *P. bifrenariae* O. L. Pereira, Glienke & Crous from the orchid *Bifrenaria harrisoniae* in Brazil (Glienke *et al.* 2011). In 2012, a cultivated orchid (*Vanda* hybrid ‘Waxy Blue’) in northern Queensland was found to have leaf spots colonized by a *Phyllosticta* sp. Morphology and molecular analysis of

---

\* e-mail: roger.shivas@daff.qld.gov.au

sequence data could not match this taxon with any known species, and thus, it is introduced as new to science in this paper.

**Tab. 1.** Species originally described as *Phyllosticta* from orchids (Orchidaceae).

Species, authority and year of publication a	Substrate and/or orchid host b	Country of type	Reference
<i>P. aplectri</i> Ellis & Everh. 1894	<i>Aplectrum hyemale</i>	USA	van der Aa (1973)
<i>P. capitalensis</i> Henn. 1908	leaves of <i>Stanhopea</i> sp.	Brazil	Hennings (1908)
<i>P. cymbidii</i> Sawada 1943 <sup>a</sup>	leaves of <i>Cymbidium aloifolium</i>	Taiwan	van der Aa (1973)
<i>P. himantoglossi</i> Petr. 1959 <sup>b</sup>	dying leaves of <i>Himantoglossum hircinum</i>	Austria	Petrak (1959)
<i>P. imbricata</i> B. R. D. Yadav & V. G. Rao 1981	living leaves of <i>Pholidota imbricata</i>	India	van der Aa & Vanev (2002)
<i>P. odontoglossi</i> Verplancke & Claessens 1934 <sup>c</sup>	living leaves of <i>Odontoglossum</i>	Belgium	van der Aa & Vanev (2002)
<i>P. phalenopsisidis</i> Marchal & Verplancke 1926 <sup>d</sup>	living leaves of <i>Dendrobium phalaenopsis</i>	Belgium	van der Aa & Vanev (2002)
<i>P. phalenopsisidis</i> var. <i>vanillae</i> Verplancke & Claessens 1934 <sup>d</sup>	living leaves of <i>Vanilla lujia</i>	Belgium	van der Aa & Vanev (2002)

<sup>a</sup> an invalid name (van der Aa 1973)

<sup>b</sup> probably a *Phoma* (van der Aa & Vanev 2002)

<sup>c</sup> probably an *Asteromella* or *Phoma* (van der Aa & Vanev 2002)

<sup>d</sup> excluded from *Phyllosticta* (van der Aa & Vanev 2002)

## Materials and methods

### Isolates and morphology

Isolates were maintained on 2% potato dextrose agar (PDA) and incubated for 14 d under 12 h ultraviolet light / 12 h dark cycle following Wulanderi *et al.* (2009). Fungal structures were mounted in lactic acid on glass slides and examined by Nomarski interference contrast microscopy. Conidial measurements are given as follows: (minimum) mean – standard deviation × mean + standard deviation (maximum) rounded to 0.5 µm for 30 conidia. Other morphological measurements reflect the range of dimensions observed. Colony colours were determined by Rayner's (1970) colour charts. A representative isolate is deposited as ex-type culture in BRIP 58044.

### DNA isolation, amplification and phylogenetic analysis


DNA was extracted from fungal mycelium with the Gentra Puregene kit (Qiagen, Melbourne, Australia) according to the manufacturer's instructions. PCR amplification was conducted using the Phusion High-Fidelity PCR

Tab. 2. Isolates used in this study. Newly deposited sequences are in bold.

Taxon	Culture Accession No. <sup>a</sup>	GenBank Accession No.	Reference
		ITS	
		TEF	
<i>Guignardia bidwellii</i>	CBS 11645	EU683672	Wikee <i>et al.</i> (2011)
<i>Guignardia gaultheriae</i>	CBS 447.70	JN592543	Su & Cai (2012)
<i>Phyllosticta bifrenariae</i>	CBS 128855 <sup>b</sup>	JF343565	Glienke <i>et al.</i> (2011)
<i>Phyllosticta brazilianiae</i>	CBS 126270	JF343572	Glienke <i>et al.</i> (2011)
<i>Phyllosticta capitalensis</i>	CBS 128856	JF261465	Glienke <i>et al.</i> (2011)
<i>Phyllosticta citriasiana</i>	CBS 120486	FJ538360	Wulandari <i>et al.</i> (2009)
<i>Phyllosticta citribraziliensis</i>	CBS 100098	FJ538352	Glienke <i>et al.</i> (2011)
<i>Phyllosticta citricarpa</i>	BRIP 52614 CBS 127454	JF343583	Glienke <i>et al.</i> (2011)
<i>Phyllosticta citrichaensis</i>	CBS 130529	JN791597	Wang <i>et al.</i> (2012)
<i>Phyllosticta hostae</i>	CGMCC 3.14355	JN692535	Su & Cai (2012)
<i>Phyllosticta hubeiensis</i>	CGMCC 3.14986	JX025037	Zhang <i>et al.</i> (2012)
<i>Phyllosticta ilicis-aquifolii</i>	CGMCC 3.14358	JN692538	Su & Cai (2012)
<i>Phyllosticta kerriae</i>	MAFF 240047 MUCC 0017	AB454266	Motohashi <i>et al.</i> (2009)
<i>Phyllosticta schimae</i>	CGMCC 3.14354	JN692534	Su & Cai (2012)
<i>Phyllosticta styracicola</i>	CGMCC 3.14985	JX025040	Zhang <i>et al.</i> (2012)
<i>Phyllosticta speewahensis</i> sp. nov.	BRIP 58044	<b>KF017269</b>	This study
<i>Saccharata capensis</i>	CBS 122693	EU552130	Marincowitz <i>et al.</i> 2008

<sup>a</sup> BRIP: Plant Pathology Herbarium, Brisbane, Australia; CBS: Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands; CGMCC: China General Microbial Culture Collection; MAFF: National Institute of Agrobiological Sciences, Tsukuba, Ibaraki Prefecture, Japan; MUCC: Laboratory of Plant Pathology, Mie University, Tsu, Mie Prefecture, Japan

<sup>b</sup> not an ex-type/ex-epitype isolate



**Fig. 1.** Maximum likelihood tree, with bootstrap (1000 replicates) values incorporated of nucleotide sequences of the ITS and TEF gene regions. The scale bar shows the number of substitutions per site. The tree was rooted to *S. capensis* (CBS 122693). The alignment is deposited in TreeBASE (S14236). All sequences are from ex-type cultures with the exception of *P. bifrenariae* (Tab. 2).

Master Mix (ThermoFisher Scientific, Scoresby, Vic., Australia), which consisted of 12.5  $\mu$ L of 2  $\times$  Master Mix with HF Buffer, 0.5  $\mu$ L each of 10 mM of forward and reverse primers, and 1  $\mu$ L of DNA template. The internal transcribed spacer (ITS) region was amplified with primers V9G (de Hoog & Gerrits van den Ende 1998) and ITS4 (White *et al.* 1990), and part of the translation elongation factor 1- $\alpha$  (TEF) was amplified with primers EF1-728 F (Carbone & Kohn 1999) and EF2 (O'Donnell *et al.* 1998). PCR products were amplified in a Bio-Rad C1000 thermal cycler (Bio-Rad Laboratories, Hercules, California, USA) using the following conditions: 98  $^{\circ}$ C for 30 s, 30 cycles at 98  $^{\circ}$ C for 10 s, 55  $^{\circ}$ C for 30 s, 72  $^{\circ}$ C for 30 s, followed by a 5 min final extension at 72  $^{\circ}$ C. The products were sequenced by Macrogen Incorporated (Seoul, Korea) using the AB 3730xl DNA Analyser (Applied Biosystems, Forster City, California, USA). The ITS and TEF sequences of BRIP 58044 were deposited in GenBank as KF017269 and KF017268, respectively.

The ITS and TEF sequences from isolate BRIP 58044 were assembled using Vector NTI Advance v.11 (Life Technologies, Mulgrave, Vic., Australia) and aligned against published sequences from GenBank. The ITS and TEF sequences of ex-type cultures from 16 taxa, along with sequences from *P.*

*bifrenariae* were included in the alignment (Tab. 2). *Saccarata capensis* was selected as the outgroup taxon as it has been shown to reside in the same clade as *Phyllosticta* within the family Phyllostictaceae (Liu *et al.* 2012). The ITS and TEF sequences were initially aligned separately using ClustalW in MEGA 5.2 (Tamura *et al.* 2011) and manually adjusted where needed. The most suitable maximum likelihood (ML) nucleotide substitution model for each gene was determined using the model test function in MEGA 5.2, and then used for the phylogenetic analysis of the ITS and TEF sequences individually, as well as for the concatenated ITS and TEF data. A ML tree using the concatenated data was generated in MEGA 5.2 using the Kimura-2 parameter substitution model with Gamma distribution.

## Results

The ITS and TEF alignments contained 573 and 211 bases, respectively. Individual analysis of ITS and TEF resulted in ML trees with the same topology (data not shown), therefore the sequence data for both genes were concatenated. The generated ML tree shows the *Phyllosticta* species described in this study clusters close to *P. hostae* Y.Y. Su & L. Cai, but is nonetheless a distinct species (Fig. 1). *Phyllosticta hostae* was recently described from a leaf of *Hosta plantaginea* (Liliaceae) in China (Su & Cai 2012).

## Taxonomy


***Phyllosticta speewahensis*** R. G. Shivas, Y. P. Tan & K. R. E. Grice, **sp. nov.** – Figs. 2–6.

Mycobank no.: MB 804255

**Description.** – Leaf spots amphigenous, narrowly linear, widest in the middle, up to 2 cm long and 1 mm wide, pale to dark reddish brown at the margins, becoming black in the centre, solitary, or in clusters of up to 5, with raised, black, solitary pycnidia, arranged along the central axis. Pycnidia 50–80 µm diam., globose, wall composed of layers of *textura angularis*, outer layer dark reddish brown. Conidiophores reduced to conidiogenous cells. Conidiogenous cells terminal, hyaline, smooth-walled, cylindrical, 5–20 × 2–4 µm. Conidia subglobose, broadly ellipsoidal or obovoid, with a truncate to rounded base and rounded apex, hyaline, (9) 10–13.5 (15) × 7.5–9 (9.5) µm, aseptate; wall uniformly 0.5 µm thick, enclosed in a mucilaginous sheath up to 3 µm wide, with an apical hyaline tapered appendage 5–17 µm long. Appressoria single, medium brown, ellipsoidal to clavate, or irregularly lobed, mostly curved, 7–14 × 5–8 µm, with a horizontal median septum, or aseptate, apex rounded, smooth-walled. Teleomorph not observed.

**Holotypus.** – AUSTRALIA, Queensland, Speewah, in leaf spots on *Vanda* hybrid ‘Waxy Blue’, 12 September 2012, leg. K. R. E. Grice, BRIP 58044, sequences ex-type in GenBank KF017269 (ITS), KF017268 (TEF).

**Culture characteristics** (after 4 d in the dark on PDA and a further 2 wk under 12 h ultraviolet light / 12 h dark cycle, at 23 °C). – Colonies



**Figs. 2-6.** *Phyllosticta speewahensis* (holotype). **2.** In leaf lesions on cultivated *Vanda* hybrid (bar = 1 cm). **3-4.** Colony on PDA after 14 d (bars 1 cm and 1 mm, respectively). **5.** Conidia (bar 10  $\mu$ m). **6.** Conidia (hyaline) and appressoria (brown) (bar 10  $\mu$ m).

1.5 to 2 cm diam., flat, without aerial mycelium, Greenish Black (Rayner 1970); reverse Dark Slate Blue (Rayner 1970).

**Etymology.** – Named after the town nearest to the location of the fungus.

**Host plant.** – *Vanda* hybrid (Orchidaceae).

**Distribution.** – Speewah, Queensland, Australia.

*Phyllosticta speewahensis* has similar sized conidia to *P. bifrenariae* (10–16  $\times$  7–9  $\mu$ m) and is morphologically and culturally similar to this species. Two morphological differences in the description of these species are that appressoria were found in cultures of *P. speewahensis* but not *P. bifrenariae*; and spermatia formed in cultures of *P. bifrenariae* but not *P. speewahensis*. The presence or absence of these morphological features is clearly an unreliable character in species differentiation as their production will be influenced by media and other environmental conditions. *Phyllosticta speewahensis* can only be reliably differentiated from *P. bifrenariae* by DNA

sequence data. *Phyllosticta speewahensis* has larger or wider conidia, (9) 10–13.5 (15) × 7.5–9 (9.5) μm, than the two other species reported from orchids, namely *P. aplectri* (5–8 × 4–6 μm) and *P. capitalensis* (10–14 × 5–7 μm) (van der Aa 1973, Glienke *et al.* 2011).

## Discussion

Of the numerous species of *Phyllosticta* that have been described from orchids, only *P. bifrenariae* has been demonstrated to be a pathogen (Silva *et al.* 2008 as *P. capitalensis*). Leaf spots caused by *Phyllosticta* spp. are common on a wide range of orchid genera and species (Cash and Watson 1955 as *P. pyriformis*, van der Aa 1973 as *P. capitalensis*). *Phyllosticta capitalensis sensu* Glienke *et al.* (2011) is now generally recognized as an endophyte of a wide range of plants from around the world (Wikee *et al.* 2013), although it was first described from an orchid (*Stanhopea*). There are likely to be a number of cryptic species of *Phyllosticta*, both endophytes and pathogens, yet to be discovered on orchids, as molecular phylogenetic methods are applied to culture specimens (Ko-Ko *et al.* 2011). The discovery of these hidden species of *Phyllosticta* spp. will certainly precede knowledge about their pathogenicity, distribution and biosecurity importance.

## References

- Aa H. A. van der (1973) Studies in *Phyllosticta* I. *Studies in Mycology* **5**: 1–110.
- Aa H. A. van der, Vanev S. (2002) *A revision of the species described in Phyllosticta*. Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands.
- Carbone I., Kohn L. M. (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* **91**: 553–556.
- Cash E. K., Watson A. J. (1955) Some fungi on Orchidaceae. *Mycologia* **47**: 729–747.
- Crous P. (2012) Fungal planet description sheet: 117. *Phyllosticta ericarum*. *Persoonia* **28**: 160–161.
- de Hoog G. S., Gerrits van den Ende A. H. G. (1998) Molecular diagnostics of clinical strains of filamentous Basidiomycetes. *Mycoses* **41**: 183–189.
- Glienke C., Pereira O. L., Stringari D., Fabris J., Kava-Cordeiro V., Galli-Terasawa L., Cunningham J. H., Shivas R. G., Groenewald J. Z., Crous P. W. (2011) Endophytic and pathogenic *Phyllosticta* species, with reference to those associated with Citrus Black Spot. *Persoonia* **26**: 47–56.
- Hennings P. (1908) Fungi S. Paulenses IV a cl. Puttemans collecti. *Hedwigia* **48**: 1–20.
- Ko-Ko T. W., McKenzie E. H. C., Bahkali A. H., To-Anun C., Chukeatirote E., Promputtha I., Abd-Elsalam K. A., Soyong, K., Wulandari N. F., Sanoamuang N., Jonglaekha N., Kobsueb R., Cheewangkoon R., Wikee S., Chamyuang S., Hyde, K. D. (2011) The need for re-inventory of Thai phytopathogens. *Chiang Mai Journal of Science* **38**: 1–13.
- Liu J.-K., Phookamsak R., Doilom M., Wikee S., Li Y.-M., Ariyawansa H., Boonmee S., Chomnunti P., Dai D.-Q., Bhat J. D., Romero A. I., Zhuang W.-Y., Monkai J., Jones G. E. B., Chukeatirote E., Ko Ko T. W., Zhao Y.-C., Wang Y., Hyde K. D. (2012) Towards a natural classification of *Botryosphaeriales*. *Fungal Diversity* **57**: 149–210.
- Marincowitz S., Groenewald J. Z., Wingfield M. J., Crous P. W. (2008) Species of *Botryosphaeriaceae* occurring on Proteaceae. *Persoonia* **21**: 111–118.
- Motohashi K., Inaba S., Anzai K., Takamatsu S., Kakashima C. (2009) Phylogenetic analyses of Japanese species of *Phyllosticta sensu stricto*. *Mycoscience* **50**: 291–302.

- O'Donnell K., Kistler H. C., Cigelnik E., Ploetz R. C. (1998) Multiple evolutionary origins of the fungus causing Panama disease of banana: Concordant evidence from nuclear and mitochondrial gene genealogies. *Proceedings of the National Academy of Sciences of the United States of America* **95**: 2044–2049.
- Petrak F. (1959) Beiträge zur Österreichischen Pilzflora. *Sydowia* **13**: 67–86.
- Rayner R. W. (1970) *A mycological colour chart*. CMI & British Mycological Society, Kew, Surrey, England.
- Shivas R. G., Tan Y. P., Grice K. R. E. (2012) Fungal planet description sheet: 131. *Phyllosticta aristolochiicola*. *Persoonia* **29**: 154–155.
- Silva M., Pereira O. L., Braga I. F., Leli S. M. (2008) Leaf and pseudobulb diseases on *Bifrenaria harrisoniae* (Orchidaceae) in Brazil. *Australasian Plant Disease Notes* **3**: 53–56.
- Su Y. Y., Cai L. (2012) Polyphasic characterisation of three new *Phyllosticta* spp. *Persoonia* **28**: 76–84.
- Tamura K., Peterson D., Peterson N., Stecher G., Nei M., Kumar G. (2011) MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony methods. *Molecular Biology and Evolution* **28**: 2731–2739.
- Wang X., Chen G., Huang F., Zhang J., Hyde K. D., Li H. (2012) *Phyllosticta* species associated with citrus diseases in China. *Fungal Diversity* **52**: 209–224.
- White, T. J., Bruns, T., Lee, S., Taylor, J. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR Protocols: A Guide to Methods and Applications* (eds. Innis M. A., Gelfand D. H., Sninsky J. J., White T. J.), Academic Press, San Diego: 315–322.
- Wikee S., Udayanga D., Crous P. W., Chukeatirote E., McKenzie, E. H. C. Bahkali A. H., Hyde K. D. (2011) *Phyllosticta* – An overview of current status of species recognition. *Fungal Diversity* **51**: 43–61.
- Wikee S., Lombard L., Crous P. W., Nakashima C., Motohashi K., Chukeatirote E., Alias S. A., McKenzie E. H. C., Hyde K. D. (2013) *Phyllosticta capitalensis*, a widespread endophyte of plants. *Fungal Diversity* (online: DOI 10.1007/s13225-013-0235-8).
- Wong M.-H., Crous P. W., Henderson J., Groenewald J. Z., Drenth A. (2012) *Phyllosticta* species associated with freckle disease of banana. *Fungal Diversity* **56**: 173–187.
- Wulandari N. F., To-Anun C., Hyde K. D., Duong L. M., de Gruyter J., Meffert J. P., Groenewald J. Z., Crous P. W. (2009) *Phyllosticta citriasiana* sp. nov., the cause of Citrus tan spot of *Citrus maxima* in Asia. *Fungal Diversity* **34**: 23–39.
- Zhang K., Su Y. Y., Cai L. (2012) Morphological and phylogenetic characterisation of two new species of *Phyllosticta* from China. *Mycological Progress* (online: DOI 10.1007/s11557-012-0861-7).

(Manuscript accepted 15 May 2013; Corresponding Editor: Kevin D. Hyde)



# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Sydowia](#)

Jahr/Year: 2013

Band/Volume: [65](#)

Autor(en)/Author(s): Shivas Roger Graham, Tan Yu Pei, Grice Kathy R. E.

Artikel/Article: [Phyllosticta species on orchids \(Orchidaceae\), introducing Phyllosticta speewahensis sp. nov. \(Phyllostictaceae, Ascomycota\) from northern Australia 139-146](#)