



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

19713-690

Date of Issuance:

12/21/16

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Chlorothalonil 720

Name and Address of Registrant (include ZIP Code):

Ms. Luz Chan
Registration Manager
Drexel Chemical Company
1700 Channel Avenue; PO Box 13327
Memphis, Tennessee 38113

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Tony Kish, Product Manager 22
Fungicide Branch, Registration Division (7505P)

Date:

12/21/16

2. You are required to comply with the data requirements described in the DCI identified below and in the following link:

<https://www.regulations.gov/document?D=EPA-HQ-OPP-2011-0840-0026>

- a. Chlorothalonil GDCI-081901-1301

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division: <http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. A one year study is required to satisfy these data requirements. You must submit these data by 12/21/2017.
4. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 19713-690."
 - Add an appropriate EPA Establishment Number & Net Contents Information
5. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 08/12/2016

If you have any questions, please contact Tony Kish by phone at 703 308-9443, or via email at kish.tony@epa.gov; or Craig Reeves by phone at 703 347-0486, or via email at reeves.craig@epa.gov

Enclosure: Stamped Accepted Label

ACCEPTED

Dec 21, 2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 19713-690

GROUP M FUNGICIDE

Drexel

Chlorothalonil 720

Fungicide

For control of diseases in listed agricultural, turf and ornamental crops.

ACTIVE INGREDIENT:

Chlorothalonil..... 54.0%

OTHER INGREDIENTS: 46.0%

TOTAL:100.0%

*This product contains 6 pounds of chlorothalonil per gallon equivalent to 720 grams per liter.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

**SHAKE WELL BEFORE USING
[RECIRCULATE CONTENTS BEFORE USE]**

See FIRST AID Below

EPA Reg. No. 19713-AON
EPA Est. No. 19713-XX-XXX

Net Content: _____ Gals. (_____ L)

FIRST AID

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call CHEMTREC at 800-424-9300 for emergency medical treatment information.

NOTE TO PHYSICIAN:

Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids.

Manufactured By:



Drexel Chemical Company

P.O. BOX 13327, MEMPHIS, TN 38113-0327

SINCE 1972

690SP-1216*P

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed absorbed through skin or inhaled Causes moderate eye irritation. Avoid contact with skin eyes or clothing. Avoid breathing spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

Mixers, loaders, applicators and all others who handle this pesticide must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVS) or viton, and shoes plus socks.

For uses in Turf and Ornamentals: Mixers, loaders, applicators and all others who handle this pesticide must wear: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVS) or viton, shoes plus socks and a NIOSH approved dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any N, R, P, or HE filter for applicators and handlers in enclosed areas such as a greenhouse.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: 1) Wash hands thoroughly before eating, drinking, chewing gum, using tobacco or using the toilet. 2) Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. 3) Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters frequently flooded areas, areas overlaying extremely shallow ground water areas with infield canals or ditches that drain to surface water areas not separated from adjacent surface waters with vegetated filter strips and areas overlaying tile drainage systems that drain to surface water.

PRODUCT INFORMATION

CHLOROTHALONIL 720 fungicide is an effective product when used according to label directions to control a broad spectrum of plant diseases. This product is suitable for use in programs that are compatible with the principles of Integrated Pest Management (IPM) which include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

This product can be used effectively in dilute or concentrate sprays. Thorough uniform coverage is essential for disease control.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the WPS.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the WPS and that involves contact with anything that has been treated, such as plants, soil or water is: Long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as nitrile or barrier laminate, and protective eyewear.

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the REI expires after 12 hours for the next 65 days, entry is permitted only when the following safety measures are provided.

1. At least one container designed specifically for flushing eyes must be available in operating condition at the WPS required decontamination site intended for workers entering the treated area.
2. Workers must be informed in a manner they can understand that:
 - that residues in the treated area may be highly irritating to their eyes
 - that they should take precautions such as refraining from rubbing their eyes to keep the residues out of their eyes
 - that if they do get residues in their eyes, they should immediately flush their eyes using the eye flush container that is located at the decontamination site or using other readily available clean water and
 - how to operate the eye flush container.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS), 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and greenhouses.

Do not enter or allow others to enter treated area until sprays have dried.

DISEASE RESISTANCE MANAGEMENT

GROUP	M	FUNGICIDE
-------	----------	-----------

This product is a Group M* fungicide. Fungal isolates with acquired resistance to Group M fungicide may eventually dominate the fungal population if Group M fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by this product or other Group M fungicides.

To delay fungicide resistance, consider:

- Avoiding consecutive use of this product or other target site of action Group M fungicides that have a similar target site of action on the same pathogens.
- Using tank-mixtures or pre-mixes with fungicides from different target site of action Groups as long as the involved products are all registered for the same use and are both effective at the mix or pre-pack rate on the pathogens(s) of concern.
- Basing fungicide use on a comprehensive IPM program.
- Monitoring treated fungal populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for fungicide resistance management and/or IPM recommendations for specific crops and resistant pathogens.

*The multi-site activity grouping, designated by the symbol "M", comprises a collection of various chemicals that act as general toxophores with several sites of action. These sites may differ between group members.

USE RESTRICTIONS

- Do not use this product on crops grown in the greenhouse.
- Do not combine this product in spray tank with pesticides surfactants or fertilizers unless prior use has shown the combination to be physically compatible, effective and non-injurious under your conditions of use.
- Do not combine this product with *Bacillus thuringiensis* (e.g., Dipel®) or spreader/sticker (e.g., Latron B-1956® or Latron AG 98®) as phytotoxicity may result from the combination when applied to crops listed on this label.
- Do not apply within 150 feet (for aerial applications) or 25 feet (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

SPRAY DRIFT PRECAUTIONS

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 75% the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the “*SPRAY DRIFT MANAGEMENT*” section.

Spray Drift Management

This section is advisory in nature and does not supersede the mandatory label requirements.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the Wind, Temperature and Humidity, and Temperature Inversions sections of this label).

Controlling Droplet Size

- Volume: Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.
- Pressure: Use the lower spray pressures listed for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles: Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation: Orienting nozzles so that the spray is released backwards, parallel to the airstream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle Type: Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length

For some use patterns, reducing the effective boom length to less than three-fourths of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Do not make applications at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Increase swath adjustment distance increase, with increasing drift potential (higher wind, smaller droplets, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Avoid application below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

Apply this product when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

APPLICATION INSTRUCTIONS

Dosage rates on this label indicate pints of this product per acre unless otherwise stated. Under conditions favoring disease development, use the higher rate specified and shortest application interval.

Note: Slowly invert container several times to assure uniform mixture.

Add slowly the required amount of this product into the spray tank during filling. With concentrate sprays, pre-mix the required amount of this product in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations

Apply this product in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth.

For field and row crops, spray volume usually will range from 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground sprays and aircraft applications.

For tree and orchard crops, apply this product in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. For fruit and nut bearing crops, the maximum volume is 300 gallons per acre unless otherwise stated in the specific use directions. For Conifers, the maximum volume is 100 gallons per acre

APPLICATION AND CALIBRATION TECHNIQUES FOR CHEMIGATION

Apply this product only through sprinkler irrigation systems including center pivot, motorized lateral move, traveling gun, solid set or portable (wheel move side roll end tow or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.

Crop injury lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other irrigation experts.

Do not apply this product through irrigation systems connected to a public water system Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments should the need arise.

The irrigation water pipeline must be fitted with a functional automatic quick closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure dram located between the irrigation water pump and the check valve to prevent back-siphoning of treated irrigation water into the water source.

Always inject this product into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional normally closed solenoid-operated valve located on the intake side of the injection pump Interlock this valve to the power system so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

This product may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump of either diaphragm or piston type constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2 to 3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems

Thoroughly mix specified amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run but continue to operate irrigation system until this product has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move Side Roll, End Tow or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides. However, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a 30 to 45 minute period. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this product plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Agitation is recommended. This product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until this product has been cleared from last sprinkler head.

APPLICATION DIRECTIONS

ASPARAGUS

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Cercospora blight (<i>Cercospora asparagi</i>)	2 to 4 pts. (1.5 to 3.0 lbs. a.i.)	By ground applications only. Apply in water volumes of 25 to 50 gals./Ac. Begin applications after final harvest of spears. Repeat at 14 to 28 day intervals. Minimum retreatment interval is 14 days depending on disease pressure. Use the higher rate and shorter interval if disease severity begins to increase or weather conditions favor disease development.
Purple spot (<i>Pleospora herbarum</i>)		
Rust (<i>Puccinia asparagi</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> • Do not apply more than 12 pts. of this product (9 lbs. a.i.) per acre per year. • Do not apply within 190 days (120 days in AZ and CA) of harvest of spears in the following year. 		

BEANS (DRY)

Adzuki, Blackeyed pea, Broad, Catjang, Chickpea (Garbanzo), Cowpea, Dry, Jackbean, Kidney, Lablab, Lima, Lupin, Lupin grain, Moth, Mungbean, Navy, Pink, Pinto, Rice, Runner, Southern pea, Tepary, Urd, Yardlong (Except Soybeans)

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Anthracnose (<i>Colletotrichum lindemuthianum</i>)	1.37 to 2 pts. (1 to 1.5 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Begin applications at first onset of disease which may occur as early as 2 to 4 weeks before flowering. Repeat at 7 to 10 day intervals. For use only on Beans to be harvested dry with pods removed.
Ascochyta blight (<i>Ascochyta phaseolorum</i>)		
Cercospora leaf blotch (<i>Cercospora cruenta</i>)		
Downy mildew (<i>Phytophthora nicotianae</i>)		
Rust (<i>Uromyces appendiculatus</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 8 pts. of this product (6 lbs. a.i.) per acre per year. Do not apply within 14 days of harvest. 		

BEANS (SNAP)

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Botrytis blight (Gray mold) (<i>Botrytis cinerea</i>)	1.37 to 3 pts. (1 to 2.25 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens. Repeat as necessary. Minimum retreatment interval is 7 days to maintain control.
Rust (<i>Uromyces appendiculatus</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 12 pts. of this product (9 lbs. a.i.) per acre per year. Do not apply within 7 days of harvest. 		

BLUEBERRIES

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Suppression: Anthracnose (Ripe rot) (<i>Colletotrichum gloeosporoides</i>)	3 to 4 pts. (2.25 to 3 lbs. a.i.)	By ground or air applications only. Use as part of an overall disease management strategy which includes alternation with a fungicide with a different mode of action. Diseases may only be suppressed and russetting may occur under heavy disease pressure or unfavorable environmental conditions. Apply with sufficient water to obtain adequate coverage, normally 20 to 100 gals./Ac. Begin applications at budbreak (green tip) and repeat at 10 day intervals through early bloom. Under heavy disease pressure, use the higher rate.
Suppression: Mummy berry (<i>Monilia vaccinii-corymbosi</i>)		
Septoria leaf spot (<i>Septoria albopunctata</i>)	3 to 4 pts. (2.25 to 3 lbs. a.i.)	Post-harvest Foliar Use (after all berries have been harvested): To maintain healthy leaves for the following year, apply with sufficient water to obtain adequate coverage (normally 20 to 100 gals./Ac.). Repeat at 10 to 14 day intervals.
Rust (<i>Pucciniastrum vaccinia</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 12 pts. of this product (9 lbs. a.i.) per acre per year. Do not apply after full bloom (except for foliar use after harvest) or within 42 days of harvest. 		

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, CAVALO (BROCCOLO, KOHLRABI), CHINESE BROCCOLI, CHINESE CABBAGE (Tight-headed Var., Napa), CHINESE MUSTARD

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Alternaria leaf spot (<i>Alternaria</i> spp.)	1.5 pts. (1.125 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Begin applications after transplants are set in field or shortly after emergence of field-seeded crop or when conditions favor disease development. Repeat at 7 to 10 day intervals.
Downy mildew (<i>Peronospora parasitica</i>)		
Ringspot (CA only) (<i>Mycosphaerella brassicicola</i>)	2 pts. (1.5 lbs. a.i.)	For field-seeded Brussels sprouts, begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals.
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 16 pts. of this product (12 lbs. a.i.) per acre per year. Do not apply within 7 days of harvest. 		

CARROTS

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Cercospora leaf spot (<i>Cercospora carotae</i>)	1.5 to 2 pts. (1.125 to 1.5 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Begin applications when disease threatens. Repeat 7 to 10 day intervals.
Alternaria leaf blight (<i>Alternaria dauci</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 20 pts. of this product (15 lbs. a.i.) per acre per year. May be applied to the day of harvest. 		

CELERY

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Basal stalk rot (<i>Rhizoctonia solani</i>)	2 to 3 pts. (1.5 to 2.25 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Start applications when transplants are set in the field. Repeat at 7 day interval as needed to maintain control.
Early blight (<i>Cercospora apii</i>)		
Late blight (<i>Septoria apicola</i>)		
Suppression (7 day schedule): Pink rot (<i>Sclerotinia sclerotiorum</i>)	3 pts. (2.25 lbs. a.i.)	
Early blight (<i>Cercospora apii</i>)	1.5 to 2 pts. (1.25 to 1.5 lbs. a.i./100 gals.)	For Celery seedbeds, apply in a spray volume of 125 gals./Ac. twice weekly or as needed to maintain control. Start applications shortly after crop emergence. Use the higher rate under severe disease conditions.
Late blight (<i>Septoria apicola</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 24 pts. of this product (18 lbs. a.i.) per acre per year. Do not apply within 7 days of harvest. 		

CORN (Sweet, Grown for Seed)

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Helminthosporium leaf blights (<i>Helminthosporium</i> spp.)	0.75 to 2 pts. (0.6 to 1.5 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development. Repeat at 7 day intervals. Under severe disease conditions, use 1.5 to 2 pts. of this product per acre.
Rust (<i>Puccinia</i> spp.)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 12 pts. of this product (9 lbs. a.i.) per acre per year. Do not apply within 14 days of harvest. Do not apply to Sweet corn to be processed. Do not allow livestock to graze in treated areas. Do not ensile treated Corn or use as livestock forage, 		

CRANBERRY

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Fruit rots	4 to 6.5 pts. (3 to 4.9 lbs. a.i.)	By ground, air or chemigation applications. Apply at early bloom. Repeat at 10 to 14 day intervals. Under severe disease conditions, use the 6.5 pts./Ac. on a 10 day schedule. When applying by chemigation, apply in 300 gallons of water per acre through solid set systems only.
Lophodermium leaf/twig blight (<i>Lophodermium hypophyllum</i>)		
Upright dieback (<i>Phomopsis vaccinii</i>)	4 to 6.5 pts. (3 to 4.9 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain coverage of uprights and runners. Make the first application prior to bloom when shoots begin growth in the Spring. Make additional applications at 10 to 14 day intervals. When applying by chemigation, apply in 300 gallons of water per acre through solid set systems only.
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 20 pts. of this product (15 lbs. a.i.) per acre per year. Do not apply within 50 days of harvest. Do not apply to beds when flooded or allow release of irrigation water from beds for at least 3 days following application. 		

CUCURBITS: CANTALOUPE, CHAYOTE, CHINESE WAXGOURD (CHINESE PRESERVING MELONS), CUCUMBER, HONEYDEW MELON, *MOMORDICA* SPP. (INCLUDES BALSAM APPLE, BITTER MELON), MUSKMELON, PUMPKIN, SQUASH, WATERMELON, ZUCCHINI) INCLUDING CULTIVARS AND/OR HYBRIDS OF THESE

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Anthrachnose (<i>Colletotrichum</i> spp.)	1.5 to 2 pts. (1.125 to 1.5 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals.
Downy mildew (<i>Peronospora parasitica</i>)		
Target spot (<i>Corynespora cassiicola</i>)		
Alternaria leaf blight (<i>Alternaria cucumerina</i>)	2 to 3 pts. (1.5 to 2.25 lbs. a.i.)	Note: Spraying mature Watermelons may result in sunburn of the upper surface of the fruit. DO NOT apply this product to Watermelons when any of the following conditions are present: <ul style="list-style-type: none"> intense heat and sunlight drought conditions poor vine canopy other crop and environmental conditions which may be conducive to increased natural sunburn DO NOT combine this product with anything except water for application to Watermelons unless your prior use has shown the combination to be non-injurious to Watermelons under your conditions of use.
Alternaria leaf spot (<i>Alternaria alternata</i>)		
Cercospora leaf spot (<i>Cercospora citrullina</i>)		
Gummy stem blight / Stem decline (<i>Didymella bryoniae</i>)		
Powdery mildew (<i>Sphaerotheca</i> sp.)		
Scab (<i>Cladosporium cucumerinum</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 21 pts. of this product (15.75 lbs. a.i.) per acre per year. May be applied to the day of harvest. 		

FRUITING VEGETABLES (EXCEPT TOMATO): EGGPLANT, GROUNDCHERRY, OKRA, PEPINO, PEPPERS (INCLUDES BELL, CHILI, COOKING, PIMENTO, SWEET), TOMATILLO

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Anthrachnose (<i>Colletotrichum</i> spp.)	1.5 pts. (1.125 lbs. a.i.)	By ground, air or chemigation applications. Begin application as foliage, flower, and fruit spray when disease is anticipated. Repeat at 7 to 10 day intervals
Botrytis leaf mold (<i>Botrytis cinerea</i>)		
Cercospora leaf spot (<i>Cercospora</i> spp.)		
Powdery mildew (<i>Leveillula taurica</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 12 pts. of this product (9 lbs. a.i.) per acre per year. Do not apply within 3 days of harvest. 		

GINSENG

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Alternaria blight (<i>Alternaria panax</i>)	2 pts. (1.5 lbs. a.i.)	Use sufficient water to obtain adequate coverage. Begin applications when disease threatens. Repeat on a 7 to 10 day intervals as disease pressure warrants.
Grey mold (<i>Botrytis cinerea</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 16 pts. of this product (12 lbs. a.i.) per acre per year. Do not apply within 14 days of harvest. 		

GRASSES GROWN FOR SEED

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Bipolaris and Dreschlera leaf spots	1 to 1.5 pts. (0.75 to 1.125 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Begin applications during stem elongation when conditions favor disease development. Re-apply at flag (top) leaf emergence. Repeat at 14 day intervals.
Glume blotch		
Rust (Leaf, Stem, Stripe)		
Septoria leaf spot (<i>Septoria</i> spp.)		
Selenophoma (Eyespot)	1 to 2 pts. (0.75 to 1.5 lbs. a.i.)	
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 6 pts. of this product (4.5 lbs. a.i.) per acre per year. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated areas or feed hay produced before harvest. Feeding of treated plant parts after harvest of seed is allowed. 		

HORSERADISH

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Ramularia stem and leaf spots (<i>Ramularia armoraciae</i>)	3 pts. (2.25 lbs. a.i.)	Use sufficient water to obtain adequate coverage. Begin applications when disease threatens. Repeat at 7 to 10 day intervals as disease pressure warrants.
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 24 pts. of this product (18 lbs. a.i.) per acre per year. Do not apply within 14 days of harvest. 		

LEEK

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Botrytis leaf blight (<i>Botrytis</i> spp.)	1.5 to 3 pts. (1.125 to 2.25 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods and repeat at 7 to 10 day intervals for as long as conditions favor disease. Use the high rate and a 7 day schedule of applications when heavy dew or rain persists.
Purple blotch (<i>Alternaria porii</i>)		
Suppression: Downy mildew (<i>Peronospora destructor</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 9 pts. of this product (6.75 lbs. a.i.) per acre per year. Do not apply within 14 days of harvest. 		

MANGO

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Anthraxnose (<i>Colletotrichum</i> spp.)	2 to 3.5 pts. (1.5 to 2.6 lbs. a.i.)	By ground or air applications only. Apply in a water volume of 20 to 300 gals./Ac. Begin applications at early bloom. Begin the season with the 2 pint rate on a 14 day interval. Repeat at 7 to 14 day intervals until early fruit development. If disease pressure is severe, use the higher rate and shorter interval.
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 32 pts. of this product (24 lbs. a.i.) per acre per year. Do not apply within 21 days of harvest. 		

MINT

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Rust (<i>Puccinia menthae</i>)	1.37 pts. (1 lb. a.i.)	By ground or air applications only. Apply with sufficient water to obtain adequate coverage, normally 20 to 150 gals./Ac. for dilute sprays and 5 to 10 gals./Ac. for concentrate ground and aircraft applications. Begin applications when emerging plants are 4 to 8 inches high. Repeat at 7 to 10 day intervals to maintain control.
Septoria leaf spot (<i>Septoria menthae</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 4 pts. of this product (3 lbs. a.i.) per acre per year. Do not apply within 80 days of harvest. Do not feed fresh or extracted Mint hay from treated fields to livestock. 		

MUSHROOM

Disease(s)/Fungus	This Product	Use Instructions
Verticillium brown spot (<i>Verticillium</i> spp.)	2.75 to 5.5 fl. ozs./1,000 sq. ft.	Apply as a drench to the Mushroom bed surface in at least 125 gallons of water per 1,000 sq. ft. of Mushroom bed. Make two applications. Apply the high rate (5.5 fl. ozs.) of this product in the first application and the low rate (2.75 fl. ozs.) of this product in the second application. Make the first application within two days of top dressing the spawn colonized Mushroom compost with a casing layer. Make the second application at pinning.
Dry bubble		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 8.25 fl. ozs. of this product per 1,000 sq. ft. per cropping cycle. Do not make more than 2 applications per cropping cycle. Do not apply within 5 days of harvest. 		

ONIONS (Dry Bulb) & GARLIC

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Botrytis leaf blight (<i>Botrytis</i> spp.)	1 to 3 pts. (0.75 to 2.25 lbs. a.i.)	By ground, air or chemigation applications. Minimum retreatment interval is 7 days. Apply with sufficient water to obtain thorough coverage of tops. This product can be used with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows: • Low disease hazard and before infection: 1 pt./Ac. at 10 day intervals • Low disease hazard and some disease present: 1.37 pt./Ac. At 7 to 10 day intervals • High disease hazard: 3 pts./Ac. at 7 day intervals For suppression of Neck rot (<i>Botrytis</i> spp.) during storage, apply a minimum of 3 weekly applications prior to lifting using 1.37 to 3 pts./Ac..
Purple blotch (<i>Alternaria porii</i>)		
Suppression: Botrytis neck rot (<i>Botrytis</i> spp.)		
Suppression: Downy mildew (<i>Peronospora destructor</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 20 pts. of this product (15 lbs. a.i.) per acre per year. Do not apply within 7 days of harvest. 		

ONIONS (Green Bunching, Grown for Seed), GARLIC (Grown for Seed), SHALLOTS

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Botrytis leaf blight (<i>Botrytis</i> spp.)	1.5 to 3 pts. (1.125 to 2.25 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods. Repeat at 7 to 10 day intervals for as long as conditions favor disease. Use the high rate and a 7 day schedule of applications when heavy dew or rain persists.
Purple blotch (<i>Alternaria porii</i>)		
Suppression: Downy mildew (<i>Peronospora destructor</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 9 pts. of this product (6.75 lbs. a.i.) per acre per year. Do not apply with 7 days of harvest of Garlic. Do not apply within 14 days of harvest of green bunching Onions or Shallots. 		

PAPAYA

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Alternaria fruit spot (<i>Alternaria alternata</i>)	1.5 to 3 pts. (1.125 to 2.25 lbs. a.i.)	By ground applications only. Apply with sufficient water to obtain adequate coverage of fruit and leaves. Begin applications when conditions favor development of disease. Continue treatments at 14 day intervals until weather conditions no longer favor disease development.
Anthracoise (<i>Colletotrichum</i> spp.)		
Stem end rot (<i>Colletotrichum</i> spp.)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 9 pts. of this product (6.75 lbs. a.i.) per acre per year. May be applied to the day of harvest. 		

PARSNIP

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Alternaria leaf spot (<i>Alternaria</i> spp.)	1.5 to 2 pts. (1.125 to 1.5 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage of fruit and leaves. Begin applications at the first sign of disease or when conditions are favorable for infection. Continue treatments at 7 to 10 day intervals.
Anthracoise (<i>Colletotrichum</i> spp.)		
Botrytis blight (Gray mold) (<i>Botrytis cinerea</i>)		
Bottom rot (<i>Rhizoctonia</i> spp.)		
Down mildew (<i>Plasmopara crustosa</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 8 pts. of this product (6 lbs. a.i.) per acre per year. Do not apply within 10 days of harvest. 		

PEANUTS

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Early leaf spots (<i>Cercospora arachidicola</i>)	1 to 1.5 pts. (0.75 to 1.125 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage when leaf wetness first occurs or 30 to 40 days after planting of fruit and leaves. Repeat at 14 days intervals. When conditions favor Late leaf spot or when Rust or Web blotch occur, apply 1.5 pts./Ac. of this product at 14 day intervals for the remainder of the season.
Late leaf spot (<i>Cercosporidium personatum</i>)		
Pepper spot (<i>Leptosphaerulina crassiasca</i>)		
Rust (<i>Puccinia arachidis</i>)	1.5 pts. (1.125 lbs. a.i.)	
Web blotch (<i>Phoma arachidicola</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> • Do not apply more than 12 pts. of this product (9 lbs. a.i.) per acre per year. • Do not apply within 14 days of harvest. • Do not allow livestock to graze treated areas. • Do not feed hay or threshings from treated fields to livestock. 		

POTATOES

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Black dot (<i>Colletotrichum coccodes</i>)	0.75 pt. (0.6 lbs. a.i.)	By ground, air or chemigation applications. Begin applications at the low rate (0.75 pt./A) when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals. Then begin applying the higher label rates (1 to 1.5 pts./Ac.) at 5 to 10 days intervals when any one of the following events occur: <ul style="list-style-type: none"> • Vines close within the rows • Late blight forecasting measures 18 disease severity values (DSV) • The crop reaches 300 P-days Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe. DO NOT exceed a 10 day interval between applications when using chemigation.
Botrytis vine rot (<i>Botrytis cinerea</i>)		
Early blight (<i>Alternaria solani</i>)	1 to 1.5 pts. (0.75 to 1.125 lbs. a.i.)	
Late blight (<i>Phytophthora infestans</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> • Do not apply more than 15 pts. of this product (11.25 lbs. a.i.) per acre per year. • Do not apply within 7 days of harvest. 		

RHUBARB

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Ascochyta (<i>Ascochyta rhei</i>)	3 pts. (2.25 lbs. a.i.)	Use sufficient water to obtain adequate coverage. Begin applications when disease threatens. Repeat at 7 to 10 day intervals as disease pressure warrants.
Ramularia leaf spot (<i>Ramularia rhei</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> • Do not apply more than 18 pts. of this product (13.5 lbs. a.i.) per acre per year. • Do not apply within 30 days of harvest. 		

SOYBEANS

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Anthracnose (<i>Colletotrichum truncatum</i>)	1.5 to 2.25 pts. (1.125 to 1.7 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain complete coverage using at least 5 gals. of water per acre for aerial application. Minimum retreatment interval is 14 days. Two application program: For determinate varieties make the first application at early pod set stage (R3) and the second application at seed formation (R5). For indeterminate varieties, make the first application when largest pods are 1 to 1.25 inches in length. Make the second application 14 days later.
Diaprothe pod and stem rot (<i>Diaprothe phaseolorum</i>)		
Frogeye leaf spot (<i>Cercospora sojina</i>)		
Purple seed stain (<i>Cercospora kikuchii</i>)		
Cercospora leaf blight (<i>Cercospora kikuchii</i>)		
Septoria brown spot (<i>Septoria glycines</i>)	1 to 2 pts. (0.75 to 1.5 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain complete coverage using at least 5 gals. of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. Minimum retreatment interval is 14 days. Three application program: For determinate varieties, make the first application at the beginning of flowering (R1), the second at early pod set (R3) and the third at beginning of seed formation (R5). For indeterminate varieties, make the first application one week after first flowering and continue applications at 14 day intervals.
Suppression: Rust (<i>Phakopsora pachyrhizi</i>)		
Stem canker (<i>Diaporthe phaseolorum</i>)	1 pt. (0.75 lb. a.i.)	Apply with 10 to 20 gals. of water per acre as a band treatment directing spray to provide coverage of entire plant. Make the first application at time of emergence of the second trifoliate leaves (V2). If conditions favor stem canker disease, make a second and third application. Make all applications at 14 day intervals.
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 6 pts. of this product (4.5 lbs. a.i.) per acre per year. Do not apply within 6 weeks of harvest. Do not feed hay or threshing from treated fields to livestock. 		

STRAWBERRIES

Disease(s)/Fungus	This Product	Use Instructions
Ramularia leaf spot (<i>Ramularia tulasnei</i>)	1.5 pts. / Ac (1.125 lbs. a.i.)	Begin application after a rain or sprinkler irrigation application and when disease threatens. Repeat at 10 to 14 days intervals. Use the shortest specified interval when disease conditions are severe. Continue applications until runners are dug. Pre-harvest interval is not applicable.
USE RESTRICTIONS:		
<ul style="list-style-type: none"> Do not apply more than 20 pts. of this product (15 lbs. a.i.) per acre per year. Only for nursery seedlings for pre-transplant. 		

TOMATOES

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Foliage: Early blight (<i>Alternaria solani</i>)	1.37 to 2 pts. (1 to 1.5 lbs. a.i.)	By ground, air or chemigation applications. Apply with sufficient water to obtain adequate coverage. Begin applications when dew or rain occurs and disease threatens. Apply on a 7 to 10 day intervals for foliage diseases. For fruit diseases, begin at fruit set and apply on a 7 to 14 day intervals. Use the highest rate and shortest interval when disease conditions are severe.
Foliage: Gray leaf mold (<i>Fluvia fluva</i> ; <i>Cladosporium</i>)		
Foliage: Gray leaf spot (<i>Stemphyllium botryosum</i>)		
Foliage: Late blight (<i>Phytophthora infestans</i>)		
Foliage: Septoria leaf spot (<i>Septoria lycopersici</i>)		
Foliage: Target spot (<i>Corynespora cassiicola</i>)		
Fruit: Alternaria fruit rot (Black mold) (<i>Alternaria alternata</i>)	2 to 3 pts. (1.5 to 2.25 lbs. a.i.)	
Fruit: Anthracnose (<i>Colletotrichum</i> spp.)		
Fruit: Botrytis gray mold (<i>Botrytis cinerea</i>)		
Fruit: Late blight fruit rot (<i>Phytophthora infestans</i>)		
Fruit: Rhizoctonia fruit rot (<i>Rhizoctonia solani</i>)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> • Do not apply more than 20 pts. of this product (15 lbs. a.i.) per acre per year. • May be applied to the day of harvest. 		

YAM

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Anthracnose (<i>Colletotrichum gloeosporioides</i>)	1 to 1.25 pts. (0.75 to 1.125 lbs. a.i.)	Use sufficient water to obtain adequate coverage. Begin applications when disease threatens. Repeat at 10 to 14 day intervals as disease pressure warrants.
USE RESTRICTIONS:		
<ul style="list-style-type: none"> • Do not apply more than 15 pts. of this product (11.25 lbs. a.i.) per acre per year. • Do not apply within 7 days of harvest. 		

TREES AND ORCHARD CROPS

Apply this product in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. For fruit and nut bearing crops, the maximum volume is 300 gallons per acre unless indicated otherwise in the specific use directions. For Conifers, the maximum volume is 100 gallons per acre.

Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, this product may be applied with aircraft using at least 20 gallons of spray per acre. The minimum volume for application by aircraft to Conifer stands and Christmas trees is 10 gallons per acre.

When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate specified may be used. The following spray volumes may be used.

TREES AND ORCHARD CROPS	Spray Volume (Gals./Ac.)	
	Concentrate	Dilute
Apricot Cherry (Tart) Nectarine Peach Plum Prune	20	300
Cherry (Sweet)	20	400
Conifer stands	10 to 20	Not used
Christmas trees	10 to 50 (By air or ground)	100
Nursery beds	5 to 10 (By ground only)	100

USE RESTRICTION

Do not allow livestock to graze in treated areas.

ALMONDS

Disease(s)/Fungus	This Product		Use Instructions
	Per Acre	Per 100 Gals*	
Anthracnose (<i>Colletotrichum cutatum</i>)	4 pts. (3 lbs. a.i.)	1.33 pts. (1 lb. a.i.)	By ground or air applications only. Apply with water volumes of 20 to 300 gals./Ac. For Brown rot blossom and twig blight, begin application at popcorn (pink bud) and at full bloom. If weather is still conducive for disease development, another application may be made at petal fall. For control of Shothole, make an application in Autumn at leaf fall. In the Spring, make the first application at bud break followed by an application at shuck split to control nut infections and to control Scab. *To be used only with full dilute spray volume.
Brown rot blossom and twig blight (<i>Monilinia</i> spp.)			
Shot hole (<i>Wilsonomyces carpophilus</i>)			
Scab (<i>Venturia carpophila</i>)			
USE RESTRICTIONS:			
<ul style="list-style-type: none"> Do not apply more than 25 pts. of this product (18.75 lbs. a.i.) per acre per year (leaf fall through shuck split). Do not apply within 150 days of harvest. 			

APRICOT, CHERRY, NECTARINE, PEACH, PLUM, PRUNE

Disease(s)/Fungus	This Product		Use Instructions
	Per Acre	Per 100 Gals*	
Leaf curl (<i>Taphrina deformans</i>)	3.12 to 4.12 pts. (2.3 to 3.1 lbs. a.i.)	1 to 1.37 pts. (0.75 to 1 lb. a.i.)	By ground or air applications only. For best control, apply at leaf fall in late Autumn using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels, use the high rate of application. Apply once or twice more in mid to late Winter before bud swell. If leaf fall application is not practical, application of this product for control of Leaf curl may be made at any time prior to bud swell the following Spring. Where Shothole occurs, also apply at bud break to protect newly emerging leaves and at shuck split to prevent fruit infections. Minimum retreatment interval is 10 days. *Volumetric rates to be used only with full dilute spray volume.
Shot hole (<i>Coryneum blight</i>) (<i>Thyrostroma carpophilum</i>)			
Lacy (russet) scab (Plums/Prunes)	3.12 to 4.12 pts. (2.3 to 3.1 lbs. a.i.)	1 to 1.37 pts. (0.75 to 1 lb. a.i.)	Make one application at popcorn (pink red or early white bud) and a second application at full bloom. If weather conditions favor disease development make an additional application at petal fall. Minimum retreatment interval is 10 days. *Volumetric rates to be used only with full dilute spray volume.
Brown rot blossom blight (<i>Monilinia</i> spp.)			
Black knot (<i>Apiosporina morbosa</i>) (Cherries/Plums)	3.12 pts. (2.3 to 3.1 lbs. a.i.)	1 to 1.37 pts. (0.75 to 1 lb. a.i.)	By ground or air applications only. In addition to the bloom application listed Above, make one application at shuck split. Do not apply this product after shuck split and before harvest. If additional disease control is needed before harvest use, another registered fungicide. For control of Cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high Leaf spot incidence, make a second application 10 to 14 days later. Minimum retreatment interval is 10 days. *Volumetric rates to be used only with full dilute spray volume.
Cherry leaf spot (<i>Blumeriella jaapii</i>)			
Scab (<i>Cladosporium carpophilum</i>)			
Suppression: Constriction canker (<i>Phomopsis amygdali</i>)	3 to 4 pts. (2.25 to 3 lbs. a.i.)	0.96 to 1.33 pts. (0.73 to 1 lb. a.i.)	By ground or air application only. Apply after fruit is harvested when leaf drop starts. Apply in sufficient spray volume for thorough coverage (recommend 50 to 100 gallons per acre) to adequately cover exposed leaf scars. Repeat on 10 to 14 day interval until leaf drop is complete. Use in conjunction with cultural practices to reduce inoculum. *Volumetric rates to be used only with full dilute spray volume.
USE RESTRICTIONS:			
<ul style="list-style-type: none"> • Do not apply more than 20.5 pts. of this product (15.4 lbs. a.i.) per acre per year. • May be applied to the day of harvest. 			

FILBERTS (HAZELNUTS)

Disease(s)/Fungus	This Product		Use Instructions
	Per Acre	Per 100 Gals*	
Eastern Filbert blight (<i>Anisogramma anomata</i>)	4 pts. (3 lbs. a.i.)	1.33 pts. (1 lb. a.i.)	By ground or air applications only. Apply with water volumes of 20 to 300 gals./Ac. Begin applications at the onset of disease or when weather conditions favor disease development. Make applications on a 14 to 28 day schedule using the shorter interval under heavy disease pressure. *Volumetric rates to be used only with full dilute spray volume.
USE RESTRICTIONS:			
<ul style="list-style-type: none"> • Do not apply more than 12 pts. of this product (9 lbs. a.i.) per acre per year (leaf fall through shuck split). • Do not apply within 120 days of harvest. • Do not apply through irrigation. • Do not apply with oils, surfactants or fertilizers. • Do not apply within 1 week of an oil-based pesticide application. 			

PASSION FRUIT

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Alternaria fruit and leaf spots (<i>Alternaria</i> spp.)	2 pts. (1.5 lbs. a.i.)	By ground applications only. Apply with sufficient water to obtain adequate coverage of fruit and leaves. Begin applications when conditions favor development of disease. Continue treatments at 14 day intervals until weather conditions no longer favor disease development.
Anthrachnose (<i>Colletotrichum</i> spp.)		
Cercospora fruit spot (<i>Cercospora</i> spp.)		
USE RESTRICTIONS:		
<ul style="list-style-type: none"> • Do not apply more than 10 pts. of this product (7.5 lbs. a.i.) per acre per year. • Do not apply within 7 days of harvest. 		

PERSIMMON (FL & HI Only)

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Cercospora (<i>Cercospora fuliginosa</i>)	1.25 pts. (0.94 lbs. a.i.)	Use sufficient water to obtain adequate coverage. Begin applications when disease threatens. Repeat at 14 day intervals as disease pressure warrants.
USE RESTRICTIONS:		
<ul style="list-style-type: none"> • Do not apply more than 6.25 pts. of this product (4.7 lbs. a.i.) per acre per year. • Do not apply within 14 days of harvest. • Aerial applications require the use of a minimum of 10 gallons per acre. 		

PISTACHIO

Disease(s)/Fungus	This Product		Use Instructions
	Per Acre	Per 100 Gals*	
Suppression: Alternaria late blight (<i>Alternaria alternata</i>)	6 pts. (4.5 lbs. a.i.)	3 pts. (2.25 lbs. a.i.)	By ground or air applications only. Apply with water volumes of 20 to 200 gals./Ac. Make the first application at the beginning of the blossom period followed by an application at full bloom. Make additional applications as required at 28 day intervals. For Septoria and Botrytis, use the higher rate if disease pressure is severe. Note: Use of this product may result in speckling or reddening of the fruit hull (epicarp). This effect is superficial and has not resulted in change in nut quality. *Volumetric rates to be used only with full dilute spray volume.
Botryosphaera blight (<i>Botryosphaera dothidea</i>)			
Botrytis blight (<i>Botrytis cinerea</i>)	4 to 6 pts. (3 to 4.5 lbs. a.i.)	2 to 3 pts. (1.5 to 2.25 lbs.a.i.)	
Septoria leaf spot (<i>Septoria pistacina</i>)			
USE RESTRICTIONS:			
<ul style="list-style-type: none"> • Do not apply more than 30 pts. of this product (22.5 lbs. a.i.) per acre per year. • Do not apply within 14 days of harvest. 			

CONIFERS (INCLUDING CHRISTMAS TREES)

For use in: 1) Conifer nursery beds; 2) Christmas tree and bough production plantations; and 3) Tree seed orchards. **Use Restrictions:** Do not apply more than 22 pints of this product (16.5 lbs. a.i.) per acre per year. Do not use on forests.

Apply by ground or air in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. The minimum volume for application by aircraft to Conifer stands and Christmas trees is 10 gallons per acre. The maximum volume is 100 gallons per acre.

Disease(s)/Fungus	This Product Per Acre	Use Instructions
Autoecious needle rust (Weir's cushion rust) (<i>Chrysomyxa weirii</i>)	5.5 pts. (4.12 lbs. a.i.)	Begin applications when 10% of buds have broken and twice thereafter at 7 to 10 day intervals.
Botrytis seedling blight, Phoma twig blight	1.5 to 2.75 pts. (1.12 to 2.1 lbs. a.i.)	Begin applications in nursery beds when seedlings are 4 inches tall and when cool moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.
Cyclaneusma and Lophodermium needlecasts	2.75 to 5.5 pts. (2.1 to 4.12 lbs. a.i.)	Apply in early Spring prior to bud break. Repeat applications at approximately 6 to 8 week intervals until spore release ceases in late Fall. Apply monthly during periods of frequent rainfall and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended then resumed upon next occurrence of needle wetness.
Interior needle blight (<i>Mycosphaerella</i> spp. and <i>Phaeocryptopus nudus</i>)	2.75 to 5.5 pts. (2.1 to 4.12 lbs. a.i.)	One or two applications: In Christmas tree plantations or Conifer stands, make one application in the Spring when new shoot growth is 0.5 to 2 inches in length. Under high disease pressure, a second application may be made 10 to 14 days after the first application. When using aerial applications, use the highest specified rate.
	1.5 to 2.75 pts. (1.12 to 2.1 lbs. a.i.)	Multiple applications: Make the first application in Spring when new shoot growth is 0.5 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest specified rate on a 3 week schedule. When using aerial applications, use the highest specified rate.
Rhabdocline needlecast	1.5 to 2.75 pts. (1.12 to 2.1 lbs. a.i.)	Apply at bud break and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance or when irregular bud break occurs, apply weekly until all trees have broken bud then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3 week schedule.
Rhizosphaera needlecast (<i>Rhizosphaera</i> spp.)	5.5 pts. (4.12 lbs. a.i.)	Multiple applications: Make the first application in Spring when new shoot growth is 0.5 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest specified rate on a 3 week schedule. When using aerial applications, use the highest specified rate.
Scirrhia brown spot (<i>Mycosphaerella deamessii</i>)		
Scleroderris canker (<i>Gremmeniella abietina</i>)	1.5 to 2.75 pts. (1.12 to 2.1 lbs. a.i.)	
Sirococcus tip blight (<i>Sirococcus conigenus</i>)	2 to 3.5 pts. (1.5 to 2.6 lbs. a.i.)	

Swiss needlecast (<i>Phaeocryptopus gaeumannii</i>)	2.75 to 5.5 pts. (2.1 to 4.12 lbs. a.i.)	One or two applications: In Christmas tree plantations or Conifer stands, make one application in the Spring when new shoot growth is 0.5 to 2 inches in length. Under high disease pressure, a second application may be made 10 to 14 days after the first application. When using aerial applications, use the highest specified rate.
	1.5 to 2.75 pts. (1.12 to 2.1 lbs. a.i.)	Multiple applications: Make the first application in Spring when new shoot growth is 0.5 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule. When using aerial applications, use the highest specified rate.

TURF AND ORNAMENTALS

This product can be used to control diseases on Turf on Golf courses and Sod farms.

USE RESTRICTIONS

- Do not use on home lawns and Turf sites associated with apartment buildings, day care centers, playgrounds, play fields, recreational park, athletic fields, athletic fields located on or next to schools (i.e., elementary middle and high schools), campgrounds, churches and theme parks.
- Agricultural Use Sites Only (Sod farms, Golf courses, Nurseries and Greenhouses): This product must not be applied within 150 feet (for aerial applications) or 25 feet (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

USE PRECAUTIONS

Add slowly the required amount of this product into the spray tank during filling. With concentrate sprays, pre-mix the required amount of this product in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

APPLICATION DIRECTIONS

Golf Course Fairways and Roughs, Lawns Around Commercial and Industrial Buildings, and Professional and Collegiate Athletic Fields

Apply this product in sufficient amount of water to provide complete coverage. This amount may vary from 30 to 450 gallons per acre. See table below for suggested rates and timing.

Do not mow or water after treatment until spray deposited on Turfgrass is thoroughly dry. Use this product in conjunction with good Turf management practices.

Use Restrictions:

- Do not apply more than 34.7 pints of this product per acre (12.7 fl. ozs./1,000 sq. ft.) (26 lbs. a.i./Ac.) per year.
- Minimum retreatment interval for single application rates of up to 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) is 7 days.
- Do not apply more than one application of a rate greater than 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) per year.
- Maximum single application rate is 15.1 pints of this product per acre (5.5 fl. ozs./1,000 sq. ft.) (11.3 lbs. a.i./Ac.).

Golf Course Greens and Tees

Apply this product in sufficient amount of water to provide complete coverage. This amount may vary from 90 to 450 gallons per acre. See table below for suggested rates and timing. Under severe disease conditions use the highest specified rate and shortest interval corresponding with the application schedule selected from the table below.

Do not mow or water after treatment until spray deposited on Turfgrass is thoroughly dry. Use this product in conjunction with good Turf management practices.

Use Restrictions - Golf Course Greens:

- Do not apply more than 97.3 pints of this product per acre (35.7 fl. ozs./1,000 sq. ft.) (73 lbs. a.i./Ac.) per year.
- Minimum retreatment interval for single application rates of up to 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) is 7 days.
- Minimum retreatment interval after an application of a rate greater than 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) is 14 days.
- Do not apply more than two application of a rate greater than 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) per year.
- Maximum single application rate is 15.1 pints of this product per acre (5.5 fl. ozs./1,000 sq. ft.) (11.3 lbs. a.i./Ac.).

Use Restrictions - Golf Course Tees:

- Do not apply more than 69.3 pints of this product per acre (25.4 fl. ozs./1,000 sq. ft.) (52 lbs. a.i./Ac.) per year.
- Minimum retreatment interval for single application rates of up to 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) is 7 days.
- Minimum retreatment interval after an application of a rate greater than 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) is 14 days.
- Do not apply more than one application of a rate greater than 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) per year.
- Maximum single application rate is 15.1 pints of this product per acre (5.5 fl. ozs./1,000 sq. ft.) (11.3 lbs. a.i./Ac.).

Sod Farms

Apply this product in 30 to 450 gallons of water per acre. Do not mow or water after treatment until spray deposited on Turfgrass is thoroughly dry. Use this product in conjunction with good Turf management practices.

Use Restrictions:

- Sod farm turf treated with chlorothalonil prior to harvest must be mechanically cut rolled and harvested.
- Do not use this product on Sod farms at application rates greater than 17.3 pints (13 lbs. a.i.) per acre per year.
- Do not apply more than 17 pints of this product per acre (6.4 fl. ozs./1,000 sq. ft.) (13 lbs. a.i./Ac.) per year.
- The minimum retreatment interval for single application rates of up to 9.7 pints of this product per acre (3.5 fl. ozs. /1,000 sq. ft.) (7.3 lbs. a.i./Ac.) is 7 days.
- Do not apply more than one application of a rate greater than 9.75 pints of this product per acre (3.6 fl. ozs./1,000 sq. ft.) (7.3 lbs. a.i./Ac.) per year.
- Maximum single application rate is 15.1 pints of this product per acre (5.5 fl. ozs./1,000 sq. ft.) (11.3 lbs. a.i./Ac.).

APPLICATION TIMING (ALL TURF)

Begin applications when conditions favor disease development. Repeat applications as long as these conditions persist. Under severe disease conditions, use the highest specified rate and shortest interval corresponding with the application schedule selected from the table below.

Disease(s)/Fungus Controlled	Application Interval (Days)	Pre-Disease Rates			Post-Disease Rates*		
		Fl. oz./1,000 sq. ft.	Pts./Ac.	Lbs. A.I./Ac.	Fl. oz./1,000 sq. ft.	Pts./Ac.	Lbs. A.I./Ac.
Algae	7 to 14	2 to 3.6	5.5 to 9.75	4.1 to 7.3	2 to 3.6	5.5 to 9.75	4.1 to 7.3
	14	-	-	-	4 to 5.5	11 to 15.1	8.25 to 11.3
Anthracnose (<i>Colletotrichum graminicola</i>)	7 to 14	3 to 3.6	8.3 to 9.75	6.2 to 7.3	-	-	-
	14	3.6 to 5.5	9.9 to 15.1	7.4 to 11.3	-	-	-
Brown patch (<i>Rhizoctonia solani</i> , <i>R. zeae</i> , <i>R. cerealis</i>)	7 to 14	2 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	-	-	-	4 to 5.5	11 to 15.1	8.25 to 11.3
Copper spot (<i>Gloeocercospora sorghi</i>)	14	4 to 5.5	11 to 15.1	8.25 to 11.3	5.5	15.1	11.3
Dichondra leaf spot (CA only) (<i>Alternaria</i> spp.)	14	4 to 5.5	11 to 15.1	8.25 to 11.3	5.5	15.1	11.3
Dollar spot (<i>Sclerotinia homeocarpa</i> , <i>Lanzia</i> or <i>Moellerodiscus</i> spp.)	7 to 10	1.0** to 2	2.8** to 5	2.1** to 4.1	-	-	-
	7 to 21	2 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	-	-	-	4 to 5.5	11 to 15.1	8.25 to 11.3
Fusarium patch (Gerlachia) (<i>Micordochium nivale</i>)	21 to 28	5.5	15.1	11.3	-	-	-
Gray leaf spot (<i>Pyricularia grisea</i> , <i>Pyricularia oryzae</i>)	7 to 10	2 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	-	-	-	-	4 to 5.5	11 to 15.1	8.25 to 11.3
Gray snow mold*** (<i>Typhula</i> spp.)	30	5.5	15.1	11.3	-	-	-
Leaf spot , Melting out, Brown blight (<i>Bipolaris sorokiniana</i> ; <i>Drechslera</i> spp. including <i>D.</i> <i>poae</i> , <i>D. siccans</i> ; <i>Curvularia</i> spp.)	7 to 10	2	5.5	4.1	-	-	-
	7 to 21	2 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	-	-	-	4 to 5.5	11 to 15.1	8.25 to 11.3
Red thread (<i>Laetisana fuciformis</i>)	7 to 10	2 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	3.6 to 5.5	9.9 to 15.1	7.4 to 11.3	5.5	15.1	11.3
Stem rust (Bluegrass) (<i>Puccinia graminis</i>)	14	4 to 5.5	11.15.1	8.25 to 11.3	5.5	15.1	11.3
<p>*For Golf course fairways and roughs, lawns around commercial and industrial buildings, and professional and collegiate athletic fields: Limit to one application per year at rates greater than 7.3 lbs. a.i./Ac. (9.75 pts./Ac. or 3.6 fl. ozs./1,000 sq. ft.) of this product</p> <p>For Golf course greens and tees: Limit to two applications per year at rates greater than 7.3 lbs. a.i./Ac. (9.75 pts./Ac. or 3.6 fl. ozs./1,000 sq. ft.) of this product.</p> <p>**Low rate is not effective on intensively mowed Turfgrass such as Golf course greens and tees.</p>							

ALGAE

Golf Course Fairways and Roughs, Lawns Around Commercial and Industrial Buildings, and Professional and Collegiate Athletic Fields AND Golf Course Greens and Tees: For prevention of Algae on Turfgrasses, apply this product at the rate of 5.5 to 9.75 pints per acre (2 to 3.6 fl. ozs./1,000 sq. ft.) (4.1 to 7.3 lbs. a.i./Ac.) on a 7 to 14 day schedule. Under severe Algae conditions, use the 9.75 pints per acre (3.6 fl. ozs./1,000 sq. ft.) rate and apply on a 7 day schedule.

Make every attempt to dry out the afflicted area when Algae is well established. Once dry, spiking or verticutting should be done to enhance Turfgrass recovery in conjunction with an application of this product at the rate of 11 to 15.1 pints per acre (4 to 5.5 fl. ozs./1,000 sq. ft.).

On Golf Course Greens and Tees: A second application of this product at the rate of 15.1 pints per acre (5.5 fl. ozs. /1,000 sq. ft.) may be made 14 days after the first application.

On Golf Course Fairways and Roughs, Lawns Around Commercial and Industrial Buildings, and Professional and Collegiate Athletic Fields AND Golf Course Greens and Tees: Following application of the 15.1 pints per acre (5.5 fl. ozs. /1,000 sq. ft.) rate, several applications of this product at the rate of 5.5 to 9.75 pints per acre (2.0 to 3.6 fl. ozs. /1,000 sq. ft.) (4.1 to 7.3 lbs. a.i./Ac.) on a 7 to 14 day interval may be necessary for Turfgrass recovery. Only a preventive spray program with this product will prevent recurrence of the Algae when environmental conditions are favorable.

FUSARIUM (GERLACHIA) PATCH

Golf Course Fairways and Roughs, Lawns Around Commercial and Industrial Buildings, and Professional and Collegiate Athletic Fields AND Golf Course Greens and Tees: In areas where Pink snow mold (Gerlachia or Fusarium patch) is likely to occur, apply this product at 15.1 pints per acre (5.5 fl. ozs./1,000 sq. ft.) (11.3 lbs. a.i./Ac.) in combination with products containing iprodione at 88 ozs. a.i. per acre (2 ozs. a.i. /1,000 sq. ft.) of Turf area. It is the pesticide user's responsibility to ensure that the combination product is registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on the product label involved in tank-mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank-mixture.

For control of Fusarium patch only in areas where snow cover is intermittent or lacking during the Winter, apply 15.1 pints per acre (5.5 fl. ozs. /1,000 sq. ft.) of this product (11.3 lbs. a.i./Ac.). Make application in late Autumn.

Golf Course Greens and Tees: Apply a second application of 15.1 pints per acre (5.5 fl. ozs./1,000 sq. ft.) of this product 21 to 28 days after the first application unless conditions favorable for Fusarium patch no longer prevail.

GRAY SNOW MOLD CAUSED BY *TYPHULA* SPP.

Golf Course Fairways and Roughs, Lawns Around Commercial and Industrial Buildings, and Professional and Collegiate Athletic Fields AND Golf Course Greens and Tees: Apply in sufficient water to obtain adequate coverage (2 to 10 gals./1,000 sq. ft.). Apply one application of 15.1 pints per acre (5.5 fl. ozs. /1,000 sq. ft.) of this product (11.3 lbs. a.i./Ac.). Application must be made before snow cover in Autumn.

Golf Course Greens and Tees: If snow cover is intermittent or lacking during the Winter, a second application of this product at 15.1 pints per acre (5.5 fl. ozs./1,000 sq. ft.) may be applied one month after the first application.

Ornamental Plants

The following ornamental plants can be treated by this product:

Broadleaf Shrubs and Trees		
Andromeda (<i>Pieris</i>) ¹	Flowering Almond ^{2,3}	Oregon-Grape (<i>Mahonia</i>) ⁴
Ash (<i>Fraxinus</i>) ²	Flowering Cherry ^{2,3}	Photinia ²
Aspen ²	Flowering Peach ^{2,3}	Poplar ²
Azalea ^{1,2,3}	Flowering Plum ^{2,3}	Privet (<i>Ligustrum</i>) ²
Buckeye horsechestnut ²	Flowering Quince ^{2,3}	Rhododendron ^{1,2,3}
Cherry Laurel ²	Hawthorn ^{2,4}	Sand Cherry ^{2,3}
Crabapple ^{2,4,5}	Holly ²	Sequoia ²
Dogwood ²	Lilac ⁷	Spiraea ²
Eucalyptus ⁶	Magnolia ²	Sycamore planetree ²
Euonymus ²	Mountain Laurel ²	Viburnum ⁷
Firethorn (<i>Pyracantha</i>) ²	Oak (Red group only) ^{2,8}	Walnut (<i>Juglans</i>) ²
DISEASES CONTROLLED BY THIS PRODUCT:		
¹Phytophthora leaf blight/dieback		
²Leaf spots and Foliar Blights:		
Actinopelte leaf spot	Coryneum blight (Shothole)	Mycosphaerella ray blight
Alternaria leaf spot/leaf blight	Curvularia leaf spot	Myrothecium leaf spot/brown rot
Anthracnose leaf blotch/spot	Cylindrosporium leaf spot	Nematostoma leaf blight
Anthracnose blight (<i>Discula</i>)	Dactylaria leaf spot	Phyllosticta leaf spot
Ascochyta blight	Didymellina leaf spot	Ramularia leaf spot
Bipolaris leaf spot (<i>Helminthosporium</i>)	Drechslera leaf spot	Rhizoctonia web blight
Black spot on Roses	Fabraea leaf spot	Septoria leaf spot
Botrytis leaf spot/leaf blight	(<i>Entomosporium</i>)	Sphaeropsis leaf spot
Cephalosporium leaf spot	Gloeosporium black leaf spot	Stagonospora leaf scorch
Cercospora leaf spot	Ink spot (<i>Drechslera</i>)	Tan leaf spot (<i>Curvularia</i>)
Cercosporidium leaf spot	Marssonina leaf spot	Volutella leaf blight
Corynespora leaf spot	Monilinia blossom blight/twig blight	
³Flower Spots and Blight:		
Botrytis flower spot/flower blight		
Curvularia flower spot		
Monilinia blossom blight		
⁴Rusts:		
<i>Gymnosporangium</i> spp.		
<i>Pucciniastrum hydrangeae</i>		
<i>Puccinia</i> spp.		
⁵Scab		
⁶Cylindrocladium stem canker		
⁷Powdery mildew:		
<i>Erysiphe cichoracearum</i>		
<i>Microsphaera</i> spp.		
⁸Taphrina blister		

Flowering Plants* and Bulbs																																						
Arabian Violet ¹ Begonia ² Camellia ² Carnation ^{1,2} Chrysanthemum ^{1,2} Crocus ² Daffodil ² Daisy ² Geranium ^{2,3}	Gladiolus ^{1,2} Hollyhock ³ Hydrangea (Foliage only) ^{2,3} Iris ^{1,2} Lily ² Marigold ² Narcissus ² Pansy ² Petunia ^{2,5}	Phlox ² Pomsetta ^{2,**} Rose ^{2,***} Statice ² Tulip ² Zinnia ^{2,6}																																				
<p>DISEASES CONTROLLED BY THIS PRODUCT:</p> <p>¹Flower Spots and Blight: Botrytis flower spot/flower blight Curvularia flower spot Monilinia blossom blight</p> <p>²Leaf spots and Foliar Blights:</p> <table border="0"> <tr> <td>Actinopelte leaf spot</td> <td>Coryneum blight (Shothole)</td> <td>Mycosphaerella ray blight</td> </tr> <tr> <td>Alternaria leaf spot/leaf blight</td> <td>Curvularia leaf spot</td> <td>Myrothecium leaf spot/brown rot</td> </tr> <tr> <td>Anthrachnose leaf blotch/spot</td> <td>Cylindrosporium leaf spot</td> <td>Nematostoma leaf blight</td> </tr> <tr> <td>Anthrachnose blight (<i>Discula</i>)</td> <td>Dactylaria leaf spot</td> <td>Phyllosticta leaf spot</td> </tr> <tr> <td>Ascochyta blight</td> <td>Didymellina leaf spot</td> <td>Ramularia leaf spot</td> </tr> <tr> <td>Bipolaris leaf spot (<i>Helminthosporium</i>)</td> <td>Drechslera leaf spot</td> <td>Rhizoctonia web blight</td> </tr> <tr> <td>Black spot on Roses</td> <td>Fabraea leaf spot</td> <td>Septoria leaf spot</td> </tr> <tr> <td>Botrytis leaf spot/leaf blight</td> <td>(<i>Entomosporium</i>)</td> <td>Sphaeropsis leaf spot</td> </tr> <tr> <td>Cephalosporium leaf spot</td> <td>Gloeosporium black leaf spot</td> <td>Stagonospora leaf scorch</td> </tr> <tr> <td>Cercospora leaf spot</td> <td>Ink spot (<i>Drechslera</i>)</td> <td>Tan leaf spot (<i>Curvularia</i>)</td> </tr> <tr> <td>Cercosporidium leaf spot</td> <td>Marssonina leaf spot</td> <td>Volutella leaf blight</td> </tr> <tr> <td>Corynespora leaf spot</td> <td>Monilinia blossom blight/twig blight</td> <td></td> </tr> </table> <p>³Rusts: <i>Gymnosporangium</i> spp. <i>Pucciniastrum hydrangeae</i> <i>Puccinia</i> spp.</p> <p>⁴Cylindrocladium stem canker</p> <p>⁵Phytophthora leaf blight/dieback</p> <p>⁶Powdery mildew: <i>Erysiphe cichoracearum</i> <i>Microsphaera</i> spp.</p>			Actinopelte leaf spot	Coryneum blight (Shothole)	Mycosphaerella ray blight	Alternaria leaf spot/leaf blight	Curvularia leaf spot	Myrothecium leaf spot/brown rot	Anthrachnose leaf blotch/spot	Cylindrosporium leaf spot	Nematostoma leaf blight	Anthrachnose blight (<i>Discula</i>)	Dactylaria leaf spot	Phyllosticta leaf spot	Ascochyta blight	Didymellina leaf spot	Ramularia leaf spot	Bipolaris leaf spot (<i>Helminthosporium</i>)	Drechslera leaf spot	Rhizoctonia web blight	Black spot on Roses	Fabraea leaf spot	Septoria leaf spot	Botrytis leaf spot/leaf blight	(<i>Entomosporium</i>)	Sphaeropsis leaf spot	Cephalosporium leaf spot	Gloeosporium black leaf spot	Stagonospora leaf scorch	Cercospora leaf spot	Ink spot (<i>Drechslera</i>)	Tan leaf spot (<i>Curvularia</i>)	Cercosporidium leaf spot	Marssonina leaf spot	Volutella leaf blight	Corynespora leaf spot	Monilinia blossom blight/twig blight	
Actinopelte leaf spot	Coryneum blight (Shothole)	Mycosphaerella ray blight																																				
Alternaria leaf spot/leaf blight	Curvularia leaf spot	Myrothecium leaf spot/brown rot																																				
Anthrachnose leaf blotch/spot	Cylindrosporium leaf spot	Nematostoma leaf blight																																				
Anthrachnose blight (<i>Discula</i>)	Dactylaria leaf spot	Phyllosticta leaf spot																																				
Ascochyta blight	Didymellina leaf spot	Ramularia leaf spot																																				
Bipolaris leaf spot (<i>Helminthosporium</i>)	Drechslera leaf spot	Rhizoctonia web blight																																				
Black spot on Roses	Fabraea leaf spot	Septoria leaf spot																																				
Botrytis leaf spot/leaf blight	(<i>Entomosporium</i>)	Sphaeropsis leaf spot																																				
Cephalosporium leaf spot	Gloeosporium black leaf spot	Stagonospora leaf scorch																																				
Cercospora leaf spot	Ink spot (<i>Drechslera</i>)	Tan leaf spot (<i>Curvularia</i>)																																				
Cercosporidium leaf spot	Marssonina leaf spot	Volutella leaf blight																																				
Corynespora leaf spot	Monilinia blossom blight/twig blight																																					
<p>*Avoid applications during bloom period on plants where flower injury is unacceptable. **Discontinue applications prior to bract formation as phytotoxicity is possible on bracts. ***Use 1 pt. of this product per 100 gals. of water.</p>																																						

Foliage Plants		
Aglaonema ¹ Areca palm ¹ Artemesia ¹ Boston fern ¹ Dumbcane (<i>Dieffenbachia</i>) ¹ Dracaena ¹ Fatsia (<i>Aralia</i>) ¹	Ficus ¹ Florida ruffle fern ¹ Leatherleaf fern ¹ Lipstick plant ¹ Ming aralia ¹ Oyster plant (<i>Rhoeo</i>) ¹ Pachysandra ^{1,*}	Parlor palm (<i>Chamaedorea</i>) ¹ Peperomia ¹ Philodendron ^{1,2} Prayer plant (<i>Maranta</i>) ¹ Syngonium ¹ Zebra plant (<i>Aphelandra</i>) ¹
DISEASES CONTROLLED BY THIS PRODUCT:		
¹ Leaf spots and Foliar Blights:		
Actinopelte leaf spot	Coryneum blight (Shothole)	Mycosphaerella ray blight
Alternaria leaf spot/leaf blight	Curvularia leaf spot	Myrothecium leaf spot/brown rot
Anthraxnose leaf blotch/spot	Cylindrosporium leaf spot	Nematostoma leaf blight
Anthraxnose blight (<i>Discula</i>)	Dactylaria leaf spot	Phyllosticta leaf spot
Ascochyta blight	Didymellina leaf spot	Ramularia leaf spot
Bipolaris leaf spot (<i>Helminthosporium</i>)	Drechslera leaf spot	Rhizoctonia web blight
Black spot on Roses	Fabraea leaf spot	Septoria leaf spot
Botrytis leaf spot/leaf blight	(<i>Entomosporium</i>)	Sphaeropsis leaf spot
Cephalosporium leaf spot	Gloeosporium black leaf spot	Stagonospora leaf scorch
Cercospora leaf spot	Ink spot (<i>Drechslera</i>)	Tan leaf spot (<i>Curvularia</i>)
Cercosporidium leaf spot	Marssonina leaf spot	Volutella leaf blight
Corynespora leaf spot	Monilinia blossom blight/twig blight	
² Phytophthora leaf blight/dieback		
*Use 2.75 pts. of this product per 100 gals. of water.		

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container. Keep container tightly closed. This product can be stored at temperatures as low as minus 20°F. Keep away from heat and flame.

PESTICIDE DISPOSAL: To avoid waste, use all materials in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often, such programs are run by State or local governments or by industry).

CONTAINER HANDLING:

Nonrefillable Container (rigid material; less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container one-fourth full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid material; 5 gallons up to < 250 gallons): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container one-fourth full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Dispose of empty container in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Refillable Container (≥ 250 gallons & Bulk): Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

WARRANTY—CONDITIONS OF SALE

OUR DIRECTIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically directed and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable laws, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable laws, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

The DREXEL logo is a registered trademark of Drexel Chemical Company. All other brand names, product names or trademarks belong to their respective holders.