

50534-202

07-24-2009

1/39



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JUL 24 2009

Mr. Pat Dinnen
Product Registration
GB Biosciences Corp.
P.O. Box 18300
Greensboro, NC 27419

Subject: Label Notification for Pesticide Registration Notice 2007-4
(Storage & Disposal Changes)

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated June 22, 2009 for:

EPA Registration 50534-202 "Daconil SDG Fungicide"

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN 2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

PLEASE READ INSTRUCTIONS ON REVERSE BEFORE COMPLETING FORM.



United States
Environmental Protection Agency

Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Notification

Application for Pesticide - Section I

1. Company/Product Number 50534-202	2. EPA Product Manager Tony Kish	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Daconil SDG Fungicide	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) GB Biosciences Corporation P. O. Box 18300 Greensboro, NC 27419-8300 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: NOTIFICATION EPA Reg. No. _____ Product Name <u>JUL 24 2009</u>	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For Section I and Section II.).

Notification of label change per PR Notice 2007-4. This Notification is consistent with the guidance of PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

GB Biosciences Corporation is amending the Storage and Disposal section of the label by Notification according to the directions stated in PR Notice 2007-4.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
*Certification must be submitted			If "Yes" Unit Packaging wgt.	No. per Container	<input type="checkbox"/> Glass
			If "Yes" Unit Packaging wgt.	No. per container	<input type="checkbox"/> Paper
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 4 x 5 lb, 4 x 10 lb		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph	<input checked="" type="checkbox"/> Other	<u>Pressure Sensitive</u>
			<input type="checkbox"/> Paper glued		
			<input type="checkbox"/> Stenciled		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Pat Dinnen		Title Label Group Leader		Telephone No. (Include Area Code) 336-632-2494	
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.				6. Date Application Received (Stamped)	
2. Signature <i>Pat Dinnen</i>		3. Title Regulatory Specialist			
4. Typed Name Pat Dinnen		5. Date June 22, 2009			

MASTER

[Insert SuperWeatherStik® logo]

Daconil® SDG

Fungicide

For control of turf and ornamental diseases

Active Ingredient:
 Chlorothalonil (tetrachloroisophthalonitrile) 82.5%

Other Ingredients: 17.5%

Total 100.0%

(82.5% Water Dispersible Granules)

KEEP OUT OF REACH OF CHILDREN.

DANGER/PELIGRO


Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 50534-202

EPA Est.

Net Weight: _____



FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Rinse eye only with water. Do not put eye drops, drugs, or ointments in eyes unless specifically recommended by a medical doctor or a poison control center. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>NOTE TO PHYSICIAN Possible mucosal damage may contraindicate the use of gastric lavage; chemical adsorbents are recommended to reduce adsorption of the product. Persons suffering with temporary allergic skin reactions may respond to treatment with oral antihistamines and topical or oral steroids.</p> <p>If in eyes, the upper and lower lids should be retracted and irrigated, and any particulate matter should be carefully removed from the conjunctival fornix. Irrigation should be continued until the conjunctival sac is neutral on pH testing with universal indicator paper. Fluroscein staining is required to reveal the extent of corneal or conjunctival epithelial loss. Topical antibiotic ointments are indicated when corneal epithelial damage is identified. Use of steroid eye drops is not advocated unless expressly requested by an ophthalmologist.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	
<p>HOT LINE NUMBER For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals**DANGER/PELIGRO**

Corrosive. Causes irreversible eye damage. May be fatal if inhaled. Causes skin irritation. Do not get in eyes, on skin, or on clothing. Do not breathe dust. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Mixers, loaders, applicators and all other handlers must wear:

- coveralls over short-sleeved shirt and short pants
- chemical resistant gloves made of any waterproof material
- chemical resistant footwear plus socks
- protective eyewear
- chemical resistant headgear for overhead exposure
- chemical resistant apron when cleaning equipment, mixing, or loading
- dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C) or a NIOSH approved respirator with any N, R, P or HE filter
- For exposures in enclosed areas, such as a greenhouse, applicators and other handlers must wear a respirator with an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any N, R, P or HE prefilter.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This product is toxic to aquatic invertebrates and wildlife. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes towards adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of GB BIOSCIENCES CORPORATION or Seller. To the extent permitted by applicable law, Buyer and User agree to hold GB BIOSCIENCES and Seller harmless for any claims relating to such factors.

GB BIOSCIENCES warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or GB BIOSCIENCES, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, GB BIOSCIENCES MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.**

To the extent permitted by applicable law, in no event shall GB BIOSCIENCES be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF GB BIOSCIENCES AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF GB BIOSCIENCES OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

GB BIOSCIENCES and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of GB BIOSCIENCES.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Daconil SDG should be used only in accordance with recommendations on this label or in separately published GB Biosciences supplemental labeling recommendations for this product.

Agricultural Uses

For use to control diseases on turf in sod farms and commercial seed production farms and ornamentals in production operations such as farms, forests, nurseries, and greenhouses.

Do not apply this product in a way that will contact workers or other persons, or pets either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow workers to enter treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls over short-sleeved shirt and short pants
- chemical resistant gloves made of any waterproof material
- chemical resistant footwear plus socks
- protective eyewear
- chemical resistant headgear for overhead exposure

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted entry interval expires after 12 hours, for the next 6.5 days entry is permitted only when the following safety measures are provided:

- (1) At least one container designed specifically for flushing eyes must be available in

operating condition at the WPS required decontamination site intended for workers entering the treated area.

(2) Workers must be informed, in a manner they can understand:

- that residues in the treated area may be highly irritating to their eyes
- that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes
- that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water
- how to operate the eyeflush container

Non-Agricultural Uses

For use to control diseases on turf on golf courses, lawns around commercial and industrial buildings, and professional and collegiate athletic fields.

For use to control diseases on ornamentals on golf courses and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

GENERAL INFORMATION

Daconil SDG is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. Daconil SDG is recommended for use in programs which are compatible with the principles of Integrated Pest Management (IPM), which include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

Resistance Management

Daconil SDG is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. Some other fungicides which are at risk from disease resistance exhibit a single-site mode of fungicidal action. Daconil SDG, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your Federal or State Cooperative Extension Service

representatives for guidance on the proper use of Daconil SDG in programs which seek to minimize the occurrence of disease resistance to other fungicides.

Mixing Instructions

Do not combine Daconil SDG in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. Do not combine Daconil SDG with Dipel®, Latron B-1956® or Latron AG-98®, Chipco® Signature™, horticultural oil, and products containing xylene as phytotoxicity may result from the combination when applied to some species on this label.

The required amount of Daconil SDG should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of Daconil SDG in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

When tank mixing other products with Daconil SDG, follow the proper sequence of adding products to the spray tank. Wettable powders or water dispersible granules such as Daconil SDG should be added to the water in the tank first, followed by flowable products, and emulsifiable concentrates added last. Provide sufficient mechanical or bypass agitation during mixing and application.

When tank mixing, observe all directions, precautions, and limitations on labeling of all products used. Consult compatibility charts or your local or State agricultural authorities for compatibility information. It is impossible to test every species and variety of plants under all conditions.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (ie., elementary, middle and high schools), campgrounds, churches, and theme parks.

Agricultural Use Sites Only (sod farms, farms, forests, nurseries and greenhouses): This product must not be applied within 150 feet for aerial and airblast applications, or 25 feet for ground applications of marine/estuarine water bodies, unless there is an untreated buffer area of that width between the area to be treated and the water body.

Spray Drift Precautions

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information**.

Aerial Drift Reduction Advisory Information

This section is advisory in nature and does not supercede the mandatory label requirements.

Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable conditions (See Wind, Temperature).

Controlling Droplet Size

- **Volume** - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** - Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** - Orienting the nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift potential.

Boom Length

For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height

Applications should not be made at a height greater than 10 ft. above the top of the largest plants, unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the upwind and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

APPLICATION

Application and Calibration Techniques for Chemigation

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set or portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system. Do not use Daconil SDG through sprinkler irrigation equipment on golf courses.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

Always inject Daconil SDG into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Posting

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, daycare centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2½ inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

This sign is in addition to any sign posted to comply with the Worker Protection Standard.

Specific Chemigation Instructions

Daconil SDG may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Thoroughly mix recommended amount of Daconil SDG for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until Daconil SDG has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of Daconil SDG for acreage to be covered with water so that the total mixture of Daconil SDG plus water in the injection tank is equal to the quantity of water used during calibration, and operate entire system at normal pressures recommended by the manufacturer of injection equipment used, for amount of time established during calibration. Agitation is recommended. Daconil SDG can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until Daconil SDG has been cleared from last sprinkler head.

DIRECTIONS FOR APPLICATION

Turf

Always use Daconil SDG in conjunction with good turf management practices.

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (ie., elementary, middle and high schools), campgrounds, churches, and theme parks.

Sod Farms:

Do not use for sodfarms at application rates greater than 13 pounds of active ingredient, per acre, per year. Do not apply more than 15.8 lbs./acre (5.8 ozs./1000 sq. ft.) of Daconil SDG per growing season (13 lbs. a.i./acre) per growing season. The minimum re-treatment interval for single application rates **up to 8.8 lbs./acre** (3.2 ozs./1000 sq. ft.) of Daconil SDG (7.3 lbs. a.i./acre) is 7 days. Do not apply more than one application of a rate greater than 8.8 lbs./acre (3.2 ozs./1000 sq. ft.) of Daconil SDG (7.3 lbs. a.i./acre) per growing season. The maximum single application rate is 8.8 lbs./acre (3.2 ozs./1000 sq. ft.) of Daconil SDG (7.3 lbs. a.i./acre).

Apply Daconil SDG in 90 to 450 gallons of water per acre (2 to 10 gallons of water per 1,000 sq. ft.) for tees and greens and 30 to 450 gallons of water per acre (0.7 to 10 gallons of water per 1,000 sq. ft.) for all other turf. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions, use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

Diseases Controlled Pathogen(s)	Application Interval (days)	Pre-Disease Rates	Post-Disease Rates
		oz. product/ 1,000 sq. ft.	oz. product/ 1,000 sq. ft.
Dollar Spot <i>Sclerotinia homeocarpa</i>	7 to 10	1.0 ^a to 1.8	-
<i>Lanzia</i> spp.	7 to 21	1.8 to 3.25	-
<i>Moellerodiscus</i> spp.	14	-	3.7 to 5.0
Leaf Spot, Melting-Out, Brown Blight <i>Drechslera</i> spp.	7 to 10	1.8	-
(including <i>D. poae</i> , <i>D. siccans</i>)	7 to 21	1.8 to 3.25	-
<i>Bipolaris sorokiniana</i> .	14	-	3.7 to 5.0
<i>Curvularia</i> spp.			
Brown Patch <i>Rhizoctonia solani</i>	7 to 14	1.8 to 3.25	-
<i>R. zeae</i>	14	-	3.7 to 5.0
<i>R. cerealis</i>			
Gray Leaf Spot <i>Pyricularia grisea</i>	7 to 10	1.8 to 3.25	-
<i>P. oryzae</i>	14	-	3.7 to 5.0
Red Thread <i>Laetisaria fuciformis</i>	7 to 10	1.8 to 3.25	-
	14	3.25 to 5.0	5.0
Anthracnose <i>Colletotrichum graminicola</i>	7 to 14	2.75 to 3.25	-
	14	3.25 to 5.0	-
Copper Spot <i>Gloeocercospora sorghi</i>	14	3.7 to 5.0	5.0
Stem Rust <i>Puccinia graminis</i>	14	3.7 to 5.0	5.0
Dichondra Leaf Spot (CA only) <i>Alternaria</i> spp.	14	3.7 to 5.0	5.0
Gray Snow Mold ^b <i>Typhula</i> spp.	30 ^c	5.0	-
Pink Snow Mold ^d Fusarium Patch <i>Microdochium nivale</i>	21 to 28 ^e	5.0	-
Algae (algal scum)	7 to 14	1.8 to 5.0	5.0

^a Low rate is not effective on intensively mowed turfgrasses such as golf course tees and greens.

^b Apply before snow cover in autumn in 2 to 10 gallons of carrier per 1,000 sq. ft.

^c *Tees and Greens*: If snow cover is intermittent or lacking during the winter, you may make a second application.

^d Do not apply on top of snow. Apply in combination with Banner MAXX[®], Heritage[®], Medallion[®] or with products containing iprodione.

^e *Tees and Greens*: If conditions are favorable for *Fusarium*, you may make a second application.

Conversion Chart for Turf Rates

oz. product/ 1,000 sq. ft.	lbs. product/ acre	lbs. a.i./ acre
1.0	2.5	2.1
1.8	5.0	4.1
2.75	7.5	6.2
3.25	8.8	7.3
3.7	10.0	8.25
5.0	13.6	11.3

Turf Restrictions

- For rates **up to and including** 3.25 ozs./1,000 sq. ft.,
 - The minimum re-treatment interval is 7 days.
- For rates **greater than** 3.25 ozs./1,000 sq. ft.,
 - Limit of two applications per year on Tees and Greens.
 - The minimum re-treatment interval is 14 days.
 - Limit of one application per year on All Other Turf.
- Maximum single application rate of Daconil SDG: 5.0 ozs./1,000 sq. ft
- Maximum amount of Daconil SDG per growing season:

	oz./1,000 sq. ft.	lbs./acre	lbs. a.i./acre
Greens	32.5	88.5	73
Tees	23.0	63.0	52
Fairways and Roughs	11.6	31.5	26
All Other Turf	5.8	15.8	13

- Do not use Daconil SDG on fine fescue turf due to the potential for phytotoxicity or turfgrass injury.
- Do not mow or water after treatment until spray deposited on turfgrass is thoroughly dry.
- Sod farm turf treated with Daconil SDG prior to harvest must be mechanically cut, rolled and palletized.

Ornamental Plants

Daconil SDG may be used to control certain diseases of container, bench, flat, plug, bed or field-grown ornamentals in greenhouses, shade-houses, outdoor nurseries, and residential and commercial landscapes.

Apply Daconil SDG at a rate of 1.4 pounds (1.16 lbs. a.i.) per 100 gallons of water unless other directions are given in the tables below. Do not apply more than 44 pounds Daconil SDG (36.4 lbs. a.i.) per acre per growing season to field grown ornamentals.

Apply in a spray to run-off, when conditions are favorable for disease development. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable. During periods when conditions favor severe disease incidence, apply Daconil SDG at 7 day intervals. The minimum re-treatment interval is 7 days. Apply Daconil SDG when plants are dry or nearly dry.

Do not apply Daconil SDG through high pressure spray equipment. Do not use mistblowers, cold fog, or other fogging application equipment when making applications of Daconil SDG in greenhouses.

Do not eat edible parts from treated ornamental plants referred to in this Ornamental Plants section.

Table 1. Ornamentals recommended for treatment with Daconil SDG.

The numbers in parentheses refer to fungal diseases in Table 2 controlled by Daconil SDG.

Broadleaf Shrubs And Trees		
Andromeda (Pieris) (4)	Flowering Almond (1,2)	Oregon-Grape (Mahonia) (6)
Ash (Fraxinus) (1)	Flowering Cherry (1,2)	Photinia (1)
Aspen (1)	Flowering Peach (1,2)	Poplar (1)
Azalea (1,2,4)	Flowering Plum (1,2)	Privet (Ligustrum) (1)
Buckeye, Horsechestnut (1)	Flowering Quince (1,2)	Rhododendron (1,2,4)
Cherry-Laurel (1)	Hawthorn (1,6)	Sand Cherry (1,2)
Crabapple (1,6,8)	Holly (1)	Sequoia (1)
Dogwood (1)	Lilac (5)	Spiraea (1)
Eucalyptus (3)	Magnolia (1)	Sycamore, Planetree (1)
Euonymus (1)	Maple (1)	Viburnum (5)
Firethorn (Pyracantha) (1)	Mountain Laurel (1)	Walnut (Juglans) (1)
	Oak (red group only) (1,7)	
Flowering Plants^a, Bulbs and Corms		
Arabian Violet (2)	Geranium (1,6)	Narcissus (1)
Begonia (1)	Gladiolus (1,2)	Pansy (1)
Caladium (1)	Hollyhock (6)	Petunia (1,4)
Camellia (2)	Hydrangea (foliage only) (1,6)	Phlox (1)
Carnation (1,2)	Iris (1,2)	Poinsettia ^b (1)
Chrysanthemum (1,2)	Iris, Bulbous (1)	Rose ^c (1)
Crocus (1)	Lily (1)	Statice (1)
Daffodil (1)	Lily, Asiatic (1)	Tulip (1)
Daisy (1)	Marigold (1)	Zinnia (1,5)
Foliage Plants		
Aglaonema (1)	Ficus (1)	Peperomia (1)
Areca Palm (1)	Lipstick Plant (1)	Philodendron (1,4)
Artemesia (1)	Ming Aralia (1)	Prayer Plant (Maranta) (1)
Dumbcane (Dffenbachia) (1)	Oyster Plant (Rhoeo) (1)	Syngonium (1)
Dracaena (1)	Pachysandra ^d (1)	Zebra Plant (Aphelandra) (1)
Fatsia (Aralia) (1)	Parlor Palm (Chamaedorea) (1)	

^aAvoid applications during bloom period on plants where flower injury is unacceptable.

^bDiscontinue applications prior to bract formation; phytotoxicity is possible on the bracts.

^cUse 1 pound Daconil SDG (0.825 lbs. a.i.) per 100 gallons of water.

^dUse 2.5 pounds of Daconil SDG (2.1 lbs. a.i.) per 100 gallons of water.

Table 2. Diseases Controlled with Daconil SDG

1. Leaf Spots/Foliar Blights		
Actinopelte leaf spot	Coryneum blight (shothole)	Mycosphaerella ray blight
Alternaria leaf spot/leaf blight	Curvularia leaf spot	Myrothecium leaf spot, brown rot
Anthracnose leaf blotch, spot	Cylindrosporium leaf spot	Nematostoma leaf blight
Anthracnose (Discula) blight	Dactylaria leaf spot	Phyllosticta leaf spot
Ascochyta blight	Didymellina leaf spot	Ramularia leaf spot
Bipolaris (Helminthosporium) leaf spot	Drechslera leaf spot	Septoria leaf spot, Rhizoctonia web blight
Black spot on roses	Fabraea (Entomosporium) leaf spot	Sphaeropsis leaf spot
Botrytis leaf spot, leaf blight	Fusarium leaf spot	Stagonospora leaf scorch
Cephalosporium leaf spot	Gloeosporium black leaf spot	Tan leaf spot (Curvularia)
Cercospora leaf spot	Ink spot (Drechslera)	Volutella leaf blight
Cercosporidium leaf spot	Marssonina leaf spot	
Corynespora leaf spot	Monilinia blossom blight, twig blight	
2. Flower spots/blights		
Botrytis flower spot, flower blight	Monilinia blossom blight	Rhizopus blossom blight
Curvularia flower spot	Ovulinia flower blight	Sclerotinia flower blight
3. Cylindrocladium stem canker		
4. Phytophthora leaf blight, dieback		
5. Powdery mildews		
<i>Erysiphe cichoracearum</i>	<i>Microsphaera</i> spp.	
6. Rusts		
<i>Gymnosporangium</i> spp.	<i>Pucciniastrum hydrangeae</i>	<i>Puccinia</i> spp.
7. Taphrina blister		
8. Scab (<i>Venturia inaequalis</i>)		

Plant Safety

Daconil SDG has been shown to be safe when applied at the recommended rates to the ornamental plants listed in the following tables. However, due to the large number of genera, species and varieties of ornamental and nursery plants, it is impossible to test every one for tolerance to Daconil SDG. Neither the manufacturer nor the seller has determined whether Daconil SDG can be used safely on genera, species, or varieties of ornamental and nursery plants not specified on this label. The professional user should conduct small scale testing at the recommended rates to ensure plant safety prior to broad scale commercial use on plant genera and species not listed in this label. Applications made during bloom may damage flowers and/or fruits.

Do not apply Daconil SDG to either green or variegated Pittosporum or to Schefflera, as multiple applications may cause phytotoxic responses.

Do not apply Daconil SDG to ferns.

Table 3. The following ornamental plant species that have been tested with Daconil SDG at recommended rates did not exhibit phytotoxicity.

Common Name	
Aechmea	Golden Pothos, Scindapsus
Aluminum Plant	Impatiens
Aster	Jade Plant
Baby's Breath	Japanese Holly
Birdsnest Sansevieria	Natal Plum
Bleeding Heart	Norfolk Island Pine
Bougainvillea	Peacock Plant
Caladium	Piggy-back Plant
Chinese Holly	Purple Passion Vine
Christmas Cactus	Silver-nerve Plant
Croton	Spineless Yucca
False Aralia	Ti Plant
Flame Violet	Venus Fly Trap
Gerbera Daisy	Wax Plant

Bulb and Corm Dip

Daconil SDG may be used to control bulb and corm diseases of ornamental flowering plants.

CROP	DISEASES	Lbs. Product per 100 Gal.*	APPLICATION DIRECTIONS
Caladium Crocus Daffodils Iris Lily (bulb) Tulips	Basal Rot Neck Rot Other bulb rot diseases caused by: <i>Mucor</i> spp. <i>Zygorrhynchus</i> spp. <i>Rhizopus</i> spp. <i>Curvularia</i> spp. <i>Rhizoctonia</i> spp. <i>Fusarium oxysporum</i>	2.5 to 5.0	Dip bulbs from 15 minutes up to 4 hours prior to planting. Add the recommended diluted mixture of Daconil SDG to the dip tanks to maintain dip solution at levels needed to achieve complete bulb coverage.
Gladiolus	<i>Botrytis</i> spp. <i>Curvularia</i> spp.	2.5	Recharge dip tanks with 0.3 lbs. product per 20,000 corms. Treat corms once before storage and once before planting. Allow to drain and dry before handling corms.

Application of Dip Tank Treatment Water: Spent dip tank treatment water may be applied using ground equipment to bulb fields for basal, neck or other bulb rots.

Conifers

Apply Daconil SDG in sufficient water and with proper calibration to obtain uniform coverage of tree canopy.

Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, Daconil SDG may be applied with aircraft. The minimum volume for application by aircraft to forest nurseries, forest stands, and Christmas trees is 10 gallons per acre. The minimum volume for application by ground equipment to conifer nursery beds is 5 gallons per acre. For conifers, the maximum volume is 100 gallons per acre.

When concentrate sprays are used or when treating immature trees, the lower rate of Daconil SDG listed may be used. Do not allow livestock to graze in treated areas.

Do not apply Daconil SDG through high pressure spray equipment. Do not use mistblowers, cold fog, or other fogging application equipment when making applications of Daconil SDG in greenhouses.

Do not use on blue spruce.

CROP	DISEASES (Pathogen)	Lbs. Product Per (lbs. a.i. per)		APPLICATION DIRECTIONS
		Acre	100 Gal.*	
Conifers (pines, spruce, Douglas fir)	Swiss needlecast (<i>Phaeocryptopus gaeumannii</i>)	2.5 to 5.0 (2.1 to 4.1)	2.5 to 5.0 (2.1 to 4.1)	Single application technique: In Christmas plantations or forest stands, make one application in the spring when new shoot growth is ½ to 2 inches in length.
	Scleroderris canker (pines) (<i>Gremmeniella abietina</i>)	1.25 to 2.5 (1.0 to 2.1)	1.25 to 2.5 (1.0 to 2.1)	Make the first application in spring when new shoot growth is ½ to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
	Swiss needlecast (<i>P. gaeumannii</i>)			
	Sirococcus tip blight (<i>S. conigenus</i>)	1.8 to 3.2 (1.5 to 2.6)	1.8 to 3.2 (1.5 to 2.6)	
	Rhizosphaera needlecast (spruces) (<i>Rhizosphaera spp.</i>)	5.0 (4.1)	5.0 (4.1)	
	Scirrhia brown spot (pines) (<i>Mycosphaerella dearnessii</i>)			
	Cyclaneusma and Lophodermium needlecasts (pines)	2.5 to 5.0 (2.1 to 4.1)	2.5 to 5.0 (2.1 to 4.1)	Apply in early spring prior to budbreak. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.
	Rhabdocline needlecast (Douglas-fir)	1.4 to 2.5 (1.1 to 2.1)	1.4 to 2.5 (1.1 to 2.1)	Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3 week schedule.
	Botrytis seedling blight	1.4 to 2.5 (1.1 to 2.1)	1.4 to 2.5 (1.1 to 2.1)	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.
Phoma twig blight				

CROP	DISEASES (Pathogen)	Lbs. Product Per (lbs. a.i. per)		APPLICATION DIRECTIONS
		Acre	100 Gal.*	
	Autoecious needle rust (Weir's cushion) (spruce)	5.0 (4.1)	5.0 (4.1)	Begin applications when 10% of buds have broken and twice thereafter at 7 to 10 day intervals.

Specific Use Restrictions:

- Do not apply more than 20 pounds Daconil SDG (16.5 lbs. a.i.) per acre during each growing season.
- The minimum re-treatment interval for established trees is 21 days.
- The minimum re-treatment interval in nursery beds is 7 days.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

Nurseries, Gardens and Landscapes

Daconil SDG may be applied by ground application to plants in production nurseries, gardens, and landscapes to control certain diseases. Follow pre-harvest interval following applications prior to consuming fruits, nuts, or other produce from the treated areas.

Fruit and Nut Trees

Apply Daconil SDG in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. For fruit and nut trees, the maximum volume is 300 gallons per acre unless indicated otherwise in the specific use directions. Apply by ground equipment only.

Do not allow livestock to graze in treated areas.

CROP	DISEASES (Pathogen)	Lbs. Product Per (lbs. a.i. per)		APPLICATION DIRECTIONS
		Acre	100 Gal.*	
Almonds	Blossom blight/brown rot (<i>Monilinia</i> spp.)	3.6 (3.0)	1.2 (1.0)	Use water volumes of 20 to 300 gallons per acre. For blossom blight, begin application at popcorn (pink bud) and follow with an application at full bloom. If weather is still conducive for disease development, another application may be made at petal fall. For control of shothole, make an application in the autumn at leaf fall. In the spring, make the first application at budbreak, followed by an application at shuck split to control nut infections and to control scab.
	Shot hole (<i>Wilsonomyces carpophilus</i>)			
	Scab (<i>Venturia carpophila</i>)			

Specific Use Restrictions:

- Do not apply more than 22.8 pounds Daconil SDG (18.75 lbs. a.i.) per acre during each growing season (leaf fall through shuck split).
- Do not apply within 150 days of harvest.

Filberts (Hazelnuts)	Eastern filbert blight (<i>Anisogramma anomala</i>)	3.6 (3.0)	1.2 (1.0)	Use a water volume of 20 to 300 gallons per acre. Begin applications at the onset of disease or when weather conditions favor disease development. Make applications on a 14 to 28 day schedule, using the shorter interval under heavy disease pressure (the minimum re-treatment interval is 14 days).
-------------------------	--	--------------	--------------	--

Specific Use Restrictions:

- Do not apply more than 11 pounds Daconil SDG (9 lbs. a.i.) per acre during each growing season.
- Do not apply within 120 days of harvest.
- Do not apply through irrigation.
- Do not apply with oils, other pesticides, surfactants or fertilizers.
- Do not apply within one week of an oil-based pesticide application.

Peach Nectarine Apricot Cherry Plum Prune	Leaf curl (<i>Taphrina deformans</i>)	2.8 to 3.8 (2.3 to 3.1)	0.9 to 1.25 (0.75 to 1.0)	For best control of both diseases, apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels, use the high rate of application and apply once or twice more in mid to late winter before budswell. If the leaf fall application is not practical, application of Daconil SDG for control of leaf curl may be made at any time prior to budswell the following spring. Where shothole occurs, also apply at budbreak to protect newly emerging leaves and at shuck split to prevent fruit infections.
	Shot hole (<i>Wilsonomyces carpophilus</i>)			
	Lacy (russet) scab (plum/prune)	2.8 to 3.8 (2.3 to 3.1)	0.9 to 1.25 (0.75 to 1.0)	Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Brown rot blossom blight <i>Monilinia</i> spp.			

CROP	DISEASES (Pathogen)	Lbs. Product Per (lbs. a.i. per)		APPLICATION DIRECTIONS
		Acre	100 Gal.*	
	Cherry leaf spot (<i>Blumeriella jaapii</i>)	2.8 to 3.8 (2.3 to 3.1)	0.9 to 1.25 (0.75 to 1.0)	In addition to the bloom application listed above, make one application at shuck split. Do not apply Daconil SDG after shuck split and before harvest. If additional disease control is needed before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10 to 14 days later.
	Scab (<i>Cladosporium carpophilum</i>)			
	Black knot (cherry, plum) (<i>Apiosporina morbosa</i>)			

Specific Use Restrictions:

- Do not apply more than 18.8 pounds of Daconil SDG (15.5 lbs. a.i.) per acre during each growing season.
- Daconil SDG may be applied the day of harvest.
- The minimum re-treatment interval is 10 days.

Pistachio	Botryosphaeria blight (<i>B. dothidea</i>)	5.4 (4.5)	2.7 (2.2)	Use a water volume of 20 to 200 gallons per acre. Make the first application at the beginning of the blossom period followed by an application at full bloom. Make additional applications as required on a 28-day schedule (the minimum re-treatment interval is 28 days). For Septoria and Botrytis, use the higher rate if disease pressure is severe.
	Suppression: Alternaria late blight (<i>A. alternata</i>)			
	Septoria leaf spot (<i>S. pistacina</i>)	3.6 to 5.4 (3.0 to 4.5)	1.8 to 2.7 (1.5 to 2.2)	NOTE: Use of this product may result in speckling or reddening of the fruit hull (epicarp). This effect is superficial and has not resulted in any change in nut quality.
	Botrytis blight (<i>B. cinerea</i>)			

Specific Use Restrictions:

- Do not apply more than 27 pounds Daconil SDG (22.5 lbs. a.i.) per season.
- Do not apply within 14 days of harvest.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

Vegetables and Fruits

Apply Daconil SDG in sufficient water and with proper calibration to obtain uniform coverage. For vegetables and fruits, the maximum volume is 150 gallons per acre unless indicated otherwise in the specific use directions. Apply by ground equipment only.

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Asparagus	Rust (<i>Puccinia asparagi</i>)	1.8 to 3.6 (1.5 to 3.0)	Use water volumes of 25 to 50 gallons per acre. Begin applications following final harvest of spears. Repeat applications at 14 to 28 day intervals (the minimum re-treatment interval is 14 days), depending on disease pressure. Use the higher rate and shorter interval if disease severity begins to increase during the season or weather conditions are conducive for severe epidemics.
	Purple Spot (<i>Pleospora herbarum</i>)		
	Cercospora blight (<i>C. asparagi</i>)		

- Specific Use Restrictions:**
- Do not apply more than 11 pounds Daconil SDG (9.0 lbs. a.i.) per acre during each growing season.
 - Do not apply within 190 days (120 days in CA and AZ) of the harvest of spears in the following season.

Bean (Snap)	Rust (<i>Uromyces appendiculatus</i>)	1.25 to 2.7 (1.0 to 2.2)	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat as necessary (the minimum re-treatment interval is 7 days) to maintain control.
	Botrytis blight (gray mold) (<i>B. cinerea</i>)	2.7 (2.2)	

- Specific Use Restrictions:**
- Do not apply more than 10.9 pounds of Daconil SDG (9.0 lbs. a.i.) per acre during each growing season.
 - Do not apply within 7 days of harvest.

Beans (Dry) (except soybeans) bean, adzuki bean, broad bean, dry bean, lablab bean, navy bean, kidney bean, lima bean, moth bean, mung bean, pink bean, pinto bean, tepary bean, urd bean, yardlong catjang chickpea (garbanzo) cowpea lupin, grain lupin bean, rice bean, runner bean, jackbean pea, blackeyed pea, southern	Rust (<i>Uromyces appendiculatus</i>)	1.25 to 1.8 (1.0 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications at first onset of disease, which may occur as early as 2 to 4 weeks before flowering. Repeat applications at 7 to 10 day intervals (the minimum re-treatment interval is 7 days). For use only on beans to be harvested dry with pods removed.
	Anthracnose (<i>Colletotrichum lindemuthianum</i>)		
	Downy mildew (<i>Phytophthora nicotianae</i>)		
	Cercospora leaf blotch (<i>C. cruenta</i>)		
	Ascochyta blight (<i>A. phaseolorum</i>)		

- Specific Use Restrictions:**

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
<ul style="list-style-type: none"> Do not apply more than 7.3 pounds of Daconil SDG (6 lbs. a.i.) per acre during each growing season. Do not apply within 14 days before harvest. 			
Blueberries	Suppression: Anthracnose (ripe rot) <i>(C. gloeosporoides)</i> Mummy Berry <i>(M. vaccinicorymbos)</i>	2.7 to 3.6 (2.2 to 3.0)	Daconil SDG should be integrated into an overall disease management strategy which includes alternation with a fungicide with a different mode of action. Diseases may only be suppressed and russetting may occur under heavy disease pressure or unfavorable environmental conditions. Apply in sufficient water to obtain adequate coverage, normally 20 to 100 gallons per acre. Begin applications at budbreak (green tip) and repeat at 10-day intervals through early bloom (the minimum re-treatment interval is 10 days). Under heavy disease pressure, use the higher rate.
	Septoria leaf spot <i>(Septoria albopunctata)</i> Rust <i>(Pucciniastrum vaccinii)</i>	3 to 4 (2.25 to 3.0)	Foliar Use After Harvest (after all berries are harvested): To maintain healthy leaves for the following season, apply in sufficient water to obtain adequate coverage (normally 20 to 100 gallons per acre). Repeat at 10 to 14 day intervals (the minimum re-treatment interval is 10 days).
Specific Use Restrictions: <ul style="list-style-type: none"> Do not apply more than 10.9 pounds Daconil SDG (9.0 lbs. a.i.) per acre during each growing season. Do not apply after full bloom (except for foliar use after harvest) or within 42 days of harvest. 			
Cabbage Chinese Cabbage (tight-headed varieties only) Cauliflower Broccoli Chinese Broccoli Brussels Sprouts	Alternaria leaf spot <i>(Alternaria spp.)</i> Downy mildew <i>(Peronospora parasitica)</i>	1.4 (1.2)	Use in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control.
	Ring spot (California only)	1.4 (1.2)	For field-seeded brussels sprouts, begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control.
Specific Use Restrictions: <ul style="list-style-type: none"> Do not apply more than 14.5 pounds Daconil SDG (12 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest. 			
Carrot	Cercospora leaf spot <i>(C. carotae)</i> Alternaria leaf blight <i>(A. dauci)</i>	1.4 to 1.8 (1.2 to 1.5)	Use in sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals (the minimum re-treatment interval is 7 days) to maintain control.
Specific Use Restrictions: <ul style="list-style-type: none"> Do not apply more than 18.2 pounds of Daconil SDG (15 lbs. a.i.) per acre during each growing season. Daconil SDG may be applied the day of harvest. 			

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Celery	Early blight (<i>Cercospora apii</i>)	1.8 to 2.7 (1.5 to 2.2)	Use in sufficient water to obtain adequate coverage. Start applications when transplants are set in the field and repeat at a 7 day interval as needed to maintain control (the minimum re-treatment interval is 7 days).
	Late blight (<i>Septoria apicola</i>)		
	Basal stalk rot (<i>Rhizoctonia solani</i>)		
	Suppression (7 day schedule): Pink rot (<i>Sclerotinia sclerotiorum</i>)	2.7 (2.2)	
	Early blight (<i>Cercospora apii</i>) Late blight (<i>Septoria apicola</i>)	1.4 to 1.8 lbs. per 100 gal. (1.2 to 1.5 lbs. a.i. per 100 gal.)	For celery seedbeds, apply in a spray volume of 125 gallons per acre twice weekly or as needed to maintain control. Start applications shortly after crop emergence. Use the higher rate under severe disease conditions.

Specific Use Restrictions:

- Do not apply more than 21.8 pounds Daconil SDG (18 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

Corn (sweet)	Helminthosporium leaf blights	0.7 to 1.8 (0.6 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at a 7 day interval as required to maintain control (the minimum re-treatment interval is 7 days). Under severe disease conditions, use 1.4 to 1.8 pounds Daconil SDG per acre.
	Rust (<i>Puccinia</i> spp.)		

Specific Use Restrictions:

- Do not apply more than 10.9 pounds Daconil SDG (9 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest.
- Do not apply to sweet corn to be processed.
- Do not allow livestock to graze in treated fields.
- Do not ensile treated corn or use as livestock forage.

Cranberry	Fruit rots	3.8 to 6.0 (3.1 to 5.0)	Apply at early bloom and repeat at 10 to 14 day intervals (the minimum re-treatment interval is 10 days). Under severe disease conditions, use 6.0 pounds Daconil SDG (5.0 lbs. a.i.) per acre rate on a 10 day schedule.
	Lophodermium leaf/twig blight (<i>L. hypophyllum</i>)		
	Upright Dieback (<i>Phomopsis vaccinii</i>)	3.8 to 6.0 (3.1 to 5.0)	Apply in sufficient water to obtain coverage of uprights and runners. Make the first application before bloom, at the time shoots begin growth in the spring. Make additional applications at 10-14 day intervals.

Specific Use Restrictions:

- Do not apply more than 18.2 pounds Daconil SDG (15 lbs. a.i.) per acre during each growing season.
- Do not apply within 50 days before harvest.
- Do not apply to beds when flooded or allow release of irrigation water from beds for at least 3 days following application.

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Cucurbits Cucumber Cantaloupe Muskmelon Honeydew melon Watermelon Squash Pumpkin	Target spot (<i>Corynespora cassicola</i>)	1.4 to 1.8 (1.2 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals (the minimum re-treatment interval is 7 days).
	Anthrachnose (<i>Colletotrichum</i> spp.)		
	Downy mildew (<i>Pseudoperonospora cubensis</i>)	1.8 to 2.7 (1.5 to 2.2)	<p>Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply Daconil SDG to watermelons when any of the following conditions are present:</p> <ol style="list-style-type: none"> 1. Intense heat and sunlight 2. Drought conditions 3. Poor vine canopy 4. Other crop and environmental conditions which may be conducive to increased natural sunburn. <p>Do not combine Daconil SDG with anything except water for application to watermelons unless prior use has shown the combination to be non-injurious to watermelons under your conditions of use.</p>
	Cercospora leaf spot (<i>C. citrullina</i>)		
	Gummy stem blight/ vine decline (<i>Didymella bryoniae</i>)		
	Alternaria leaf blight (<i>A. cucumerina</i>)		
Alternaria leaf spot (<i>A. alternata</i>)			
Scab (<i>Cladosporium cucumerinum</i>)	1.2 (1.0)	Use in sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground applications. Begin applications when emerging plants are 4 to 8 inches high. Repeat applications at 7 to 10 day intervals to maintain control (the minimum re-treatment interval is 7 days).	
Powdery mildew (<i>Sphaerotheca</i> only)			
Specific Use Restrictions:			
<ul style="list-style-type: none"> ▪ Do not apply more than 19.1 pounds Daconil SDG (15.75 lbs. a.i.) per acre during each growing season. ▪ Daconil SDG may be applied the day of harvest. 			
Mango	Anthrachnose (<i>Colletotrichum</i> spp.)	1.8 to 3.1 (1.5 to 2.6)	Use a water volume of 20 to 300 gallons per acre. Begin applications at early bloom and repeat on a 7 to 14 day interval until early fruit development. Begin the season with the 1.8 lb. rate on a 14-day interval (the minimum re-treatment interval is 7 days). If disease pressure is severe, use the higher rate and shorter interval.
Specific Use Restrictions:			
<ul style="list-style-type: none"> ▪ Do not apply more than 29 pounds Daconil SDG (24.0 lbs. a.i.) per acre during each growing season. ▪ Do not apply within 21 days of harvest. 			
Mint (Indiana, Michigan and Wisconsin only)	Rust (<i>Puccinia menthae</i>) Septoria leaf spot (<i>S. menthae</i>)	1.2 (1.0)	Use in sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground applications. Begin applications when emerging plants are 4 to 8 inches high. Repeat applications at 7 to 10 day intervals to maintain control (the minimum re-treatment interval is 7 days).

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
------	------------------------	---------------------------------	------------------------

Specific Use Restrictions:

- Do not apply more than 3.6 pounds Daconil SDG (3 lbs. a.i.) per acre during each growing season.
- Do not apply within 80 days of harvest.
- Do not feed fresh or extracted mint hay from treated fields to livestock.

Onion (dry bulb) and Garlic	Botrytis leaf blight (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) Suppression: Botrytis neck rot Downy mildew (<i>Peronospora destructor</i>)	0.9 to 2.7 (0.75 to 2.2)	Use in sufficient water to obtain adequate coverage of tops. Daconil SDG is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply Daconil SDG as follows:			
				Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard
			Rate per Acre	0.9 lbs.	1.25 lbs.	2.7 lbs.
			Frequency	10 days	7 to 10 days	7 days
			For suppression of neck rot (<i>Botrytis</i> spp.) during storage, a minimum of three weekly applications prior to lifting, using 1.25 to 2.7 lbs. of Daconil SDG per acre, is recommended. The minimum re-treatment interval is 7 days.			

Specific Use Restrictions:

- Do not apply more than 18.2 pounds Daconil SDG (15 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

Onion (green bunching) Leek Shallots Onion and Garlic (grown for seed)	Botrytis leaf blight (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) Suppression: Downy mildew (<i>Peronospora destructor</i>)	1.4 to 2.7 (1.2 to 2.2)	Use in sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods, and repeat at 7 to 10 day intervals for as long as conditions favor disease (the minimum re-treatment interval is 7 days). Use the high rate and a 7 day schedule of applications when heavy dew or rain persist.
--	---	----------------------------	---

Specific Use Restrictions:

- Do not apply more than 8.2 pounds Daconil SDG (6.75 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest on garlic.
- Do not apply within 14 days of harvest on green bunching onions, leeks or shallots.

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Papaya	Alternaria fruit spot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum</i> spp.) Stem end rot (<i>A. alternata</i> , <i>Colletotrichum</i> spp.)	1.4 to 2.7 (1.2 to 2.2)	Apply with ground equipment only, in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days).

Specific Use Restrictions:

- Do not apply more than 8.2 pounds Daconil SDG (6.75 lbs. a.i.) per acre during each growing season.
- Daconil SDG may be applied the day of harvest.

Parsnip	Alternaria leaf spot (<i>Alternaria</i> spp.) Downy mildew (<i>Plasmopara</i> <i>crustosa</i>) Anthracnose (<i>Colletotrichum</i> spp.) Botrytis blight (gray mold) (<i>B. cinerea</i>) Bottom rot (<i>Rhizoctonia</i>)	1.4 to 1.8 (1.2 to 1.5)	Apply in sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule (the minimum re-treatment interval is 7 days).
---------	--	----------------------------	---

Specific Use Restrictions:

- Do not apply more than 7.3 pounds of Daconil SDG (6 lbs. a.i.) per acre during each growing season.
- Do not apply within 10 days of harvest.

Passion Fruit (Hawaii only)	Alternaria fruit and leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Cercospora fruit spot	1.8 (1.5)	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin applications during late bloom and repeat at 14 day intervals until weather conditions no longer favor disease development (the minimum re-treatment interval is 14 days.)
--------------------------------	---	--------------	---

Specific Use Restrictions:

- Do not apply more than 9.1 pounds Daconil SDG (7.5 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
Peanut	Early leaf spot (<i>Cercospora arachidicola</i>)	0.9 to 1.36 (0.75 to 1.1)	Apply in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting; repeat at 14 day intervals (the minimum re-treatment interval is 14 days). When conditions favor late leaf spot or when rust or web blotch occur, apply 1.36 pounds Daconil SDG per acre at 14 day intervals for the remainder of the season.
	Late leaf spot (<i>Cercosporidium personatum</i>)		
	Pepper Spot (<i>Leptosphaerulina crassiasca</i>)		
	Rust (<i>Puccinia arachidis</i>)	1.36 (1.1)	
Potato	Late blight (<i>Phytophthora infestans</i>)	0.7 (0.6)	Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum re-treatment interval is 5 days).
	Early blight (<i>Alternaria solani</i>)	-then-	
	Botrytis vine rot (<i>B. cinerea</i>)	0.9 to 1.36 (0.75 to 1.1)	Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occur:
	Black dot (<i>Colletotrichum coccodes</i>)		

Specific Use Restrictions:

- Do not apply more than 10.9 pounds Daconil SDG (9 lbs. a.i.) per acre during each growing season.
- Do not apply within 14 days of harvest.
- Do not allow livestock to graze in treated areas.
- Do not feed hay or threshings from treated fields to livestock.

Specific Use Restrictions:

- Do not apply more than 13.6 pounds of Daconil SDG (11.25 lbs. a.i.) per acre during each growing season.
- Do not apply within 7 days of harvest.

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
------	------------------------	---------------------------------	------------------------

Tomato

<p>FOLIAGE</p> <p>Early blight (<i>Alternaria solani</i>)</p> <p>Late blight (<i>Phytophthora infestans</i>)</p> <p>Gray leaf spot (<i>Stemphyllium botryosum</i>)</p> <p>Gray leaf mold (<i>Fluvia fluva</i>; <i>Cladosporium</i>)</p> <p>Septoria leaf spot (<i>S. lycopersici</i>)</p> <p>Target spot (<i>Corynespora cassicola</i>)</p>	<p>1.3 to 1.8 (1.1 to 1.5)</p>
--	------------------------------------

Apply in sufficient water to obtain adequate coverage. Begin applications when dew or rain occur and disease threatens. Apply on a 7 to 10 day interval for foliage diseases. For fruit diseases, begin at fruit set and apply on a 7 to 14 day interval. Use the highest rate and shortest interval specified when disease conditions are severe. The minimum re-treatment interval is 7 days.

CROP	DISEASES (Pathogen)	Lbs. Product/A (lbs. a.i./A)	APPLICATION DIRECTIONS
	FRUIT Anthracnose (<i>Colletotrichum</i> spp.) Alternaria fruit rot (black mold) (<i>A. alternata</i>) Botrytis gray mold (<i>B. cinerea</i>) Late blight fruit rot (<i>P. infestans</i>) Rhizoctonia fruit rot (<i>R. solani</i>)	1.8 to 2.6 (1.5 to 2.2)	

Specific Use Restrictions:

- Do not apply more than 18.3 pounds Daconil SDG (15.1 lbs. a.i.) per acre during each growing season.
- Daconil SDG may be applied the day of harvest.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage

Store in a dry place.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Daconil®, Daconil Ultrex®, Turf Care®, SuperWeatherStik®, Banner MAXX®, Heritage®, and Medallion® are trademarks of a Syngenta Group Company

Chipco® and Signature™ are trademarks of Bayer

Dipel® is a registered trademark of Valent BioSciences Corporation.

Latron B-1956® and Latron AG-98® are trademarks of Dow AgroSciences LLC.

© 200_ Syngenta

For non-emergency (e.g., current product information) call
Syngenta Crop Protection at 1-800-334-9481.

Manufactured for :
GB Biosciences Corporation
P.O. Box 18300
Greensboro, North Carolina 27419-8300
www.syngenta-us.com